Parameter estimation of the combined SDLINARmodel of order p and further sideways

Miodrag S. Djordjević¹, Aleksandar S. Nastić¹, and Miroslav M. Ristić¹

¹University of Niš, Faculty of Sciences and Mathematics, Department of Mathematics, dmiodrag@pmf.ni.ac.rs, anastic78@gmail.com, miristic72@gmail.com

The combined SDLINAR model of order p is a model of time series that is suitable for modeling phenomena whose values change over time through integer increments or decrements, and which, with certain probabilities, can be the result of the near or distant past. Under the spotlight of this work are results of parameter estimations of such models, as well as the problems that have arisen and possible directions for solving those problems.

References

- P. J. Brockwell, R. A. Davis, Time Series Theory and Methods. Springer, New York, 1991.
- [2] M. S. Djordjević, An extension on *INAR* models with discrete Laplace marginal distributions, Communication in Statistics - Theory and Methods, 46(12) (2017), 5896–5913.
- [3] M. S. Djordjević, A combined SDLINAR(p) model and identification and prediction of its latent components, 31(5), (2017), 919–946.
- [4] R. K. Freeland, B. McCabe, Asymptotic properties of CLS estimators in the Poisson AR(1) model, Statist. Probab. Lett. 73 (2005), 147–153.
- [5] A. S. Nastić, M. M. Ristić, H. S. Bakouch, A combined geometric *INAR(p)* model based on negative binomial thinning, Mathematical and Computer Modelling, **55** (2012), 1665–1672.
- [6] M. M. Ristić, A. S. Nastić, A mixed *INAR(p)* model, J. Time Ser. Anal. 33 (2012), 903–915.

- [7] I. Silva, M. E. Silva, Asymptotic distribution of the Yule-Walker estimator for INAR(p) processes, Statistic and Probability Letters, 76(15), (2006) 1655–1663.
- [8] D. Tjstheim, Estimation in nonlinear time series models, Stochastic Process Appl. 21 (1986), 251–273.
- [9] H. White, Asymptotic theory for econometrics, Academic Press, San Diego, 2001.