

On Internality of Generalized Averaged Gaussian Quadrature Rules and Their Truncations

Dušan Lj. Djukić¹, Rada M. Mutavdžić Djukić¹, Lothar Reichel², and Miodrag M. Spalević¹

¹University of Belgrade, Faculty of Mechanical Engineering, Department of Mathematics, ddjukic@mas.bg.ac.rs, rmutavdzic@mas.bg.ac.rs, mspalevic@mas.bg.ac.rs

²Kent State University, Department of Mathematical Sciences, reichel@math.kent.edu

Generalized (also called optimal) averaged Gauss quadrature formulas may yield higher accuracy than Gauss quadrature rules that use the same moment information. This is illustrated in [3]. They therefore may be attractive to use when moments or modified moments are cumbersome to evaluate. However, generalized averaged Gauss quadrature formulas may be not internal, i. e., they may have nodes outside the convex hull of the support of the measure that defines the associated Gauss rules; see, e.g., [2, 4, 1] for examples and analyses. It may therefore not be possible to use generalized averaged Gauss quadrature formulas with integrands that only are defined on the convex hull of the support of the measure. A survey of our results on internality of generalized averaged Gaussian quadrature rules and their truncations will be presented.

References

- [1] D. Lj. Djukić, R. M. Mutavdžić Djukić, L. Reichel, M. M. Spalević, Internality of generalized averaged Gauss quadrature rules and truncated variants for modified Chebyshev measures of the third and fourth kinds, *Numer. Algorithms* **92** (2023), 523–544.
- [2] D. P. Laurie, Anti-Gaussian quadrature formulas, *Math. Comp.* **65** (1996), 739–747.
- [3] L. Reichel, M. M. Spalević, Generalized averaged Gaussian quadrature formulas: Properties and applications, *J. Comput. Appl. Math.* **410** (2022), Art. 114232.

- [4] M. M. Spalević, On generalized averaged Gaussian formulas, *Math. Comp.* **76** (2007), 1483–1492.