

Different approaches in the study of φ -contraction

Marija Cvetković¹

¹University of Niš, Faculty of Sciences and Mathematics, Department of Mathematics,
marija.cvetkovic@pmf.edu.rs

The study of φ -contractions dates to 1968. and results of eminent researchers like F. E. Browder, D. W. Boyd and J. S. Wong. In the following half of century various modifications and improvements of this concept were in the spotlight of research in the area of Metric Fixed Point Theory. We intend to discuss on the relation of Boyd-Wong contraction, Matkowski contraction, Meir-Keeler contraction, Proinov contraction, F -contraction and θ -contraction. It is also of great importance to study possible applications and benefits of applying nonlinear contractions over Banach contraction and its modifications.

References

- [1] D.W. Boyd and J.S.W. Wong, *On nonlinear contractions*, Proc. Amer. Math. Soc. **20** (1969), 458-464.
- [2] F.E. Browder, *On the convergence of successive approximations for nonlinear functional equations*, Nederl. Akad. Wetensch. Proc. Ser. A Indag. Math., **30** (1968), 27-35.
- [3] M. Cvetković, *The relation between F -contraction and Meir-Keeler contraction*, Rev. R. Acad. Cienc. Exactas Fs. Nat. Ser. A Mat. **117**(1) (2023)
- [4] M. Cvetković, *Behind the concept of F -contraction*, Fixed Point Theory, in press
- [5] M. Jleli and B. Samet, *A new generalization of the Banach contraction principle*, J. Inequal. Appl. **38** (2014)
- [6] A. Meir and E. Keeler, *A theorem on contraction mappings*, J. Math. Anal. Appl. **28**(1969), 326–329.
- [7] P. D. Proinov, *Fixed point theorems for generalized contractive mappings in metric spaces*. J. Fixed Point Theory Appl. **22**(21) (2020)