

A TOTALLY RELAXED SELF-ADAPTIVE SUBGRADIENT EXTRAGRADIENT SCHEME FOR EQUILIBRIUM AND FIXED POINT PROBLEMS IN A BANACH SPACE

OLAWALE KAZEEM OYEWOLE^{1,2}, HAMMED ANUOLUWAPO ABASS^{1,2},
AND OLUWATOSIN TEMITOPE MEWOMO¹

ABSTRACT. The goal of this paper is to introduce a Totally Relaxed Self adaptive Subgradient Extragradient Method (TRSSEM) together with an Halpern iterative method for approximating a common solution of Fixed Point Problem (FPP) and Equilibrium Problem (EP) in 2-uniformly convex and uniformly smooth Banach space. We prove the strong convergence of the sequence generated by our proposed method. The proposed method does not require the computation of a projection onto a feasible set, it instead requires a projection onto a finite intersection of sub-level sets of convex functions. Our result generalizes, unifies and extends some related results in the literature.

1. INTRODUCTION

Let C be a nonempty, closed and convex subset of a real Banach space E with dual space E^* . Let E be endowed with the duality pairing $\langle \cdot, \cdot \rangle$ of element from E and E^* , and also the corresponding norm $\| \cdot \|$. Let $f : C \times C \rightarrow \mathbb{R} \cup \{+\infty\}$ be a bifunction such that $C \subset \text{int}(\text{dom}(f, \cdot))$, then for every $x \in C$, the Equilibrium Problem (EP) (see [3, 14]), is to find a point $x^* \in C$ such that

$$(1.1) \quad f(x^*, y) \geq 0, \quad \text{for all } y \in C.$$

We denote the EP and its solution set by $EP(C, f)$ and $Sol(C, f)$, respectively.

Key words and phrases. Equilibrium problem, strongly pseudomonotone, strong convergence, Banach space, quasi- ϕ -nonexpansive mapping, fixed point.

2020 *Mathematics Subject Classification.* Primary: 47H09. Secondary: 49J35, 90C47.

DOI

Received: September 21, 2021.

Accepted: February 02, 2022.

The EP is a generalization of many important optimization problems, such as Variational Inequality Problem (VIP), Fixed Point Problem (FPP) and so on (see [6, 14] and the references therein). In particular, if $f(x, y) = \langle Ax, y - x \rangle$, where $A : C \rightarrow E^*$, is a nonlinear mapping, then $EP(C, f)$ (1.1) reduces to the classical VIP introduced by Stampacchia [47] (see also [36, 38, 41, 52]), which is to find a point $x^* \in C$ such that

$$(1.2) \quad \langle Ax^*, y - x^* \rangle \geq 0, \quad \text{for all } y \in C.$$

There are two important directions of research on EP: These are the existence of solution of EP and other related problems (see [14, 29] for more details) and the development of iterative algorithms for approximating the solution of EP, its several generalizations and related optimization problems (see [1, 12, 13, 33, 34, 42–44] and the references therein).

In 2018, Hieu [24] introduced some methods for solving strongly pseudomonotone and Lipschitz type bifunction EPs. We note that a bifunction f satisfies the Lipschitz type condition, if there exist positive constants $c_1, c_2 \in \mathbb{R}$ such that for all $x, y, z \in C$, the inequality

$$f(x, y) + f(y, z) \geq f(x, z) - c_1 \|x - y\|^2 - c_2 \|y - z\|^2$$

holds.

In general EP, the Lipschitz type condition does not hold and when it does, finding the constants c_1 and c_2 is always not an easy task. This does have effect on the efficiency of the method involved. In addition, in the method of Hieu [24], there is the need to first solve at least one strongly convex programming problem. Also, if the bifunction and the feasible sets have complex structures, the computations could be expensive and time consuming.

Furthermore, the problem of finding a common point in the set of solutions of different generalizations of EP and the fixed point set of a nonlinear mapping in Hilbert, Banach and Hadamard spaces have been considered by several authors in literature (see [25, 39, 40, 46, 51, 57]) and the references therein for further reading.

In 2013, Anh [9] introduced an extragradient algorithm for finding a common element of the fixed point set of a nonexpansive mapping and solution set of an EP involving pseudomonotone and Lipschitz type continuous bifunction in real Hilbert space. The author proved a strong convergence result of the sequence generated by his method under some standard conditions, see [8–10] for related results.

However, in Banach spaces, just like the extragradient method employed by Hieu [24], many existing methods for approximating a common solution FPP and EP involving a pseudomonotone bifunctions requires that a strongly convex programming is solved (see [26, 27] and the references therein).

To avoid the assumptions of Lipschitz continuity on the bifunction and solving strongly convex programming, Vinh and Gibali [53] introduced two gradient-type iterative algorithms involving a one-step projection method for solving $EP(C, f)$ (1.1) and proved strong convergence results for both algorithms with an adaptive step-size

rule which does not require the Lipschitz condition of the associated method. The method proposed in [53] involves a projection onto a feasible set, and is known to be computationally expensive, time and memory consuming if the feasible set is not simple.

In an attempt to overcome this setback, Censor et al. [17] introduced the subgradient extragradient method which uses a projection onto a halfspace. Also, He et al. [23] introduced a TRSSEM for solving the VIP (1.2) in a real Hilbert space. Let $C^i := \{x \in H : h_i(x) \leq 0\}$, where $h_i : H \rightarrow \mathbb{R}$ for $i = 1, 2, \dots, m$, are convex functions. In the TRSSEM, the feasible set is given as

$$C := \bigcap_{i=1}^m C^i.$$

On the other hand, for approximating a fixed point of a nonexpansive mapping T , Mainge [31] introduced an inertial Krasnoselskij-Mann Algorithm as follows:

$$(1.3) \quad \begin{cases} w_n = x_n + \theta_n(x_n - x_{n-1}), \\ x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T w_n, \quad n \geq 1, \end{cases}$$

and proved a weak convergence theorem under some mild assumptions on the sequences $\{\theta_n\}$ and $\{\alpha_n\}$. The term $\theta_n(x_n - x_{n-1})$ as given (1.3) is referred to as the inertial extrapolation term. It is known that the introduction of the inertial term helps to speed up the convergence rate of the algorithm. Due to its importance, lots of researchers have adopted the use of the inertial technique in their quest for approximating the solutions of fixed point and optimization problem (see [4, 5, 31] and the references therein).

In this paper, motivated by the works of He et al. [23], Vinh and Gibali [53] and other related results in literature, we introduce a TRSSEM for approximating a common solution of FPP and EP in 2-uniformly convex and uniformly smooth Banach space. We prove a strong convergence result for the sequence generated by the proposed method under some conditions. Finally, we give some applications of our main result. The rest of the section is organized as follows. In Section 2, we recall some important results and definitions that will be useful in establishing our main result. In Section 3, we state our proposed method and then discuss its convergence analysis. We give some theoretical application of our main result in Section 4 and give a concluding remark Section 5.

2. PRELIMINARIES

We denote the weak and the strong convergence of a sequence $\{x_n\}$ to a point x by $x_n \rightharpoonup x$ and $x_n \rightarrow x$, respectively.

Let E be a real Banach space, given a function $g : E \rightarrow \mathbb{R}$.

• The function g is called Gâteaux differentiable at $x \in E$, if there exists an element E , denoted by $g'(x)$ or $\nabla g(x)$ such that

$$\lim_{t \rightarrow \infty} \frac{g(x + ty) - g(x)}{t} = \langle y, g'(x) \rangle, \quad y \in E,$$

where g' or $\nabla g(x)$ is called Gâteaux differential or gradient of g at x . We say g is Gâteaux on E if for each $x \in E$, g is Gâteaux differentiable at x .

- The function g is called weakly lower semicontinuous at $x \in E$, if $x_n \rightarrow x$ implies $g(x) \leq \liminf_{n \rightarrow \infty} g(x_n)$. We say that a function g is weakly lower semicontinuous on E , if for each $x \in E$, g is weakly lower semicontinuous at x .

- If g is a convex function, then it is said to be differentiable at a point $x \in E$ if the following set

$$(2.1) \quad \partial g(x) = \{f \in E : g(y) - g(x) \geq \langle f, y - x \rangle, y \in E\}$$

is nonempty. Each element $\partial g(x)$ is called a subgradient of g at x or the subdifferential of g and the inequality (2.1) is said to be the subdifferential inequality of g at x .

The function g is subdifferentiable at x , if g is subdifferentiable at every $x \in E$. It is well known that if g is Gâteaux differentiable at x , then g is subdifferentiable at x and $\partial g(x) = \{g'(x)\}$, that is, $\partial g(x)$ is just a singleton set. For more details on Gâteaux differentiable functions on Banach space, see [15].

Let C be a nonempty, closed and convex subset of a real Banach space with norm $\|\cdot\|$ and let $J : E \rightarrow 2^{E^*}$ be the normalized duality mapping defined by

$$J(x) = \{x^* \in E^* : \langle x, x^* \rangle = \|x\|^2 = \|x^*\|^2 \text{ for all } x \in E\},$$

where E^* denotes the dual space of E and $\langle \cdot, \cdot \rangle$ the duality pairing between the elements of E and E^* . Alber [7], introduced a generalized projection operator Π_C an analogue of the metric projection $P_C : H \rightarrow C$ in the Hilbert space H . He defines $\Pi_C : E \rightarrow C$ by

$$\Pi_C(x) = \inf_{y \in C} \{\phi(y, x) \text{ for all } x \in E\}.$$

In Hilbert spaces $P_C(x) \equiv \Pi_C(x)$.

Consider the Lyapunov functional $\phi : E \times E \rightarrow \mathbb{R}^+$ defined by

$$\phi(x, y) = \|x\|^2 - 2\langle x, Jy \rangle + \|y\|^2, \quad \text{for all } x, y \in E.$$

In the real Hilbert space, we observe that $\phi(x, y) = \|x - y\|^2$. It is easy to see that

$$(\|x\| - \|y\|)^2 \leq \phi(x, y) \leq (\|x\| + \|y\|)^2.$$

The functional ϕ also satisfies the following important properties:

$$(2.2) \quad \phi(x, y) = \phi(x, z) + \phi(z, y) + 2\langle x - z, Jz - Jy \rangle$$

and

$$(2.3) \quad \phi\left(x, J^{-1}(\lambda Jy + (1 - \lambda)Jz)\right) \leq \lambda\phi(x, y) + (1 - \lambda)\phi(x, z),$$

for all $x, y, z \in E$ and $\lambda \in (0, 1)$.

Note. If E is a reflexive, strictly convex, and smooth Banach space, then for $x, y \in E$, $\phi(x, y) = 0$ if and only if $x = y$, see [18, 48].

We are also concerned with the functional $V : E \times E^* \rightarrow \mathbb{R}$ defined by

$$(2.4) \quad V(x, x^*) = \|x\|^2 - 2\langle x, x^* \rangle + \|x^*\|^2,$$

for all $x \in E$ and $x^* \in E^*$. That is, $V(x, x^*) = \phi(x, J^{-1}x^*)$ for all $x \in E$ and $x^* \in E^*$. It is well known that if E is a reflexive, strictly convex and smooth Banach space, then

$$V(x, x^*) \leq V(x, x^* + y^*) - 2 \langle J^{-1}x^* - x, y^* \rangle,$$

for all $x \in E$ and all $x^*, y^* \in E^*$, see [50].

Let C be a closed and convex subset of E and $T : C \rightarrow C$ be a mapping, a point $x \in C$ is called a fixed point of T , if $x = Tx$. We denote the set of fixed points of T by $F(T)$. Let $T : C \rightarrow C$ be a mapping, a point $p \in C$ is called an asymptotic fixed point of T (see [45]) if C contains a sequence $\{x_n\}$ such that $x_n \rightarrow p$ and $\|x_n - Tx_n\| \rightarrow 0$ as $n \rightarrow \infty$. We denote by $\hat{F}(T)$ the set of asymptotic fixed points of T . A mapping $T : C \rightarrow C$ is said to be relatively nonexpansive if $\hat{F}(T) = F(T)$ and $\phi(p, Tx) \leq \phi(p, x)$ for all $x \in C$ and $p \in F(T)$ (see [16, 48]). T is said to be ϕ -nonexpansive if $\phi(Tx, Ty) \leq \phi(x, y)$ for all $x, y \in C$ and quasi- ϕ -nonexpansive if $F(T) \neq \emptyset$ and $\phi(p, Tx) \leq \phi(p, x)$ for all $x \in C$ and $p \in F(T)$.

The class of quasi- ϕ -nonexpansive mappings is more general than the class of relatively nonexpansive mapping which requires the strict condition $F(T) = \hat{F}(T)$ (see [16, 45, 48]).

Let E be a real Banach space. The modulus of convexity of E is the function $\delta_E : (0, 2] \rightarrow [0, 1]$ defined by

$$\delta_E(\epsilon) = \inf \left\{ 1 - \frac{1}{2} \|x + y\| : \|x\| = \|y\| = 1, \|x - y\| \geq \epsilon \right\}.$$

Recall that E is said to be uniformly convex if $\delta_E(\epsilon) > 0$ for any $\epsilon \in (0, 2]$. E is said to be strictly convex if $\frac{\|x+y\|}{2} < 1$ for all $x, y \in E$, with $\|x\| = \|y\| = 1$ and $x \neq y$. Also, E is p -uniformly convex if there exists a constant $c_p > 0$ such that $\delta_E(\epsilon) > c_p \epsilon^p$ for any $\epsilon \in (0, 2]$.

The modulus of smoothness of E is the function $\rho_E : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ defined by

$$\rho_E(t) = \sup \left\{ \frac{1}{2} (\|x + ty\| - \|x - ty\|) - 1 : \|x\| = \|y\| = 1 \right\}.$$

E is said to be uniformly smooth if $\lim_{t \rightarrow 0} \frac{\rho_E(t)}{t} = 0$. Let $1 < q \leq 2$, then E is q -uniformly smooth if there exists $c_q > 0$ such that $\rho_E(t) \leq c_q t^q$ for $t > 0$. It is known that E is p -uniformly convex if and only if E^* is q -uniformly smooth, where $p^{-1} + q^{-1} = 1$. It is also known that every q -uniformly smooth Banach space is uniformly smooth.

It is widely known that if E is uniformly smooth, then the duality mapping J is norm-to-norm continuous on each bounded subset of E . The following are some important and useful properties of J , for further details, see [2, 48].

Let C be a nonempty, closed and convex subset of a real Banach space E and $f : E \times E \rightarrow \mathbb{R} \cup \{+\infty\}$ be a bifunction. f is said to be

(i) strongly monotone on C , if there exists $\gamma \geq 0$ such that for any $x, y \in C$

$$f(x, y) + f(y, x) \leq -\gamma\|x - y\|^2;$$

(ii) monotone on C , if

$$f(x, y) + f(y, x) \leq 0, \quad \text{for all } x, y \in C;$$

(iii) pseudomonotone on C , if

$$f(x, y) \geq 0 \Rightarrow f(y, x) \leq 0, \quad \text{for all } x, y \in C;$$

(iv) strongly γ -pseudomonotone on C , if there exists $\gamma > 0$ such that for any $x, y \in C$

$$f(x, y) \geq 0 \Rightarrow f(y, x) \leq -\gamma\|x - y\|^2.$$

From the above, it is clear (i) \Rightarrow (ii) \Rightarrow (iii) \Rightarrow (iv). The converse is generally not true (see [53]).

We now give the following useful and important lemmas that are needed in establishing our main results.

Lemma 2.1 ([35]). *Let E be a 2-uniformly convex and smooth Banach space. Then for every $x, y \in E$*

$$\phi(x, y) \geq \nu\|x - y\|^2,$$

where $\nu > 0$ is the 2-uniformly convexity constant of E .

Lemma 2.2 ([28]). *Let E be a smooth and uniformly convex real Banach space and let $\{x_n\}$ and $\{y_n\}$ be two sequences in E . If either $\{x_n\}$ or $\{y_n\}$ is bounded and $\phi(x_n, y_n) \rightarrow 0$ as $n \rightarrow \infty$, then $\|x_n - y_n\| \rightarrow 0$ as $n \rightarrow \infty$.*

Lemma 2.3 ([7]). *Let C be a nonempty, closed and convex subset of a reflexive, strictly convex and smooth Banach space X . If $x \in E$ and $q \in C$, then*

$$(2.5) \quad q = \Pi_C x \iff \langle y - q, Jx - Jq \rangle \leq 0, \quad \text{for all } y \in C,$$

and

$$(2.6) \quad \phi(y, \Pi_C x) + \phi(\Pi_C x, x) \leq \phi(y, x), \quad \text{for all } y \in C, x \in X.$$

Lemma 2.4 ([55]). *Fix a number $s > 0$. A real Banach space X is uniformly convex if and only if there exists a continuous strictly increasing function $\psi : [0, \infty) \rightarrow [0, \infty)$ with $\psi(0) = 0$ such that*

$$\|tx + (1 - t)y\|^2 \leq t\|x\|^2 + (1 - t)\|y\|^2 - t(1 - t)\psi(\|x - y\|),$$

for all $x, y \in X$, $\lambda \in [0, 1]$, with $\|x\| < s$ and $\|y\| < s$.

Lemma 2.5 ([54]). *Let $\{a_n\}$ be a sequence of nonnegative real numbers satisfying the following relation*

$$a_{n+1} \leq (1 - \alpha_n)a_n + \alpha_n\sigma_n + \gamma_n, \quad n \geq 0,$$

where

- (a) $\{\alpha_n\} \subset [0, 1]$, $\lim_{n \rightarrow \infty} \alpha_n = 0$ and $\sum_{n=1}^{\infty} \alpha_n = \infty$;
- (b) $\limsup_{n \rightarrow \infty} \sigma \leq 0$;
- (c) $\gamma_n \geq 0$, $n \geq 1$, and $\sum_{n=1}^{\infty} \gamma_n < \infty$.

Then, $\lim_{n \rightarrow \infty} a_n = 0$.

Lemma 2.6 ([32]). *Let $\{a_n\}$ be a sequence of real numbers such that there exists a subsequence $\{n_j\}$ of $\{n\}$ such that $a_{n_j} < a_{n_{j+1}}$ for all $j \in \mathbb{N}$. Then, there exists a nondecreasing subsequence $\{m_n\} \subset \mathbb{N}$ such that $m_n \rightarrow \infty$ and the following properties are satisfied by all (sufficiently large) numbers $n \in \mathbb{N}$: $a_{m_n} < a_{m_{n+1}}$ and $a_n < a_{m_{n+1}}$. In fact, $m_n = \max\{i \leq k : a_i < a_{i+1}\}$.*

3. MAIN RESULT

In this section, we give a concise and precise statement of our algorithm, discuss some of its elementary properties and its convergence analysis. The convergence analysis is given in the next section.

Statement 3.1. Let C be a nonempty, closed and convex subset of a 2-uniformly convex and uniformly smooth real Banach space E with dual space E^* . For $i = 1, 2, \dots, m$, let $h_i : E \rightarrow \mathbb{R}$ be a family of convex, weakly lower semicontinuous and Gâteaux differentiable functions. Let $S : E \rightarrow E$ be a quasi- ϕ -nonexpansive mapping and $f : C \times C \rightarrow \mathbb{R} \cup \{+\infty\}$ be a strongly γ -pseudomonotone bifunction satisfying the following assumptions.

Assumption 3.2. We require the following assumptions for our operator and the solution set:

- A1. $f(x, \cdot)$ is convex and lower semi-continuous for every $x \in E$;
- A2. f is strongly γ -pseudomonotone on C ,
- A3. $Sol(C, f) \neq \emptyset$;
- A4. if $\{x_n\}_{n=0}^{\infty} \subset E$ is bounded, then the sequence $\{g(x_n) \in \partial(f(x_n, \cdot))(x_n)_{n=0}^{\infty}\}$ is bounded.

Note. The assumption A4. is quite standard assumption and it holds for example when $f(x, \cdot)$ is bounded on bounded subsets (see [11]).

Assumption 3.3. To prove a strong convergence result using Algorithm 3.4, the following conditions are needed.

- B1. The feasible set C is defined by $C := \cap_{i=1}^m C^i$, where $C^i := \{z \in E : h_i(z) \leq 0\}$;
- B2. $\lim_{n \rightarrow \infty} \alpha_n = 0$ and $\sum_{n=0}^{\infty} \alpha_n = \infty$;
- B3. $0 < \liminf_{n \rightarrow \infty} \gamma_n \leq \limsup_{n \rightarrow \infty} \gamma_n < 1$;
- B4. $\sum_{n=1}^{\infty} \phi(x_n, x_{n-1}) < \infty$.
- B5. $\lim_{n \rightarrow \infty} \frac{\theta_n}{\alpha_n} = 0$.

Algorithm 3.4. (TRSSEM) for EP(C, f)

Step 0. Choose the sequences $\{\theta_n\}$, $\{\alpha_n\}$ and $\{\gamma_n\} \subset (0, 1)$ satisfying Assumption 3.3, let $\mu \in (0, 1)$ and $\beta_0 > 0$. For $u \in C$, select initial points x_0 and x_1 in C . Set $n = 1$.

Step 1. For $i = 1, 2, \dots, m$, and given the current iterate w_n , construct the family of half spaces

$$C_n^i := \{z \in E : h_i(w_n) + \langle h'_i(w_n), z - w_n \rangle \leq 0\}$$

and set

$$C_n = \bigcap_{i=1}^m C_n^i.$$

Let $w_n := J^{-1}(Jx_n + \theta_n(Jx_{n-1} - Jx_n))$. Take $g(w_n) \in \partial(f(w_n, \cdot))(w_n)$, $n \geq 1$, and compute

$$(3.1) \quad z_n = \Pi_{C_n} J^{-1}(Jw_n - \beta_n g(w_n)),$$

where β_n is given by

$$(3.2) \quad \beta_{n+1} = \begin{cases} \min \left\{ \beta_n, \frac{\mu \|w_n - z_n\|}{\|g(w_n) - g(z_n)\|} \right\}, & \text{if } g(w_n) \neq g(z_n), \\ \beta_n, & \text{otherwise.} \end{cases}$$

Step 2. If $w_n = z_n$ ($w_n \in \text{Sol}(C, f)$), then set $w_n = y_n$ and go to Step 3. Otherwise, compute the next iterate by

$$(3.3) \quad y_n = \Pi_{Q_n} J^{-1}(Jw_n - \beta_n g(z_n)),$$

where

$$Q_n = \{w \in E : \langle w - z_n, Jw_n - \beta_n g(w_n) - Jz_n \rangle \leq 0\}.$$

Step 3. Compute

$$(3.4) \quad x_{n+1} = J^{-1}((1 - \alpha_n)Ju + \alpha_n(1 - \gamma_n)Jy_n + \gamma_n JSy_n).$$

Step 4. Set $n := n + 1$ and go to Step 1.

Lemma 3.1. *If $w_n = z_n$, then $w_n \in \text{Sol}(C, f)$.*

Proof. Suppose $w_n = z_n$, then by (2.5) and (3.1), we have

$$\langle Jw_n - \beta_n g(w_n) - Jw_n, y - z_n \rangle \leq 0, \quad y \in C,$$

or equivalently

$$(3.5) \quad \langle g(w_n), y - w_n \rangle \geq 0, \quad \text{for all } y \in C.$$

Therefore, from (3.5) and the definition of the subdifferential f in the second argument, we obtain

$$f(w_n, y) = f(w_n, y) - f(w_n, w_n) \geq \langle g(w_n), y - w_n \rangle \geq 0.$$

Hence, $w_n \in \text{Sol}(C, f)$. □

Lemma 3.2 ([56]). *The sequence $\{\beta_n\}$ generated by (3.2) is a monotonically decreasing sequence and*

$$\lim_{n \rightarrow \infty} \beta_n = \beta \geq \min \left\{ \frac{\mu}{L}, \beta_0 \right\}.$$

Remark 3.1. Note that if $w_n = z_n$ and $w_n = Sw_n$ we are at a common solution of the $EP(C, f)$ and fixed point of the mapping S . In our convergence analysis, we will assume implicitly that this does not occur after finitely many iterations so that our Algorithm 3.4 generates an infinite sequence satisfying, in particular $w_n \neq z_n$ and $w_n \neq Sw_n$ for all $n \in \mathbb{N}$.

We now prove some lemmas which are required components of the main result.

Lemma 3.3. *The sequence $\{x_n\}$ generated by Algorithm 3.4 is bounded.*

Proof. Let $x^* \in \text{Sol}(C, f)$, then we have from (2.6), that

$$\begin{aligned} \phi(x^*, y_n) &= \phi(x^*, \Pi_{Q_n} J^{-1}(Jw_n - \beta_n g(w_n))) \\ &\leq \phi(x^*, J^{-1}(Jw_n - \beta_n g(z_n))) - \phi(y_n, J^{-1}(Jw_n - \beta_n g(w_n))) \\ &= \|x^*\|^2 - 2\langle x^*, Jw_n - \beta_n g(z_n) \rangle - \|y_n\|^2 + 2\langle y_n, Jw_n - \beta_n g(z_n) \rangle \\ &= \phi(x^*, w_n) - \phi(y_n, w_n) + 2\beta_n \langle x^* - y_n, g(z_n) \rangle \\ &= \phi(x^*, w_n) - (\phi(y_n, z_n) + \phi(z_n, w_n)) \\ &\quad + 2\langle y_n - z_n, Jz_n - Jw_n \rangle + 2\beta_n \langle x^* - y_n, g(z_n) \rangle \\ &= \phi(x^*, w_n) - \phi(y_n, z_n) - \phi(z_n, w_n) \\ (3.6) \quad &\quad + 2\langle y_n - z_n, Jw_n - Jz_n \rangle + 2\beta_n \langle x^* - y_n, g(z_n) \rangle. \end{aligned}$$

Now, we have from (3.6) that

$$\begin{aligned} 2\beta_n \langle x^* - y_n, g(z_n) \rangle &= 2\beta_n \langle x^* - z_n, g(z_n) \rangle + 2\beta_n \langle z_n - y_n, g(z_n) \rangle \\ (3.7) \quad &= 2\beta_n \langle x^* - z_n, g(z_n) \rangle + 2\langle y_n - z_n, -\beta_n g(z_n) \rangle. \end{aligned}$$

Substituting (3.7) into (3.6) and using the strongly pseudomonotonicity of f , we obtain

$$\begin{aligned} \phi(x^*, y_n) &= \phi(x^*, w_n) - \phi(y_n, z_n) - \phi(z_n, w_n) + 2\langle y_n - z_n, Jw_n - Jz_n \rangle \\ &\quad + 2\beta_n \langle x^* - z_n, g(z_n) \rangle + 2\langle y_n - z_n, -\beta_n g(z_n) \rangle \\ &= \phi(x^*, w_n) - \phi(y_n, z_n) - \phi(z_n, w_n) + 2\langle y_n - z_n, Jw_n - \beta_n g(z_n) - Jz_n \rangle \\ &\quad + 2\beta_n \langle x^* - z_n, g(z_n) \rangle \\ &\leq \phi(x^*, w_n) - \phi(y_n, z_n) - \phi(z_n, w_n) \\ &\quad + 2\beta_n \langle y_n - z_n, Jw_n - \beta_n g(z_n) - Jz_n \rangle + 2\beta_n f(z_n, x^*) \\ &\leq \phi(x^*, w_n) - \phi(y_n, z_n) - \phi(z_n, w_n) \\ &\quad - 2\beta_n \gamma \phi(x^*, z_n) + 2\langle y_n - z_n, Jw_n - \beta_n g(z_n) - Jz_n \rangle \\ (3.8) \quad &\leq \phi(x^*, w_n) - \phi(y_n, z_n) - \phi(z_n, w_n) + 2\langle y_n - z_n, Jw_n - \beta_n g(z_n) - Jz_n \rangle. \end{aligned}$$

By the definition of Q_n and Cauchy-Schwartz inequality, we have

$$\begin{aligned}
\langle y_n - z_n, Jw_n - \beta_n g(z_n) - Jz_n \rangle &= 2\langle y_n - z_n, Jw_n - \beta_n g(z_n) - Jz_n \rangle \\
&\quad + 2\beta_n \langle y_n - z_n, g(w_n) - g(z_n) \rangle \\
(3.9) \qquad \qquad \qquad &\leq 2\beta_n \|y_n - z_n\| \|g(w_n) - g(z_n)\|.
\end{aligned}$$

Using (3.2) and Lemma 2.1 in (3.9), we get

$$\begin{aligned}
\langle y_n - z_n, Jw_n - \beta_n g(z_n) - Jz_n \rangle &\leq 2 \frac{\mu\beta_n}{\beta_{n+1}} \|y_n - z_n\| \|w_n - z_n\| \\
&\leq 2 \frac{\mu\beta_n}{\beta_{n+1}} \sqrt{\frac{\phi(y_n, z_n)}{\nu}} \sqrt{\frac{\phi(z_n, w_n)}{\nu}} \\
(3.10) \qquad \qquad \qquad &\leq \frac{\mu\beta_n}{\nu\beta_{n+1}} (\phi(y_n, z_n) + \phi(z_n, w_n)).
\end{aligned}$$

Therefore, from (3.8) and (3.10), we have

$$(3.11) \quad \phi(x^*, y_n) \leq \phi(x^*, w_n) - \left(1 - \frac{\mu\beta_n}{\nu\beta_{n+1}}\right) (\phi(y_n, z_n) + \phi(z_n, w_n)).$$

From (2.3) and (3.4), we have

$$\begin{aligned}
\phi(x^*, x_{n+1}) &= \phi(x^*, J^{-1}(\alpha_n Ju + (1 - \alpha_n)(1 - \gamma_n)Ju_n + \gamma_n JSy_n)) \\
&= \phi(x^*, J^{-1}(\alpha_n Ju + (1 - \alpha_n)(1 - \gamma_n)Jy_n + (1 - \alpha_n)\gamma_n JSy_n)) \\
&\leq \alpha_n \phi(x^*, u) + (1 - \alpha_n)(1 - \gamma_n)\phi(x^*, y_n) + (1 - \alpha_n)\gamma_n \phi(x^*, Sy_n) \\
&\leq \alpha_n \phi(x^*, u) + (1 - \alpha_n)\phi(x^*, y_n) \\
&\leq \alpha_n \phi(x^*, u) + (1 - \alpha_n)\phi(x^*, w_n) \\
&\quad - \left(1 - \frac{\mu\beta_n}{\nu\beta_{n+1}}\right) (\phi(y_n, z_n) + \phi(z_n, w_n)) \\
(3.12) \qquad \qquad &\leq \alpha_n \phi(x^*, u) + (1 - \alpha_n)\phi(x^*, w_n).
\end{aligned}$$

From Algorithm 3.4, we have

$$\begin{aligned}
\phi(x^*, w_n) &= \phi(x^*, J^{-1}(Jx_n + \theta_n(Jx_{n-1} - Jx_n))) \\
&\leq (1 - \theta_n)\phi(x^*, x_n) + \theta_n\phi(x^*, x_{n-1}),
\end{aligned}$$

hence

$$\begin{aligned}
\phi(x^*, x_{n+1}) &\leq \alpha_n \phi(x^*, u) + (1 - \alpha_n)((1 - \theta_n)\phi(x^*, x_n) + \theta_n\phi(x^*, x_{n-1})) \\
&\leq \alpha_n \phi(x^*, u) + (1 - \alpha_n)(\phi(x^*, x_n) + \phi(x^*, x_{n-1})) \\
&\leq \max\{\phi(x^*, u), (\phi(x^*, x_n) + \phi(x^*, x_{n-1}))\} \\
&\quad \vdots \\
(3.13) \qquad \qquad &\leq \max\{\phi(x^*, u), (\phi(x^*, x_1) + \phi(x^*, x_0))\}, \quad n \geq 1.
\end{aligned}$$

This implies that $\{\phi(x^*, x_n)\}$ is bounded. Therefore, $\{x_n\}$ is bounded. Consequently, $\{g(y_n)\}$ is bounded and by the nonexpansiveness of the projection operator and the mapping S , we have that $\{z_n\}$, $\{w_n\}$, $\{y_n\}$ and $\{Sy_n\}$ are bounded. \square

The boundedness of $\{x_n\}$ implies that there is at least one weak limit point. The next result provides a condition under which each of such weak limit is in the solution set of the equilibrium problem.

Lemma 3.4. *Let $\{x_{n_k}\}$ be a subsequence of $\{x_n\}$ converging weakly to a point $p \in C$ and suppose that the conditions $\|w_{n_i} - z_{n_i}\| \rightarrow 0$ and $\|w_{n_i} - x_{n_i}\| \rightarrow 0$ as $i \rightarrow \infty$ hold on this subsequence. Then $p \in \text{Sol}(C, f)$.*

Proof. From Lemma 2.5 and the definition of subdifferential, we have

$$\begin{aligned}
 0 &\leq \langle x - z_{n_i}, Jz_{n_i} - (Jw_{n_i} - \beta_{n_i}g(w_{n_i})) \rangle \\
 &= \langle x - z_{n_i}, Jz_{n_i} - Jw_{n_i} \rangle + \langle x - z_{n_i}, \beta_{n_i}g(w_{n_i}) \rangle \\
 &= \langle x - z_{n_i}, Jz_{n_i} - Jw_{n_i} \rangle + \langle x - w_{n_i}, \beta_{n_i}g(w_{n_i}) \rangle + \langle w_{n_i} - z_{n_i}, \beta_{n_i}g(w_{n_i}) \rangle \\
 (3.14) \quad &\leq \langle x - z_{n_i}, Jz_{n_i} - Jw_{n_i} \rangle + \langle w_{n_i} - z_{n_i}, \beta_{n_i}g(w_{n_i}) \rangle + f(w_{n_i}, x).
 \end{aligned}$$

Passing limit to the inequality in (3.14), we have

$$f(p, x) \geq 0, \quad \text{for all } x \in C. \quad \square$$

In proving the strong convergence of our Algorithm 3.4, the underlying idea relies on certain estimate and other classical properties of the iterates which are given in the next lemmas below.

Lemma 3.5. *The sequence $\{x_n\}$ generated by Algorithm 3.4 satisfies the following estimates:*

- (i) $a_{n+1} \leq (1 - \alpha_n)a_n + \alpha_n b_n$;
- (ii) $-1 \leq \limsup_{n \rightarrow \infty} b_n < +\infty$,

where $a_n = \phi(x^*, x_n)$ and $b_n = \frac{\theta_n}{\alpha_n} \phi(x^*, x_{n-1}) + 2\langle Ju - Jx^*, x_{n+1} - x^* \rangle$.

Proof. Let $p_n = (1 - \gamma_n)Jy_n + \gamma_nJSy_n$, then from (2.4), we have

$$\begin{aligned}
 \phi(x^*, x_{n+1}) &= \phi(x^*, J^{-1}(\alpha_n Ju + (1 - \alpha_n)Jp_n)) \\
 &\leq V(x^*, \alpha_n Ju + (1 - \alpha_n)Jp_n - \alpha_n(Ju - Jx^*)) \\
 &\quad - 2\langle -\alpha_n(Ju - Jx^*), J^{-1}(\alpha_n Ju + (1 - \alpha_n)Jp_n) \rangle \\
 &\leq V(x^*, \alpha_n Jx^* + (1 - \alpha_n)Jp_n) + 2\alpha_n \langle Ju - Jx^*, x_{n+1} - x^* \rangle \\
 &\leq \alpha_n V(x^*, Jx^*) + (1 - \alpha_n)V(x^*, Jp_n) + 2\alpha_n \langle Ju - Jx^*, x_{n+1} - x^* \rangle \\
 &\leq \alpha_n \phi(x^*, x^*) + (1 - \alpha_n)\phi(x^*, p_n) + 2\alpha_n \langle Ju - Jx^*, x_{n+1} - x^* \rangle \\
 &\leq (1 - \alpha_n)\phi(x^*, p_n) + 2\alpha_n \langle Ju - Jx^*, x_{n+1} - x^* \rangle \\
 &\leq (1 - \alpha_n)(1 - \gamma_n)\phi(x^*, y_n) + \gamma_n(1 - \alpha_n)\phi(x^*, Sy_n) \\
 &\quad + 2\alpha_n \langle Ju - Jx^*, x_{n+1} - x^* \rangle
 \end{aligned}$$

$$\begin{aligned}
&\leq (1 - \alpha_n)\phi(x^*, y_n) + 2\alpha_n\langle Ju - Jx^*, x_{n+1} - x^* \rangle \\
&\leq (1 - \alpha_n)\phi(x^*, w_n) + 2\alpha_n\langle Ju - Jx^*, x_{n+1} - x^* \rangle \\
&= (1 - \alpha_n)((1 - \theta_n)\phi(x^*, x_n) + \theta_n\phi(x^*, x_{n-1}) + 2\alpha_n\langle Ju - Jx^*, x_{n+1} - x^* \rangle) \\
&\leq (1 - \alpha_n)\phi(x^*, x_n) + \alpha_n \left(\frac{\theta_n}{\alpha_n}\phi(x^*, x_{n-1}) + 2\langle Ju - Jx^*, x_{n+1} - x^* \rangle \right).
\end{aligned}$$

This established (i). Next we proof (ii). Since $\{x_n\}$ is bounded, then we have

$$\sup_{n \geq 0} b_n \leq \sup \left(\frac{\theta_n}{\alpha_n}\phi(x^*, x_{n-1}) + 2\|Ju - Jx^*\|\|x_{n+1} - x^*\| \right) < \infty.$$

This implies that $\limsup_{n \rightarrow \infty} b_n < \infty$. Next we show that $\limsup_{n \rightarrow \infty} b_n \geq -1$. Assume the contrary, that is $\limsup_{n \rightarrow \infty} b_n \leq -1$. Then there exists $n_0 \in \mathbb{N}$ such that $b_n < -1$ for all $n \geq n_0$. Then for all $n_0 \in \mathbb{N}$, we get from (i), that

$$\begin{aligned}
a_{n+1} &\leq (1 - \alpha_n)a_n + \alpha_n b_n \\
&< (1 - \alpha_n)a_n - \alpha_n \\
&= a_n - \alpha_n(a_n + 1) \leq a_n - \alpha_n.
\end{aligned}$$

Taking lim sup of both sides in the last inequality, we have

$$\limsup_{n \rightarrow \infty} a_n \leq a_{n_0} - \lim_{n \rightarrow \infty} \sum_{i=n_0}^n \alpha_i = -\infty.$$

This contradicts the definition of $\{a_n\}$ as a nonnegative integer.

Therefore, $\limsup_{n \rightarrow \infty} b_n \geq -1$. □

We now present our main theorem.

Theorem 3.5. *Let C be a nonempty, closed and convex subset of a 2-uniformly convex and uniformly smooth real Banach space E and $h_i : E \rightarrow \mathbb{R}$ be a family of convex, weakly lower semicontinuous and Gâteaux differentiable functions, for $i = 1, 2, \dots, m$. Let $f : E \times E \rightarrow \mathbb{R} \cup \{+\infty\}$ be a bifunction satisfying conditions A1-A4, let $S : C \rightarrow C$ be a quasi- ϕ -nonexpansive mapping such that $\Gamma = \{Sol(C, f) \cap F(S)\} \neq \emptyset$. Let $\{\theta_n\}$, $\{\beta_n\}$ and $\{\alpha_n\}$ be sequences in $(0, 1)$ satisfying Assumption 3.3, then the sequence $\{x_n\}$ generated by Algorithm 3.4 converges strongly to $p = \Pi_\Gamma u$, where Π_Γ is the projection of C onto Γ .*

Proof. Let $p \in \Gamma$, we divide the proof into two cases.

Case I Suppose that there exists $n_0 \in \mathbb{N}$ such that $\{\phi(x^*, x_n)\}$ is monotone non-increasing. Since $\{\phi(x^*, x_n)\}$ is bounded, then it is convergent and

$$(3.15) \quad \phi(x^*, x_n) - \phi(x^*, x_{n+1}) \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Since $p_n = J^{-1}((1 - \gamma_n)Jy_n + \gamma_nJSy)$, then from Lemma 2.4, we have

$$\phi(x^*, p_n) = \phi(x^*, J^{-1}((1 - \gamma_n)Jy_n + \gamma_nJSy))$$

$$\begin{aligned}
&= V(x^*, (1 - \gamma_n)Jy_n + \gamma_nJSy) \\
&= \|x^*\|^2 - 2\langle x^*, (1 - \gamma_n)Jy_n + \gamma_nJSy \rangle + \|(1 - \gamma_n)Jy_n + \gamma_nJSy\|^2 \\
&= \|x^*\|^2 - 2(1 - \gamma_n)\langle x^*, Jy_n \rangle - 2\gamma_n\langle x^*, JSy_n \rangle + (1 - \gamma_n)\|y_n\|^2 + \gamma_n\|Sy_n\|^2 \\
&\quad - \gamma_n(1 - \gamma_n)\psi(\|Jy_n - JSy_n\|) \\
&= \phi(x^*, y_n) + \phi(x^*, Sy_n) - \gamma_n(1 - \gamma_n)\psi(\|Jy_n - JSy_n\|) \\
(3.16) \quad &\leq \phi(x^*, y_n) - \gamma_n(1 - \gamma_n)\psi(\|Jy_n - JSy_n\|).
\end{aligned}$$

Therefore, from (3.4), (3.11) and (3.16), we have

$$\begin{aligned}
\phi(x^*, x_{n+1}) &= \phi(x^*, J^{-1}(\alpha_n Ju + (1 - \alpha_n)Jp_n)) \\
&\leq \alpha_n\phi(x^*, u) + (1 - \alpha_n)\phi(x^*, p_n) \\
&\leq \alpha_n\phi(x^*, u) + (1 - \alpha_n)\phi(x^*, y_n) - \gamma_n(1 - \gamma_n)\psi(\|Jy_n - JSy_n\|) \\
&\leq \alpha_n\phi(x^*, u) + (1 - \alpha_n)\phi(x^*, w_n) - \gamma_n(1 - \gamma_n)\psi(\|Jy_n - JSy_n\|) \\
&= \alpha_n\phi(x^*, u) + (1 - \alpha_n)((1 - \theta_n)\phi(x^*, x_n) \\
&\quad + \theta_n\phi(x^*, x_{n-1})) - \gamma_n(1 - \gamma_n)\psi(\|Jy_n - JSy_n\|) \\
&\leq \alpha_n\phi(x^*, u) + (1 - \alpha_n)\phi(x^*, x_n) + \theta_n\phi(x^*, x_{n-1}) \\
(3.17) \quad &\quad - \gamma_n(1 - \gamma_n)\psi(\|Jy_n - JSy_n\|).
\end{aligned}$$

Hence,

$$\begin{aligned}
\gamma_n(1 - \gamma_n)\psi(\|Jy_n - JSy_n\|) &\leq \alpha_n \left(\frac{\theta_n}{\alpha_n}\phi(x^*, x_{n-1}) + \phi(x^*, u) \right) \\
&\quad + (1 - \alpha_n)\phi(x^*, x_n) - \phi(x^*, x_{n-1}).
\end{aligned}$$

By using $\alpha_n \rightarrow 0$, we obtain $\gamma_n(1 - \gamma_n)\psi(\|Jy_n - JSy_n\|) \rightarrow 0$ as $n \rightarrow \infty$. Therefore, by condition B3 and the property of ψ , we get

$$\lim_{n \rightarrow \infty} \|Jy_n - JSy_n\| = 0.$$

Since J^{-1} is norm-to-norm continuous on bounded subsets of E , we obtain

$$(3.18) \quad \lim_{n \rightarrow \infty} \|y_n - Sy_n\| = 0.$$

Furthermore, from (3.12), we have

$$\phi(x^*, y_n) \leq \phi(x^*, w_n) - \left(1 - \frac{\mu\beta_n}{\nu\beta_{n+1}}\right) (\phi(y_n, z_n) + \phi(z_n, w_n)).$$

Therefore, it follows from (3.4) that

$$\begin{aligned}
\phi(x^*, x_{n+1}) &\leq \alpha_n\phi(x^*, u) + (1 - \alpha_n)\phi(x^*, p_n) \\
&\leq \alpha_n\phi(x^*, u) + (1 - \alpha)\phi(x^*, y_n) \\
&\leq \alpha_n\phi(x^*, u) + (1 - \alpha_n)\phi(x^*, w_n) \\
&\quad - (1 - \alpha_n) \left(1 - \frac{\mu\beta_n}{\nu\beta_{n+1}}\right) (\phi(y_n, z_n) + \phi(z_n, w_n))
\end{aligned}$$

$$\begin{aligned} &\leq \alpha_n \phi(x^*, u) + (1 - \alpha_n)((1 - \theta_n)\phi(x^*, x_n) + \theta_n\phi(x^*, x_{n-1})) \\ &\quad - (1 - \alpha_n) \left(1 - \frac{\mu\beta_n}{\nu\beta_{n+1}}\right) (\phi(y_n, z_n) + \phi(z_n, w_n)). \end{aligned}$$

This implies that

$$\begin{aligned} (1 - \alpha_n) \left(1 - \frac{\mu\beta_n}{\nu\beta_{n+1}}\right) (\phi(y_n, z_n) + \phi(z_n, w_n)) &\leq \alpha_n \left(\phi(x^*, u) + \frac{\theta_n}{\alpha_n}\phi(x^*, x_{n-1})\right) \\ &\quad + (1 - \alpha_n)\phi(x^*, x_n) - \phi(x^*, x_{n+1}). \end{aligned}$$

By condition B2 and (3.15), we have $(\phi(y_n, z_n) + \phi(z_n, w_n)) \rightarrow 0$, as $n \rightarrow \infty$, thus

$$\lim_{n \rightarrow \infty} \phi(y_n, z_n) = \lim_{n \rightarrow \infty} \phi(z_n, w_n) = 0.$$

Since the sequences $\{y_n\}$, $\{z_n\}$ and $\{w_n\}$ are bounded, we obtain by Lemma 2.2, that

$$(3.19) \quad \lim_{n \rightarrow \infty} \|y_n - z_n\| = \lim_{n \rightarrow \infty} \|z_n - w_n\| = 0.$$

From Algorithm 3.4 and condition B4, we obtain

$$\lim_{n \rightarrow \infty} \phi(w_n, x_n) = \lim_{n \rightarrow \infty} \theta_n \phi(x_n, x_{n-1}) = 0,$$

and by Lemma 2.2, we get

$$(3.20) \quad \lim_{n \rightarrow \infty} \|w_n - x_n\| = 0.$$

It is easy to see from (3.19) and (3.20), that

$$(3.21) \quad \lim_{n \rightarrow \infty} \|x_n - z_n\| = \|x_n - y_n\| = 0.$$

Observe also that

$$(3.22) \quad \phi(y_n, p_n) = \phi(y_n, J^{-1}((1 - \gamma_n)Jy_n + \gamma_n)JSy_n) \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Hence, by Lemma 2.2, we obtain

$$\lim_{n \rightarrow \infty} \|y_n - p_n\| = 0.$$

This and (3.21), imply

$$\lim_{n \rightarrow \infty} \|x_n - p_n\| = 0.$$

Furthermore,

$$\|Jx_{n+1} - Jp_n\| = \alpha_n \|Ju - Jp_n\| = \alpha_n \|Ju - Jp_n\| \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Since J^{-1} is norm-to-norm continuous on bounded subsets of E , we have $\|x_{n+1} - p_n\| \rightarrow 0$, as $n \rightarrow \infty$. Hence,

$$(3.23) \quad \|x_{n+1} - x_n\| \leq \|x_{n+1} - p_n\| + \|p_n - x_n\| \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Now, since the sequence $\{x_n\}$ is bounded there exists a subsequence $\{x_{n_i}\}$ of $\{x_n\}$ such that $x_n \rightharpoonup q \in E$. Then, by (3.19), (3.20) and Lemma 3.4, we obtain $q \in \text{Sol}(C, f)$. Also, since $\|y_n - Sy_n\| \rightarrow 0$ and $\|x_n - y_n\| \rightarrow 0$ as $n \rightarrow \infty$, then we have $q \in \hat{F}(S) = F(S)$. Therefore, $q \in \Gamma$.

We now show that $\{x_n\}$ converges strongly to a point $x^* = \Pi_{\Gamma}u$. Let $\{x_{n_i}\}$ be a subsequence of $\{x_n\}$ such that $x_{n_i} \rightharpoonup q$ and

$$\limsup_{n \rightarrow \infty} \langle Ju - Jx^*, x_{n+1} - x^* \rangle = \lim_{i \rightarrow \infty} \langle Ju - Jx^*, x_{n_i+1} - x^* \rangle.$$

Since $\|x_{n+1} - x_n\| \rightarrow 0$ as $n \rightarrow \infty$, we have by (2.5), that

$$(3.24) \quad \limsup_{n \rightarrow \infty} \langle Ju - Jx^*, x_{n+1} - x^* \rangle = \lim_{i \rightarrow \infty} \langle Ju - Jx^*, x_{n_i+1} - x^* \rangle \\ = \langle Ju - Jx^*, q - x^* \rangle \leq 0.$$

It follows from Lemma 2.5, Lemma 3.5 (i) and (3.24), that $\phi(p, x_n) \rightarrow$ as $n \rightarrow \infty$. Therefore, by Lemma 2.2, we obtain

$$\lim_{n \rightarrow \infty} \|x_n - x^*\| = 0.$$

Case II Suppose there exists a subsequence $\{x_{n_j}\}$ of $\{x_n\}$ such that

$$\phi(x^*, x_{n_j+1}) > \phi(x^*, x_{n_j}), \quad \text{for all } n \in \mathbb{N}.$$

From Lemma 2.6, there exists a non-decreasing sequence $\{m_n\} \subset \mathbb{N}$ such that $m_n \rightarrow \infty$ and the following inequalities hold for all $n \in \mathbb{N}$:

$$(3.25) \quad \phi(x^*, x_{m_n}) \leq \phi(x^*, x_{m_n+1}) \quad \text{and} \quad \phi(p, x_n) \leq \phi(x^*, x_{m_n+1}).$$

We note from (3.11) and (3.12), that

$$\begin{aligned} \phi(x^*, x_{m_n}) &\leq \phi(x^*, x_{m_n+1}) \leq \alpha_{m_n} \phi(x^*, u) \\ &\quad + (1 - \alpha_{m_n}) \left[\phi(x^*, w_{m_n}) - \left(1 - \frac{\mu\beta_{m_n}}{\nu\beta_{m_n+1}}\right) (\phi(y_{m_n}, z_{m_n}) + \phi(z_{m_n}, w_{m_n})) \right] \\ &\leq \alpha_{m_n} \phi(x^*, u) + (1 - \alpha_{m_n}) ((1 - \theta_{m_n})\phi(x^*, x_{m_n}) + \theta_{m_n}\phi(x^*, x_{m_n-1})) \\ &\quad - (1 - \alpha_{m_n}) \left(1 - \frac{\mu\beta_{m_n}}{\nu\beta_{m_n+1}}\right) (\phi(y_{m_n}, z_{m_n}) + \phi(z_{m_n}, w_{m_n})) \\ &\leq \alpha_{m_n} \left(\phi(x^*, u) + \frac{\theta_{m_n}}{\alpha_{m_n}} \phi(x^*, x_{m_n-1}) \right) \\ &\quad - (1 - \alpha_{m_n}) \left(1 - \frac{\mu\beta_{m_n}}{\nu\beta_{m_n+1}}\right) (\phi(y_{m_n}, z_{m_n}) + \phi(z_{m_n}, w_{m_n})). \end{aligned}$$

Hence,

$$\begin{aligned} &(1 - \alpha_{m_n}) \left(1 - \frac{\mu\beta_{m_n}}{\nu\beta_{m_n+1}}\right) \times (\phi(y_{m_n}, z_{m_n}) + \phi(z_{m_n}, w_{m_n})) \\ &\leq \alpha_{m_n} \left(\phi(x^*, u) + \frac{\theta_{m_n}}{\alpha_{m_n}} \phi(x^*, x_{m_n-1}) \right) + (1 - \alpha_{m_n})\phi(x^*, x_{m_n}) - \phi(x^*, x_{m_n}). \end{aligned}$$

Since $\alpha_{m_n} \rightarrow 0$ as $n \rightarrow \infty$, it follows that

$$\left(1 - \frac{\mu\beta_{m_n}}{\nu\beta_{m_n+1}}\right) (\phi(y_{m_n}, z_{m_n}) + \phi(z_{m_n}, w_{m_n})) \rightarrow 0, \quad \text{as } n \rightarrow \infty,$$

hence

$$\lim_{n \rightarrow \infty} \phi(y_{m_n}, z_{m_n}) = \lim_{n \rightarrow \infty} \phi(z_{m_n}, w_{m_n}) = 0.$$

Since $\{x_{m_n}\}$, $\{y_{m_n}\}$ and $\{w_{m_n}\}$ are bounded, we have

$$\lim_{n \rightarrow \infty} \|y_{m_n} - z_{m_n}\| = \lim_{n \rightarrow \infty} \|z_{m_n} - w_{m_n}\| = 0.$$

Following similar method as in Case I, we obtain

$$(3.26) \quad \lim_{n \rightarrow \infty} \|w_{m_n} - Sw_{m_n}\| = \lim_{n \rightarrow \infty} \|x_{m_{n+1}} - x_{m_n}\| = 0.$$

By Lemma 3.4 and (3.26), we obtain a weak limit $q \in E$ of $\{x_{m_n}\}$ such that $q \in \Gamma$.

Again, since $\{x_{m_n}\}$ is bounded, we can choose a sequence $\{x_{m_n}\}$ of $\{x_{m_n}\}$, subsequence if necessary such that $x_{m_n} \rightarrow q$ as $n \rightarrow \infty$ and

$$\limsup_{n \rightarrow \infty} \langle Ju - Jx^*, x_{m_{n+1}} - x^* \rangle = \lim_{n \rightarrow \infty} \langle Ju - Jx^*, x_{m_{n+1}} - x^* \rangle.$$

Hence, from (2.5), we have

$$(3.27) \quad \begin{aligned} \limsup_{n \rightarrow \infty} \langle Ju - Jx^*, x_{m_{n+1}} - x^* \rangle &= \lim_{n \rightarrow \infty} \langle Ju - Jx^*, x_{m_{n+1}} - x^* \rangle \\ &\leq \langle Ju - Jx^*, q - x^* \rangle \leq 0. \end{aligned}$$

From (3.25), we have

$$\begin{aligned} 0 &\leq \phi(x^*, x_{m_{n+1}}) - \phi(x^*, x_{m_n}) \\ &\leq (1 - \alpha_{m_n})\phi(x^*, x_{m_n}) \\ &\quad + \alpha_{m_n} \left(\frac{\theta_{m_n}}{\alpha_{m_n}} \phi(x^*, x_{m_{n-1}}) + 2\langle Ju - Jx^*, x_{m_{n+1}} - x^* \rangle \right) - \phi(x^*, x_{m_n}). \end{aligned}$$

That is

$$(3.28) \quad \phi(x^*, x_{m_n}) \leq \frac{\theta_{m_n}}{\alpha_{m_n}} \phi(x^*, x_{m_{n-1}}) + 2\langle Ju - Jx^*, x_{m_{n+1}} - x^* \rangle.$$

Hence, by condition (B5) and (3.27), we obtain $\phi(x^*, x_{m_n}) \rightarrow 0$ as $n \rightarrow \infty$ and Lemma 2.2 implies $\|x_{m_n} - x^*\| \rightarrow 0$ as $n \rightarrow \infty$. Consequently, $\|x_n - x^*\| \rightarrow 0$ as $n \rightarrow \infty$. Therefore, the sequence $\{x_n\}$ converges strongly to $x^* = \Pi_{\Gamma}u$. \square

4. APPLICATIONS

In this section, we present some theoretical applications of our main result.

4.1. Variational Inequalities Problem. Suppose we define the f in $EP(C, f)$ (1.1), by:

$$(4.1) \quad f(x, y) := \begin{cases} \langle Ax, y - x \rangle, & \text{if } x, y \in C, \\ +\infty, & \text{otherwise,} \end{cases}$$

where $A : C \rightarrow E^*$ is a strongly γ -pseudomonotone mapping. Then $EP(C, f)$ (1.1) reduces to $VIP(C, A)$, that is to find $x^* \in C$ such that

$$(4.2) \quad \langle Ax^*, y - x^* \rangle \geq 0, \quad \text{for all } y \in C.$$

We denote the set of solution of (4.2) by $Sol(C, A)$. Recall an operator A is said to be strongly γ -pseudomonotone, if there exists $\gamma > 0$ such that for any $x, y \in C$

$$\langle Ax, y - x \rangle \geq 0 \Rightarrow \langle Ay, y - x \rangle \geq \gamma\phi(y, x).$$

In this situation, Algorithm 3.4 when modified provides a new method for solving variational inequality problems and fixed point problem for a quasi- ϕ -nonexpansive mapping. We give the new method as follows.

Algorithm 4.1. (TRSSEM) for $VIP(C, A)$

Step 0. Choose the sequences $\{\theta_n\}$, $\{\alpha_n\}$ and $\{\gamma_n\} \subset (0, 1)$ satisfying Assumption 3.3, take $\eta, \rho \in (0, 1)$ and $\beta_0 > 0$. For $u \in C$, select initial points x_0 and x_1 in C . Set $n = 1$.

Step 1. For $i = 1, 2, \dots, m$, and given the current iterate w_n , construct the family of half spaces

$$C_n^i := \{z \in E : h_i(w_n) + \langle h_i'(w_n), z - w_n \rangle \leq 0\}$$

and set

$$C_n = \bigcap_{i=1}^m C_n^i.$$

Let $w_n := J^{-1}(Jx_n + \theta_n(Jx_{n-1} - Jx_n))$. Compute

$$(4.3) \quad z_n = \Pi_{C_n} J^{-1}(Jw_n - \beta_n Aw_n),$$

where β_n is given by

$$(4.4) \quad \beta_{n+1} = \begin{cases} \min \left\{ \beta_n, \frac{\mu \|w_n - z_n\|}{\|g(w_n) - g(z_n)\|} \right\}, & \text{if } g(w_n) \neq g(z_n), \\ \beta_n, & \text{otherwise.} \end{cases}$$

Step 2. If $w_n = z_n$ ($w_n \in Sol(C, A)$), then set $w_n = y_n$ and go to Step 3. Otherwise, compute the next iterate by

$$(4.5) \quad y_n = \Pi_{Q_n} J^{-1}(Jw_n - \beta_n Az_n),$$

where

$$Q_n = \{w \in E : \langle w - z_n, Jw_n - \beta_n Aw_n - Jz_n \rangle \leq 0\}.$$

Step 3. Compute

$$(4.6) \quad x_{n+1} = J^{-1}((1 - \alpha_n)Ju + \alpha_n(1 - \gamma_n)Jy_n + \gamma_n JSy_n).$$

Step 4. Set $n := n + 1$ and go to Step 1.

A convergence result for solving $VIP(C, A)$ (4.2) is given below without proof.

Theorem 4.2. *Let C be a nonempty, closed and convex subset of a 2-uniformly convex and uniformly smooth real Banach space E and $h_i : E \rightarrow \mathbb{R}$ be a family of convex, weakly lower semicontinuous and Gâteaux differentiable functions, for $i = 1, 2, \dots, m$. Let $A : C \rightarrow E^*$ be a strongly γ -pseudomonotone operator that is bounded on bounded sets, let $S : E \rightarrow E$ be a quasi- ϕ -nonexpansive mapping such that $\Gamma = \{Sol(C, A) \cap F(S)\} \neq \emptyset$. Let $\{\theta_n\}$, $\{\beta_n\}$ and $\{\alpha_n\}$ be sequences in $(0, 1)$ satisfying Assumption 3.3, then the sequence $\{x_n\}$ generated by Algorithm 4.1 converges strongly to $p = \Pi_\Gamma u$, where Π_Γ is the projection of C onto Γ .*

4.2. Fixed Point Problem (FPP). Given a closed set $C \subset E$, a fixed point of a mapping $T : C \rightarrow C$ is any point $x^* \in C$ such that $x^* = Tx^*$. Finding a fixed point amounts to solving $EP(C, f)$ with

$$f(x, y) = \langle x - Tx, y - x \rangle, \quad \text{for all } y \in C.$$

In this case, we define the operator $T = I - A$, where I is the identity mapping on C and A is the operator defined in Subsection 4.1. The method and result given in 4.1, thus apply.

5. CONCLUSION

We considered an iterative approximation of a common solution of EP and FPP. We introduced a totally relaxed self adaptive inertial subgradient extragradient method, Mann and Halpern iterative technique for solving this problem in 2-uniformly convex Banach space, which is also uniformly smooth. Our method uses a carefully selected adaptive stepsize which does not depend on any Lipschitz-type condition neither does it require the knowledge of the Lipschitz constant of the gradient of pseudomonotone operator.

Acknowledgements. The authors sincerely thank the anonymous referee for his careful reading, constructive comments and fruitful suggestions that improved the manuscript. The first and second authors acknowledge with thanks the bursary and financial support from Department of Science and Innovation and National Research Foundation, Republic of South Africa Center of Excellence in Mathematical and Statistical Sciences (DSI-NRF COE-MaSS) Doctoral Bursary. The third author is supported by the National Research Foundation (NRF) of South Africa Incentive Funding for Rated Researchers (Grant Number 119903). Opinions expressed and conclusions arrived are those of the authors and are not necessarily to be attributed to the CoE-MaSS and NRF.

REFERENCES

- [1] H. A. Abass, C. Izuchukwu, O. T. Mewomo and Q. L. Dong, *Strong convergence of an inertial forward-backward splitting method for accretive operators in real Banach space*, Fixed Point Theory **21**(2) (2020), 397–411. <https://doi.org/10.24193/fpt-ro.2020.2.28>

- [2] R. P. Agarwal, D. O'Regan and D. R. Saha, *Fixed Point Theory for Lipschitzian-Type Mappings with Applications*, Springer, New York, 2009. <https://link.springer.com/book/10.1007/978-0-387-75818-3>
- [3] T. O. Alakoya, L. O. Jolaoso and O. T. Mewomo, *Strong convergence theorems for finite families of pseudomonotone equilibrium and fixed point problems in Banach spaces*, Afr. Mat. **32** (2021), 897–923. <https://doi.org/10.1007/s13370-020-00869-z>
- [4] T. O. Alakoya and O. T. Mewomo, *Viscosity S -iteration method with inertial technique and self-adaptive step size for split variational inclusion, equilibrium and fixed point problems*, Comput. Appl. Math. (2021), Article ID 39. <https://doi.org/10.1007/s40314-021-01749-3>
- [5] T. O. Alakoya, A. O. E. Owolabi and O. T. Mewomo, *An inertial algorithm with a self-adaptive step size for a split equilibrium problem and a fixed point problem of an infinite family of strict pseudo-contractions*, J. Nonlinear Var. Anal. **5** (2021), 803–829. <https://doi.org/10.1080/02331934.2021.1895154>
- [6] T. O. Alakoya, A. Taiwo and O. T. Mewomo, *On system of split generalised mixed equilibrium and fixed point problems for multivalued mappings with no prior knowledge of operator norm*, Fixed Point Theory **23**(1) (2022), 45–74. <https://doi.org/10.24193/fpt-ro.2022.1.04>
- [7] Y. I. Alber, *Metric and generalized projection operators in Banach spaces: properties and applications*, in: A. G. Kartsatos (Ed.), *Theory and Applications of Nonlinear Operators and Accretive and Monotone Type*, Lecture Notes in Pure and Applied Mathematics **178**, Dekker, New York, 1996, 15–50.
- [8] P. N. Anh, *A hybrid extragradient method extended to fixed point problems and equilibrium problems*, Optimization **62** (2013), 271–283. <https://doi.org/10.1080/02331934.2011.607497>
- [9] P. N. Anh and H. A. Le Thi, *An Armijo-type method for pseudomonotone equilibrium problems and its applications*, J. Glob. Optim. **57** (2013), 803–820. <https://doi.org/10.1007/s10898-012-9970-8>
- [10] P. N. Anh, *Strong convergence theorems for nonexpansive mappings and Ky Fan inequalities*, J. Optim. Theory Appl. **154** (2012), 303–320. <https://doi.org/10.1007/s10957-012-0005-x>
- [11] H. H. Bauschke and J. M. Borwein, *On projection algorithms for solving convex feasibility problems*, SIAM Rev. **38**(3) (1996), 367–426. <https://doi.org/10.1137/S0036144593251710>
- [12] J. Y. Bello Cruz, P. S. M. Santos and S. Scheimberg, *A two-phase algorithm for a variational inequality formulation of equilibrium problems*, J. Optim. Theory Appl. **159** (2013), 562–575. <https://doi.org/10.1007/s10957-012-0181-8>
- [13] G. Bigi, M. Castellani, M. Pappalardo and M. Passacantando, *Existence and solution methods for equilibria*, Eur. J. Oper. Res. **227** (2013), 1–11. <https://doi.org/10.1016/j.ejor.2012.11.037>
- [14] E. Blum and W. Oettli, *From optimization and variational inequalities to equilibrium problems*, Math. Student **63** (1994), 123–145.
- [15] Y. Censor and A. Lent, *An iterative row-action method for interval complex programming*, J. Optim. Theory Appl. **34** (1981), 321–353. <https://doi.org/10.1007/BF00934676>
- [16] Y. Censor and S. Reich, *Iterations of paracontractions and firmly nonexpansive operators with applications to feasibility and optimization*, Optimization **37** (1996), 323–339. <https://doi.org/10.1080/02331939608844225>
- [17] Y. Censor, A. Gibali and S. Reich, *The subgradient extragradient method for solving variational inequalities in Hilbert spaces*, J. Optim. Theory Appl. **148** (2011), 318–335. <https://doi.org/10.1007/s10957-010-9757-3>
- [18] I. Cioranescu, *Geometry of Banach Spaces, Duality Mappings and Nonlinear*, Kluwer, Dordrecht, 1990.
- [19] P. L. Combettes and S. A. Hirstoaga, *Equilibrium programming in Hilbert spaces*, J. Nonlinear Convex Anal. **6** (2005), 117–136.

- [20] P. Debnath, N. Konwar and S. Radenović, *Metric Fixed Point Theory: Applications in Science, Engineering and Behavioural Science*, Springer Verlag, Singapore, 2021. <https://doi.org/10.1007/978-981-16-4896-0>
- [21] P. M. Duc, L. D. Muu and N. V. Quy, *Solution-existence and algorithms with their convergence rate for strongly pseudomonotone equilibrium problems*, Pacific J. Optim. **12** (2016), 833–845.
- [22] M. Fukushima, *A relaxed projection method for variational inequalities*, Math. Program **35** (1986), 58–70. <https://doi.org/10.1007/BF01589441>
- [23] S. He, T. Wu, A. Gibali and Q.-L. Dong, *Totally self relaxed self-adaptive algorithm for solving variational inequalities over the intersection of sub-level sets*, Optimization **67**(9) (2018), 1487–1504. <https://doi.org/10.1080/02331934.2018.1476515>
- [24] D. V. Hieu, *New extragradient method for a class of equilibrium problems*, App. Anal. **97**(5) (2018), 811–824. <https://doi.org/10.1080/00036811.2017.1292350>
- [25] C. Izuchukwu, K. O. Aremu, O. K. Oyewole, O. T. Mewomo and S. H. Khan, *On mixed equilibrium problems in Hadamard spaces*, J. Math. **2019** (2019), Article ID 3210649, 13 pages. <https://doi.org/10.1155/2019/3210649>
- [26] Z. Jouymandi and F. Moradlou, *Extragradient and linesearch algorithms for solving equilibrium problems and fixed point problems in Banach space*, Fixed Point Theory **20**(2) (2019), 523–539. <https://doi.org/10.24193/fpt-ro.2019.2.34>
- [27] Z. Jouymandi and F. Moradlou, *Retraction algorithms for solving variational inequalities, pseudomonotone equilibrium problems and fixed point problems in Banach spaces*, Numer. Algorithms **78**(2) (2018), 1153–1182. <https://doi.org/10.1007/s11075-017-0417-7>
- [28] S. Kamimura and W. Takahashi, *Strong convergence of a proximal-type algorithm in a Banach space*, SIAM J. Optim. **13** (2002), 938–945. <https://doi.org/10.1137/S105262340139611X>
- [29] G. Kassay, M. Miholca and N. T. Vinh, *Vector quasi-equilibrium problems for the sum of two multivalued mappings*, J. Optim. Theory Appl. **169** (2016), 424–442. <https://doi.org/10.1007/s10957-016-0919-9>
- [30] S. H. Khan, T. O. Alakoya and O. T. Mewomo, *Relaxed projection methods with self-adaptive step size for solving variational inequality and fixed point problems for an infinite family of multivalued relatively nonexpansive mappings in Banach spaces*, Math. Comput. Appl. **25** (2020), Article ID 54, 25 pages. <https://doi.org/10.3390/mca25030054>
- [31] P. E. Maingé, *Convergence theorem for inertial KM-type algorithms*, J. Comput. Appl. Math. **219** (2008), 223–236. <https://doi.org/10.1016/j.cam.2007.07.021>
- [32] P. E. Maingé, *Strong convergence of projected subgradient methods for nonsmooth and nonstrictly convex minimization*, Set-Valued Var. Anal. **16** (2008), 899–912. <https://doi.org/10.1007/s11228-008-0102-z>
- [33] O. T. Mewomo and F. U. Ogbuisi, *Convergence analysis of iterative method for multiple set split feasibility problems in certain Banach spaces*, Quaest. Math. **41**(1) (2018), 129–148. <https://doi.org/10.2989/16073606.2017.1375569>
- [34] O. T. Mewomo and O. K. Oyewole, *An iterative approximation of common solutions of split generalized vector mixed equilibrium problem and some certain optimization problems*, Demonstr. Math. **54**(1) (2021), 335–358. <https://doi.org/10.1515/dema-2021-0019>
- [35] K. Nakajo, *Strong convergence for gradient projection method and relatively nonexpansive mappings in Banach spaces*, Appl. Math. Comput. **271** (2015), 251–258. <https://doi.org/10.1016/j.amc.2015.08.096>
- [36] G. N. Ogwo, T. O. Alakoya and O. T. Mewomo, *Iterative algorithm with self-adaptive step size for approximating the common solution of variational inequality and fixed point problems*, Optimization (2021). <https://doi.org/10.1080/02331934.2021.1981897>
- [37] G. N. Ogwo, T. O. Alakoya and O. T. Mewomo, *Inertial iterative method with self-adaptive step size for finite family of split monotone variational inclusion and fixed point problems in Banach spaces*, Demonstr. Math. (2021). <https://doi.org/10.1515/dema-2020-0119>

- [38] G. N. Ogwo, C. Izuchukwu, Y. Shehu and O. T. Mewomo, *Convergence of relaxed inertial sub-gradient extragradient methods for quasimonotone variational inequality problems*, J. Sci. Comput. (2021). <https://doi.org/10.1007/s10915-021-01670-1>
- [39] M. A. Olona, T. O. Alakoya, A. O.-E. Owolabi and O. T. Mewomo, *Inertial shrinking projection algorithm with self-adaptive step size for split generalized equilibrium and fixed point problems for a countable family of nonexpansive multivalued mappings*, Demonstr. Math. **54** (2021), 47–67. <https://doi.org/10.1515/dema-2021-0006>
- [40] M. A. Olona, T. O. Alakoya, A. O.-E. Owolabi and O. T. Mewomo, *Inertial algorithm for solving equilibrium, variational inclusion and fixed point problems for an infinite family of strictly pseudocontractive mappings*, J. Nonlinear Funct. Anal. **2021** (2021), Article ID 10, 21 pages. <https://doi.org/10.23952/jnfa.2020.10>
- [41] A. O.-E. Owolabi, T. O. Alakoya, A. Taiwo and O. T. Mewomo, *A new inertial-projection algorithm for approximating common solution of variational inequality and fixed point problems of multivalued mappings*, Numer. Algebra Control Optim. (2021). <https://doi.org/10.3934/naco.2021004>
- [42] O. K. Oyewole, H. A. Abass and O. T. Mewomo, *Strong convergence algorithm for a fixed point constraint split null point problem*, Rend. Circ. Mat. Palermo (2) **70**(1) (2021), 389–408. <https://doi.org/10.1007/s12215-020-00505-6>
- [43] O. K. Oyewole, K. O. Aremu and O. T. Mewomo, *A multi step inertial algorithm for approximating a common solution of split generalized mixed equilibrium and minimization problems*, Ric. Mat. (2021). <https://doi.org/10.1007/s11587-021-00624-x>
- [44] O. K. Oyewole, O. T. Mewomo, L. O. Jolaoso and S. H. Khan, *An extragradient algorithm for split generalized equilibrium problem and the set of fixed points of quasi- ϕ -nonexpansive mappings in Banach spaces*, Turkish J. Math. **44**(4) (2020), 1146–1170. <https://1146\T1\textendash1170>. doi:10.3906/mat-1911-83
- [45] S. Reich, *A weak convergence theorem for alternating method with Bregman distance*, in: A. G. Kartsatos (Ed.), *Theory and Applications of Nonlinear Operators of Accretive and Monotone Type*, Marcel Dekker, New York, 1996, 313–318.
- [46] S. Saewan and P. Kumam, *A generalized f -projection method for countable families of weak relatively nonexpansive mappings and the system of generalized Ky Fan inequalities*, J. Glob. Optim. **56** (2012), 1–23. <https://doi.org/10.1007/s10898-012-9922-3>
- [47] G. Stampacchia, *Forms bilinéaires coercitives sur les ensembles convexes*, C. R. Acad. Sci. Paris **258** (1964), 4414–4416.
- [48] A. Taiwo, T. O. Alakoya and O. T. Mewomo, *Strong convergence theorem for solving equilibrium problem and fixed point of relatively nonexpansive multi-valued mappings in a Banach space with applications*, Asian-Eur. J. Math. **14**(8) (2021), Article ID 2150137, 31 pages. <https://doi.org/10.1142/S1793557121501370>
- [49] V. Todorcević, *Harmonic Quasiconformal Mappings and Hyperbolic Type Metrics*, Springer Nature, Switzerland AG, 2019. <https://doi.org/10.1007/978-3-030-22591-9>
- [50] R. T. Rockfellar, *Monotone operators and the proximal point algorithm*, SIAM J. Control Optim. **14** (1977), 877–808. <https://doi.org/10.1137/0314056>
- [51] G. C. Ugwunnadi, C. Izuchukwu and O. T. Mewomo, *On nonspreading-type mappings in Hadamard spaces*, Bol. Soc. Parana. Mat. (3) **39**(5) (2021), 175–197. <https://doi.org/10.5269/bspm.41768>
- [52] V. A. Uzor, T. O. Alakoya and O. T. Mewomo, *Strong convergence of a self-adaptive inertial Tseng’s extragradient method for pseudomonotone variational inequalities and fixed point problems*, Open Math. (2022). <https://doi.org/10.1515/math-2022-0429>
- [53] N. T. Vinh and A. Gibali, *Gradient projection-type algorithms for solving equilibrium problems and its applications*, Comput. Appl. Math. **38** (2019), Article ID 119. <https://doi.org/10.1007/240314-019-0894-5>

- [54] H. K. Xu, *Another control condition in an iterative method for nonexpansive mappings*, Bull. Aust. Math. Soc. **65** (2002), 109–113. <https://doi.org/10.1017/S0004972700020116>
- [55] H. K. Xu, *Inequalities in Banach spaces with applications*, Nonlinear Anal. **16** (1991), 1127–1138. [https://doi.org/10.1016/0362-546X\(91\)90200-K](https://doi.org/10.1016/0362-546X(91)90200-K)
- [56] J. Yang and H. W. Liu, *Strong convergence result for solving monotone variational inequalities in Hilbert space*, Numer. Algorithms **80** (2019), 741–752. <https://doi.org/10.1007/s11075-018-0504-4>
- [57] F. Yang, L. Zhao and J. K. Kim, *Hybrid projection method for generalized mixed equilibrium problem and fixed point problem of infinite family of asymptotically quasi- ϕ -nonexpansive mappings in Banach spaces*, Appl. Math. Comput. **218**(10) (2012), 6072–6082. <https://doi.org/10.1016/j.amc.2011.11.091>

¹SCHOOL OF MATHEMATICS, STATISTICS AND COMPUTER SCIENCE,
UNIVERSITY OF KWAZULU-NATAL,
DURBAN, SOUTH AFRICA

²DSI-NRF CENTER OF EXCELLENCE IN MATHEMATICAL AND STATISTICAL SCIENCES,
(COE-MASS) JOHANNESBURG, SOUTH AFRICA
Email address: ¹217079141@stu.ukzn.ac.za
Email address: ²216075727@stu.ukzn.ac.za
Email address: ³mewomoo@ukzn.ac.za