

## ON THE REVERSE MINKOWSKI'S INTEGRAL INEQUALITY

BOUHARKET BENAÏSSA<sup>1,2</sup>

**ABSTRACT.** The aim of this work is to obtain the reverse Minkowski integral inequality. For this aim, we first give a proposition which is important for our main results. Then we establish some reverse Minkowski integral inequalities for parameters  $0 < p < 1$  and  $p < 0$ , respectively.

### 1. INTRODUCTION

In recent years, inequalities are playing a very significant role in all fields of mathematics and present a very active and attractive field of research. As example, let us cite the field of integration which is dominated by inequalities involving functions and their integrals ([2, 3]). One of the famous integral inequalities is Minkowski's integral inequality. In particular the following statement was proved for  $p \geq 1$  (for details to see [1]).

**Theorem 1.1.** *Let  $1 \leq p \leq +\infty$ ,  $\Omega \subset \mathbb{R}^n$  and  $A \subset \mathbb{R}^m$  be a measurable sets. Suppose that  $f$  is measurable on  $\Omega \times A$  and  $f(\cdot, y) \in L_p(\Omega)$  for almost all  $y \in A$ . Then*

$$(1.1) \quad \left\| \int_A f(\cdot, y) dy \right\|_{L_p(\Omega)} \leq \int_A \|f(\cdot, y)\|_{L_p(\Omega)} dy,$$

*if the right-hand side is finite.*

*Remark 1.1.* If  $0 < p < 1$ ,  $\text{mes } A > 0$  and  $\text{mes } \Omega > 0$  inequality (1.1) is not valid (to see [1]).

In this paper we obtain some integral inequalities which are reverse versions of the inequality (1.1).

---

*Key words and phrases.* Hölder's inequality, Minkowski's integral inequality.  
*2010 Mathematics Subject Classification.* Primary: 26D15. Secondary: 26D10.  
*Received:* July, 04, 2019.  
*Accepted:* January 04, 2020.

## 2. PRELIMINARIES

**2.1. Reverse Young's and Holder's Inequalities.** The following inequalities are well-known Young inequalities. Let  $a > 0$ ,  $b > 0$  and  $\frac{1}{p} + \frac{1}{p'} = 1$ , then

$$(2.1) \quad ab \leq \frac{a^p}{p} + \frac{b^{p'}}{p'}, \quad \text{for } p \geq 1,$$

$$(2.2) \quad ab \geq \frac{a^p}{p} + \frac{b^{p'}}{p'}, \quad \text{for } 0 < p < 1.$$

**Corollary 2.1** (Reverse Young's inequality). *Let  $a > 0$ ,  $b > 0$  and  $\frac{1}{p} + \frac{1}{p'} = 1$ , then*

$$(2.3) \quad ab \geq \frac{a^p}{p} + \frac{b^{p'}}{p'}, \quad \text{for } p < 0.$$

*Proof.* We have  $\frac{p'-1}{p'} = \frac{1}{p}$ ,  $(p-1)(p'-1) = 1$  and inequality (2.3) is equivalent to

$$\frac{a^{p-1}}{bp} + \frac{b^{p'-1}}{ap'} \leq 1.$$

We take  $t = \frac{a^{p-1}}{b}$ , then

$$\frac{b^{p'-1}}{ap'} = \frac{a^{(p-1)(p'-1)}}{t^{(p'-1)ap'}} = \frac{1}{t^{(p'-1)p'}} = \frac{t^{-(p'-1)}}{p'}.$$

We obtain

$$\frac{a^{p-1}}{bp} + \frac{b^{p'-1}}{ap'} = \frac{t}{p} + \frac{t^{-(p'-1)}}{p'} = f(t), \quad t > 0.$$

For all  $t > 0$ , we have

$$f'(t) = \frac{1}{p} - \frac{p'-1}{p'} t^{-p'} = \frac{1}{p} - \frac{1}{p} t^{-p'} = \frac{1}{p} (1 - t^{-p'}),$$

for all  $p < 0$  and  $0 < p' < 1$ , we get

$$f'(t) = 0 \Leftrightarrow 1 - t^{-p'} = 0 \Leftrightarrow t = 1,$$

$$f'(t) > 0 \Leftrightarrow 1 - t^{-p'} < 0 \Leftrightarrow 0 < t < 1.$$

Hence, the function  $f$  is majored with  $f(1) = 1$  for all  $t \in (0, \infty)$ .

We deduce that

$$\frac{a^{p-1}}{bp} + \frac{b^{p'-1}}{ap'} \leq 1 \Leftrightarrow ab \geq \frac{a^p}{p} + \frac{b^{p'}}{p'}, \quad \text{for } p < 0. \quad \square$$

**Corollary 2.2** (Reverse Hölder's inequality). *Let  $\Omega \subset \mathbb{R}^n$  be a measurable set and  $p < 0$ , we suppose that  $f, g$  are measurable on  $\Omega$ .*

*If  $f \in L_p(\Omega)$  and  $g \in L_{p'}(\Omega)$  ( $p'$  is the conjugate parameter), then*

$$(2.4) \quad \int_{\Omega} |fg| dt \geq \|f\|_{L_p} \|g\|_{L_{p'}}.$$

*Proof.* Choose  $a = \frac{|f|}{\|f\|_{L_p}}$ ,  $b = \frac{|g|}{\|g\|_{L_{p'}}}$  and by using reverse Young's inequality (2.3), we write

$$\frac{|fg|}{\|f\|_{L_p} \cdot \|g\|_{L_{p'}}} \geq \frac{|f|^p}{p\|f\|_{L_p}^p} + \frac{|g|^{p'}}{p'\|g\|_{L_{p'}}^{p'}},$$

by integrand the above inequality we obtain

$$\int_{\Omega} \frac{|f(t)g(t)|}{\|f\|_{L_p} \cdot \|g\|_{L_{p'}}} dt \geq \int_{\Omega} \frac{|f(t)|^p}{p\|f\|_{L_p}^p} dt + \int_{\Omega} \frac{|g(t)|^{p'}}{p'\|g\|_{L_{p'}}^{p'}} dt = 1,$$

and thus

$$\int_{\Omega} |f(t)g(t)| dt \geq \|f\|_{L_p} \|g\|_{L_{p'}}, \quad \text{for } p < 0. \quad \square$$

*Remark 2.1.* We can write

$$\int_{\Omega} |f(t)g(t)| dt \geq \left( \int_{\Omega} |f(t)|^p dt \right)^{\frac{1}{p}} \left( \int_{\Omega} |g(t)|^{p'} dt \right)^{\frac{1}{p'}},$$

hence

$$\left( \int_{\Omega} |f(t)g(t)| dt \right)^p \leq \left( \int_{\Omega} |f(t)|^p dt \right) \left( \int_{\Omega} |g(t)|^{p'} dt \right)^{p-1}$$

(see [4]).

Now we give a proposition which will be used frequently in the proof of main theorems.

Let  $-\infty < a < b < +\infty$  and  $-\infty < c < d < +\infty$  and we defined the set  $\mathbb{E}$  by

$$\mathbb{E} = \{f \mid f : (a, b) \times (c, d) \rightarrow \mathbb{R}, f \geq 0 \text{ or } f \leq 0\}.$$

Suppose  $H : (a, b) \times (c, d) \rightarrow \mathbb{C}$  a measurable function defined by

$$H(x, y) = f_1(x, y) + i f_2(x, y),$$

where  $f_1, f_2 \in \mathbb{E}$ .

**Proposition 2.1.** (i) If  $f_1 = 0$  or  $f_2 = 0$ , then

$$(2.5) \quad \left| \int_c^d |H(x, y)| dy \right| = \left| \int_c^d H(x, y) dy \right|.$$

(ii) If  $f_1 \neq 0$  and  $f_2 \neq 0$ , then

$$(2.6) \quad \left| \int_c^d |H(x, y)| dy \right| \leq \sqrt{2} \left| \int_c^d H(x, y) dy \right|.$$

*Proof.* (i) If  $f_2 = 0$ , then

$$\left| \int_c^d |H(x, y)| dy \right| = \left| \int_c^d |f_1(x, y)| dy \right| = \left| \int_c^d f_1(x, y) dy \right| = \left| \int_c^d H(x, y) dy \right|.$$

If  $f_1 = 0$ , then

$$\left| \int_c^d |H(x, y)| dy \right| = \left| \int_c^d |i f_2(x, y)| dy \right| = \left| \int_c^d |f_2(x, y)| dy \right|$$

$$\begin{aligned}
&= \left| \int_c^d f_2(x, y) dy \right| = \left| \int_c^d i f_2(x, y) dy \right| \\
&= \left| \int_c^d H(x, y) dy \right|.
\end{aligned}$$

(ii) If  $f_1 \neq 0$  and  $f_2 \neq 0$ , then

$$\begin{aligned}
\left| \int_c^d |H(x, y)| dy \right|^2 &= \left| \int_c^d [f_1^2(x, y) + f_2^2(x, y)]^{\frac{1}{2}} dy \right|^2 \\
&= \left( \int_c^d |f_1^2 + f_2^2|^{\frac{1}{2}}(x, y) dy \right)^2 \\
&= \|f_1^2 + f_2^2\|_{L_p(c, d)}, \quad \text{with } p = \frac{1}{2}, \\
\left| \int_c^d H(x, y) dy \right|^2 &= \left| \int_c^d f_1(x, y) dy + i \int_c^d f_2(x, y) dy \right|^2 \\
&= \left( \int_c^d f_1(x, y) dy \right)^2 + \left( \int_c^d f_2(x, y) dy \right)^2 \\
&= \left( \int_c^d |f_1(x, y)| dy \right)^2 + \left( \int_c^d |f_2(x, y)| dy \right)^2 \\
&= \|f_1^2\|_{L_p(c, d)} + \|f_2^2\|_{L_p(c, d)}, \quad \text{with } p = \frac{1}{2}.
\end{aligned}$$

For all  $0 < p < 1$  we have

$$\|f_1^2 + f_2^2\|_{L_p(c, d)} \leq 2^{\frac{1}{p}-1} \left( \|f_1^2\|_{L_p(c, d)} + \|f_2^2\|_{L_p(c, d)} \right),$$

for  $p = \frac{1}{2}$  we obtain

$$\left| \int_c^d |H(x, y)| dy \right|^2 \leq 2 \left| \int_c^d H(x, y) dy \right|^2.$$

Then

$$\left| \int_c^d |H(x, y)| dy \right| \leq \sqrt{2} \left| \int_c^d H(x, y) dy \right|. \quad \square$$

In this work we consider the reverse inequality of (1.1), with  $0 < p < 1$  and  $p < 0$  for  $f : (a, b) \times (c, d) \rightarrow \mathbb{K}$ , with  $\mathbb{K}$  is  $\mathbb{C}$ ,  $\mathbb{E}$  or  $i\mathbb{E}$ .

### 3. MAIN RESULTS

In this section we obtain some reverse Minkowski type inequalities.

**Theorem 3.1.** *Let  $0 < p < 1$ ,  $-\infty < a < b < +\infty$  and  $-\infty < c < d < +\infty$ . Suppose that  $H : (a, b) \times (c, d) \rightarrow \mathbb{C}$  is measurable with  $\operatorname{Re}(H), \operatorname{Im}(H) \in \mathbb{E}$ ,  $\operatorname{Re}(H)\operatorname{Im}(H) \neq 0$  and  $H(x, y) \in L_{p,x}(a, b)$  for almost all  $y \in (c, d)$ . Then*

$$(3.1) \quad \left\| \int_c^d H(\cdot, y) dy \right\|_{L_p(a,b)} \geq (\sqrt{2})^{p-2} \int_c^d \|H(\cdot, y)\|_{L_p(a,b)} dy,$$

if left-hand side is finite.

*Proof.* We have

$$\left| \int_c^d H(x, y) dy \right| \leq \int_c^d |H(x, y)| dy.$$

Then for  $p - 1 < 0$  we get

$$\left| \int_c^d H(x, y) dy \right|^{p-1} \geq \left( \int_c^d |H(x, y)| dy \right)^{p-1}.$$

By Proposition 2.1, we obtain

$$\begin{aligned} \left| \int_c^d H(x, y) dy \right|^p &= \left| \int_c^d H(x, y) dy \right|^{p-1} \left| \int_c^d H(x, y) dy \right| \\ &\geq \left( \int_c^d |H(x, y)| dy \right)^{p-1} \left| \int_c^d H(x, y) dy \right| \\ &\geq \left( \int_c^d |H(x, y)| dy \right)^{p-1} (\sqrt{2})^{-1} \left| \int_c^d |H(x, y)| dy \right| \\ &= (\sqrt{2})^{-1} \left( \int_c^d |H(x, y)| dy \right)^{p-1} \left| \int_c^d |H(x, y)| dy \right|. \end{aligned}$$

By integrating the last inequality, we establish

$$\begin{aligned} \int_a^b \left| \int_c^d H(x, y) dy \right|^p dx &\geq (\sqrt{2})^{-1} \int_a^b \left( \int_c^d |H(x, t)| dt \right)^{p-1} \left| \int_c^d |H(x, y)| dy \right| dx \\ &= (\sqrt{2})^{-1} \int_a^b \left| \int_c^d \left( \int_c^d |H(x, t)| dt \right)^{p-1} |H(x, y)| dy \right| dx \\ &\geq (\sqrt{2})^{-1} \left| \int_a^b \left\{ \int_c^d \left( \int_c^d |H(x, t)| dt \right)^{p-1} |H(x, y)| dy \right\} dx \right| \\ &= (\sqrt{2})^{-1} \left| \int_c^d \left\{ \int_a^b \left( \int_c^d |H(x, t)| dt \right)^{p-1} |H(x, y)| dx \right\} dy \right|. \end{aligned}$$

Let

$$R_1 = \int_a^b \left( \int_c^d |H(x, t)| dt \right)^{p-1} |H(x, y)| dx$$

and suppose that  $G(x) = \left( \int_c^d |H(x, y)| dy \right)^{p-1}$ .

Therefore, we get

$$\begin{aligned} \|G(x)\|_{L_{p'}((a,b))} &= \left( \int_a^b \left| \int_c^d |H(x, y)| dy \right|^{p'(p-1)} dx \right)^{\frac{1}{p'}} \\ &= \left( \int_a^b \left| \int_c^d |H(x, y)| dy \right|^p dx \right)^{\frac{p-1}{p}} \\ &= \left\{ \left( \int_a^b \left| \int_c^d |H(x, y)| dy \right|^p dx \right)^{\frac{1}{p}} \right\}^{p-1} \\ &= \left\| \int_c^d |H(x, y)| dy \right\|_{L_p((a,b))}^{p-1}. \end{aligned}$$

The last expression is finite (see hypotheses of theorem) then  $G(x) \in L_{p'}((a, b))$ . By applying the reverse Hölder's inequality and using Proposition 2.1, we obtain

$$\begin{aligned} R_1 &\geq \left( \int_a^b \left| \int_c^d |H(x, t)| dt \right|^{p'(p-1)} dx \right)^{\frac{1}{p'}} \left( \int_a^b |H(x, y)|^p dx \right)^{\frac{1}{p}} \\ &= \left( \int_a^b \left| \int_c^d |H(x, t)| dt \right|^p dx \right)^{\frac{1}{p'}} \left( \int_a^b |H(x, y)|^p dx \right)^{\frac{1}{p}} \\ &\geq \left( \int_a^b (\sqrt{2})^p \left| \int_c^d H(x, t) dt \right|^p dx \right)^{\frac{1}{p'}} \left( \int_a^b |H(x, y)|^p dx \right)^{\frac{1}{p}} \\ &= (\sqrt{2})^{p-1} \left( \int_a^b \left| \int_c^d H(x, t) dt \right|^p dx \right)^{\frac{1}{p'}} \left( \int_a^b |H(x, y)|^p dx \right)^{\frac{1}{p}} = R_2. \end{aligned}$$

Then we get

$$\begin{aligned} \int_c^d R_1 dy &\geq \int_c^d R_2 dy, \\ R_2 > 0 &\rightarrow \left| \int_c^d R_1 dy \right| \geq \left| \int_c^d R_2 dy \right| = \int_c^d R_2 dy. \end{aligned}$$

Thus, we conclude that

$$\begin{aligned} \int_a^b \left| \int_c^d H(x, y) dy \right|^p dx &\geq (\sqrt{2})^{-1} \left| \int_c^d R_1 dy \right| \\ &\geq (\sqrt{2})^{-1} \int_c^d R_2 dy \\ &= (\sqrt{2})^{p-2} \left( \int_a^b \left| \int_c^d H(x, t) dt \right|^p dx \right)^{\frac{1}{p'}} \left( \int_a^b |H(x, y)|^p dx \right)^{\frac{1}{p}} dy. \end{aligned}$$

Therefore, we get

$$\begin{aligned} & \left( \int_a^b \left| \int_c^d H(x, y) dy \right|^p dx \right) \left( \int_a^b \left| \int_c^d H(x, t) dt \right|^p dx \right)^{-\frac{1}{p'}} \\ & \geq (\sqrt{2})^{p-2} \int_c^d \left( \int_a^b |H(x, y)|^p dx \right)^{\frac{1}{p}} dy, \end{aligned}$$

then

$$\left( \int_a^b \left| \int_c^d H(x, y) dy \right|^p dx \right)^{1-\frac{1}{p'}} \geq (\sqrt{2})^{p-2} \int_c^d \left( \int_a^b |H(x, y)|^p dx \right)^{\frac{1}{p}} dy.$$

Finally, we conclude that

$$\left( \int_a^b \left| \int_c^d H(x, y) dy \right|^p dx \right)^{\frac{1}{p}} \geq (\sqrt{2})^{p-2} \int_c^d \left( \int_a^b |H(x, y)|^p dx \right)^{\frac{1}{p}} dy,$$

which completes the proof. □

**Theorem 3.2.** *Let  $0 < p < 1$ ,  $-\infty < a < b < +\infty$  and  $-\infty < c < d < +\infty$ . Suppose that  $H : (a, b) \times (c, d) \rightarrow \mathbb{E}$  is measurable and  $H(x, y) \in L_{p,x}(a, b)$  for almost all  $y \in (c, d)$ . Then*

$$(3.2) \quad \left\| \int_c^d H(\cdot, y) dy \right\|_{L_p(a,b)} \geq \int_c^d \|H(\cdot, y)\|_{L_p(a,b)} dy,$$

*if left-hand side is finite.*

**Theorem 3.3.** *Let  $0 < p < 1$ ,  $-\infty < a < b < +\infty$  and  $-\infty < c < d < +\infty$ . Suppose that  $H : (a, b) \times (c, d) \rightarrow i\mathbb{E}$  is measurable and  $H(x, y) \in L_{p,x}(a, b)$  for almost all  $y \in (c, d)$ . Then*

$$(3.3) \quad \left\| \int_c^d H(\cdot, y) dy \right\|_{L_p(a,b)} \geq \int_c^d \|H(\cdot, y)\|_{L_p(a,b)} dy,$$

*if left-hand side is finite.*

*Proof.* The proof of Theorem 3.2 and Theorem 3.3 is similar to Theorem 3.1. □

**Theorem 3.4.** *Let  $p < 0$ ,  $-\infty < a < b < +\infty$  and  $-\infty < c < d < +\infty$ . Suppose that  $H : (a, b) \times (c, d) \rightarrow \mathbb{C}$  is measurable with  $\text{Re}(H), \text{Im}(H) \in \mathbb{E}$ ,  $\text{Re}(H)\text{Im}(H) \neq 0$  and  $H(x, y) \in L_{p,x}(a, b)$  for almost all  $y \in (c, d)$ . Then*

$$(3.4) \quad \left\| \int_c^d H(\cdot, y) dy \right\|_{L_p(a,b)} \geq (\sqrt{2})^{p-2} \int_c^d \|H(\cdot, y)\|_{L_p(a,b)} dy,$$

*if left-hand side is finite.*

*Proof.* By using the inequality

$$\left| \int_c^d H(x, y) dy \right| \leq \int_c^d |H(x, y)| dy,$$

we get

$$\left| \int_c^d H(x, y) dy \right|^p \geq \left( \int_c^d |H(x, y)| dy \right)^p, \quad \text{for } p < 0.$$

By integrating the last inequality, we get

$$\begin{aligned} \int_a^b \left| \int_c^d H(x, y) dy \right|^p dx &\geq \int_a^b \left( \int_c^d |H(x, y)| dy \right)^p dx \\ &= \int_a^b \left[ \left( \int_c^d |H(x, t)| dt \right)^{p-1} \left( \int_c^d |H(x, y)| dy \right) \right] dx \\ &= \int_a^b \left[ \int_c^d \left( \int_c^d |H(x, t)| dt \right)^{p-1} |H(x, y)| dy \right] dx \\ &= \int_c^d \left\{ \int_a^b \left( \int_c^d |H(x, t)| dt \right)^{p-1} |H(x, y)| dx \right\} dy. \end{aligned}$$

Let

$$R_3 = \int_a^b \left( \int_c^d |H(x, t)| dt \right)^{p-1} |H(x, y)| dx.$$

By the reverse Hölder's inequality and Proposition 2.1, we obtain

$$\begin{aligned} R_3 &\geq \left( \int_a^b \left| \int_c^d |H(x, t)| dt \right|^{p'(p-1)} dx \right)^{\frac{1}{p'}} \left( \int_a^b |H(x, y)|^p dx \right)^{\frac{1}{p}} \\ &= \left( \int_a^b \left| \int_c^d |H(x, t)| dt \right|^p dx \right)^{\frac{1}{p'}} \left( \int_a^b |H(x, y)|^p dx \right)^{\frac{1}{p}} \\ &\geq \left( \int_a^b (\sqrt{2})^p \left| \int_c^d H(x, t) dt \right|^p dx \right)^{\frac{1}{p'}} \left( \int_a^b |H(x, y)|^p dx \right)^{\frac{1}{p}} \\ &= (\sqrt{2})^{p-1} \left( \int_a^b \left| \int_c^d H(x, t) dt \right|^p dx \right)^{\frac{1}{p'}} \left( \int_a^b |H(x, y)|^p dx \right)^{\frac{1}{p}} = R_4. \end{aligned}$$

That is, we get

$$\int_c^d R_3 dy \geq \int_c^d R_4 dy.$$

Therefore, we obtain

$$\int_a^b \left| \int_c^d H(x, y) dy \right|^p dx \geq \int_c^d R_3 dy \geq \int_c^d R_4 dy$$



and

$$\begin{aligned} \int_c^d R_4 dy &= (\sqrt{2})^{p-1} \int_c^d \left( \int_a^b \left| \int_c^d H(x,t) dt \right|^p dx \right)^{\frac{1}{p'}} \left( \int_a^b |H(x,y)|^p dx \right)^{\frac{1}{p}} dy \\ &= (\sqrt{2})^{p-1} \left( \int_a^b \left| \int_c^d H(x,t) dt \right|^p dx \right)^{\frac{1}{p'}} \int_c^d \left( \int_a^b |H(x,y)|^p dx \right)^{\frac{1}{p}} dy. \end{aligned}$$

It follows that

$$\begin{aligned} &\left( \int_a^b \left| \int_c^d H(x,y) dy \right|^p dx \right) \left( \int_a^b \left| \int_c^d H(x,t) dt \right|^p dx \right)^{-\frac{1}{p'}} \\ &\geq (\sqrt{2})^{p-1} \int_c^d \left( \int_a^b |H(x,y)|^p dx \right)^{\frac{1}{p}} dy. \end{aligned}$$

Consequently, we get

$$\begin{aligned} \left( \int_a^b \left| \int_c^d H(x,y) dy \right|^p dx \right)^{\frac{1}{p}} &\geq (\sqrt{2})^{p-1} \int_c^d \left( \int_a^b |H(x,y)|^p dx \right)^{\frac{1}{p}} \\ &\geq (\sqrt{2})^{p-2} \int_c^d \left( \int_a^b |H(x,y)|^p dx \right)^{\frac{1}{p}}. \end{aligned}$$

This completes the proof. □

**Theorem 3.5.** *Let  $p < 0$ ,  $-\infty < a < b < +\infty$  and  $-\infty < c < d < +\infty$ . Suppose that  $H : (a, b) \times (c, d) \rightarrow \mathbb{E}$  is measurable and  $H(x, y) \in L_{p,x}(a, b)$  for almost all  $y \in (c, d)$ . Then*

$$(3.5) \quad \left\| \int_c^d H(\cdot, y) dy \right\|_{L_p(a,b)} \geq \int_c^d \|H(\cdot, y)\|_{L_p(a,b)} dy,$$

*if left-hand side is finite.*

**Theorem 3.6.** *Let  $p < 0$ ,  $-\infty < a < b < +\infty$  and  $-\infty < c < d < +\infty$ . Suppose that  $H : (a, b) \times (c, d) \rightarrow i\mathbb{E}$  is measurable and  $H(x, y) \in L_{p,x}(a, b)$  for almost all  $y \in (c, d)$ . Then*

$$(3.6) \quad \left\| \int_c^d H(\cdot, y) dy \right\|_{L_p(a,b)} \geq \int_c^d \|H(\cdot, y)\|_{L_p(a,b)} dy,$$

*if left-hand side is finite.*

*Proof.* The proof of Theorem 3.5 and Theorem 3.6 is similar to Theorem 3.4. □

**Acknowledgements.** The author is grateful to Pr. Abdelkader Senouci (Algeria) and Ass. Pr. Hüseyin Bubak (Turkey) for helpful comments. The author wish to thank the support of DG-RSDT.

## REFERENCES

- [1] V. I. Burenkov, *Functional Spaces. Basic Integral Inequalities Connected with  $L_p$ -Spaces*, Peoples' Friendship University of Russia, Moscow, 1989, 5–12.
- [2] S. Marinković, P. Rajković and M. Stanković, *The inequalities for some types  $q$ -integrals*, *Comput. Math. Appl.* **56** (2008), 2490–2498.
- [3] B. G. Pachpatte, *Mathematical Inequalities*, North Holland Mathematical Library **67**, Oxford, 2005.
- [4] B. Yang, *On a new Hardy-type integral inequality*, *International Mathematical Forum* **67**(2) (2007), 3317–3322.

<sup>1</sup>FACULTY OF MATERIAL SCIENCES,  
LABORATORY OF INFORMATICS AND MATHEMATICS,  
UNIVERSITY OF TIARET, ALGERIA

<sup>2</sup>UNIV-INSCRIPTION: DJILLALI LIABES UNIVERSITY,  
SIDI BEL ABBÈS, 22000, ALGERIA  
*Email address:* bouharket.benaïssa@univ-tiaret.dz