

SOME REMARKS ON THE RANDIĆ ENERGY OF GRAPHS

Š. B. BOZKURT ALTINDAĞ¹, I. MILOVANOVIĆ², AND E. MILOVANOVIĆ²

ABSTRACT. Let G be a graph of order n . The Randić energy of G is defined as $RE(G) = \sum_{i=1}^n |\rho_i|$, where $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$ are the Randić eigenvalues of G . In this study, we present improved bounds for $RE(G)$ as well as a relationship between (ordinary) graph energy and $RE(G)$.

1. INTRODUCTION

Let $G = (V, E)$, $V = \{v_1, v_2, \dots, v_n\}$, be a simple connected graph of order n and size m , with vertex degree sequence $\Delta = d_1 \geq d_2 \geq \dots \geq d_n = \delta$, $d_i = d(v_i)$. Denote by $D = \text{diag}(d_1, d_2, \dots, d_n)$ the diagonal matrix of its vertex degrees. If vertices v_i and v_j are adjacent in G , it will be denoted as $i \sim j$.

Let $A = (a_{ij})$, be the $(0, 1)$ adjacency matrix of G . The eigenvalues of matrix A , $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$, are the (ordinary) eigenvalues of G [4]. Some well known properties of these eigenvalues are [4]:

$$\sum_{i=1}^n \lambda_i = \text{tr}(A) = 0, \quad \sum_{i=1}^n \lambda_i^2 = \text{tr}(A^2) = \sum_{i=1}^n d_i = 2m, \quad \prod_{i=1}^n \lambda_i = \det A.$$

Denote with $|\lambda_1^*| \geq |\lambda_2^*| \geq \dots \geq |\lambda_n^*|$ the non-increasing arrangement of the absolute values of eigenvalues of G . The notion of (ordinary) graph energy was introduced in [12]. It is defined to be

$$E(G) = \sum_{i=1}^n |\lambda_i| = \sum_{i=1}^n |\lambda_i^*|.$$

Key words and phrases. Graph spectrum, Randić spectrum, graph energy, Randić energy.
 2020 *Mathematics Subject Classification.* Primary: 05C50. Secondary: 05C90.
 DOI

Received: March 15, 2022.

Accepted: June 22, 2022.

The Randić matrix of G [2] is defined as

$$R = R(G) = D^{-1/2}AD^{-1/2}.$$

The eigenvalues of matrix R , $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$, form the Randić spectrum of G . Some properties of Randić eigenvalues are (see, e.g., [2]):

$$\sum_{i=1}^n \rho_i = \text{tr}(R) = 0, \quad \sum_{i=1}^n \rho_i^2 = \text{tr}(R^2) = 2R_{-1}(G),$$

where $R_{-1}(G)$ is a vertex-degree based graph invariant introduced in [3] defined as

$$R_{-1}(G) = \sum_{i \sim j} \frac{1}{d_i d_j}.$$

It is known as general Randić index R_{-1} , as well as modified second Zagreb index [24].

In [14] it was proven that the following identity is valid

$$(1.1) \quad \det R = \frac{\det A}{\prod_{i=1}^n d_i}.$$

The other two vertex-degree based topological indices that are of interest for the present paper are the first Zagreb index [17]

$$M_1(G) = \sum_{i=1}^n d_i^2 = \sum_{i \sim j} (d_i + d_j),$$

and the inverse degree index [9] defined as

$$ID(G) = \sum_{i=1}^n \frac{1}{d_i} = \sum_{i \sim j} \left(\frac{1}{d_i^2} + \frac{1}{d_j^2} \right).$$

Denote with $|\rho_1^*| \geq |\rho_2^*| \geq \dots \geq |\rho_n^*|$ the non-increasing arrangement of the absolute values of Randić eigenvalues of G . The Randić energy of G is defined as [2]

$$RE(G) = \sum_{i=1}^n |\rho_i| = \sum_{i=1}^n |\rho_i^*|.$$

More on its mathematical properties can be found in [1–3, 5, 7, 14, 20, 22].

In this paper, we obtain improved bounds for $RE(G)$ as well as a relationship between $E(G)$ and $RE(G)$.

2. PRELIMINARIES

In this section we recall some results from spectral graph theory and analytical inequalities that are of interest for the present paper.

Lemma 2.1 ([20]). *The Randić spectral radius is $\rho_1 = 1$.*

Remark 2.1. In [14] it was observed that when $G \cong \overline{K}_n$ then $\rho_1 = 0$. Therefore, if G has at least one edge, then $\rho_1 = 1$.

Let $G_1 \vee G_2$ denote the complete product of two graphs G_1 and G_2 . This graph is obtained from $G_1 \cup G_2$ by joining every vertex of G_1 with every vertex of G_2 .

Lemma 2.2 ([8]). *Let G be a connected graph of order n with maximum vertex degree $\Delta = n - 1$. Then $|\rho_2| = |\rho_3| = \dots = |\rho_n|$ if and only if $G \cong K_n$, or $G \cong K_1 \vee r K_2$, with $n = 2r + 1$ ($r \geq 2$).*

Lemma 2.3 ([20]). *Let G be a connected graph of order n . Then*

$$(2.1) \quad RE(G) \leq 1 + \sqrt{(n-1)(2R_{-1}(G) - 1)}.$$

Remark 2.2. The inequality (2.1) was also proved in [19, 21], as well as in [5] as a special case of one more general result. In [8] it was proved that when $\Delta = n - 1$, equality in (2.1) holds if and only if $G \cong K_n$, or $G \cong K_1 \vee r K_2$, with $n = 2r + 1$ ($r \geq 2$).

Lemma 2.4 ([1]). *Let G be a connected bipartite graph of order $n \geq 2$. Then*

$$(2.2) \quad RE(G) \leq 2 + \sqrt{(n-2)(2R_{-1}(G) - 2)}.$$

Remark 2.3. The inequality (2.2) was also proved in [21]. In [8] it was proven that equality in (2.2) holds if and only if $G \cong K_{p,q}$, $p + q = n$, for odd n .

Lemma 2.5 ([11]). *Let G be a connected bipartite graph of order $n \geq 3$ with Randić eigenvalues $\rho_1 = 1 \geq \rho_2 \geq \dots \geq \rho_{n-1} \geq \rho_n = -1$ and let $\rho = \max_{2 \leq i \leq n-1} \{|\rho_i|\}$. Then, for any real k , $\rho \geq k \geq \sqrt{\frac{2R_{-1}(G)-2}{n-2}}$, holds*

$$(2.3) \quad RE(G) \leq 2 + k + \sqrt{(n-3)(2R_{-1}(G) - 2 - k^2)}.$$

Equality holds if G is a complete bipartite graph, in which case $k = 0$.

Remark 2.4. In [18, Theorem 3.4] it was claimed that when

$$(2.4) \quad \frac{1}{\Delta} \geq \sqrt{\frac{2R_{-1}(G) - 1}{n - 1}},$$

then

$$(2.5) \quad RE(G) \leq 1 + \frac{1}{\Delta} + \sqrt{(n-2) \left(2R_{-1}(G) - 1 - \frac{1}{\Delta^2} \right)},$$

which would mean that (2.5) is stronger than (2.1). However, if (2.4) is true, then $\Delta \geq n - 1$, which is not possible. Therefore, the inequality (2.5) is not correct.

Lemma 2.6 ([6, 18]). *Let G be a connected graph of order n . Then*

$$(2.6) \quad RE(G) \geq 1 + (n-1) \left(\frac{|\det A|}{\prod_{i=1}^n d_i} \right)^{\frac{1}{n-1}}.$$

The following analytical inequality would be used in proofs of theorems in the present paper.

Lemma 2.7 ([23]). *Let $p = (p_i)$, $i = 1, 2, \dots, n$, be a sequence of positive real numbers and $a = (a_i)$ and $b = (b_i)$, $i = 1, 2, \dots, n$, two similarly ordered sequences of non-negative real numbers. Then*

$$(2.7) \quad \sum_{i=1}^n p_i \sum_{i=1}^n p_i a_i b_i \geq \sum_{i=1}^n p_i a_i \sum_{i=1}^n p_i b_i.$$

When $a = (a_i)$ and $b = (b_i)$, $i = 1, 2, \dots, n$, are of different monotonicity, then opposite inequality is valid. Equality holds if and only if $a_1 = \dots = a_n$ or $b_1 = \dots = b_n$.

3. MAIN RESULTS

In the next theorem we establish a lower bound on $RE(G)$.

Theorem 3.1. *Let G be a connected graph of order n . Then, for any real k , $|\rho_2^*| \geq k \geq \sqrt{\frac{2R_{-1}(G)-1}{n-1}}$, holds*

$$(3.1) \quad RE(G) \geq 1 + k + (n-2) \left(\frac{|\det A|}{k \prod_{i=1}^n d_i} \right)^{\frac{1}{n-2}}.$$

Equality holds if and only if G is a graph with the property $\rho_1 = |\rho_1^*| = 1$, $|\rho_2^*| = k$, and $|\rho_i^*| = \sqrt{\frac{2R_{-1}(G)-1-k^2}{n-2}}$, for $i = 3, 4, \dots, n$.

Proof. Using arithmetic-geometric mean inequality (see, e.g., [23]), Lemma 2.1 and (1.1) we obtain

$$\begin{aligned} RE(G) &= \sum_{i=1}^n |\rho_i^*| = 1 + |\rho_2^*| + \sum_{i=3}^n |\rho_i^*| \\ &\geq 1 + |\rho_2^*| + (n-2) \left(\frac{|\det R|}{|\rho_2^*|} \right)^{\frac{1}{n-2}} \\ &= 1 + |\rho_2^*| + (n-2) \left(\frac{|\det A|}{|\rho_2^*| \prod_{i=1}^n d_i} \right)^{\frac{1}{n-2}}. \end{aligned} \quad (3.2)$$

Let us consider the following function defined by

$$f(k) = x + (n-2) \left(\frac{|\det A|}{x \prod_{i=1}^n d_i} \right)^{\frac{1}{n-2}}.$$

Observe that f is increasing for $x \geq \left(\frac{|\det A|}{\prod_{i=1}^n d_i} \right)^{\frac{1}{n-1}}$. Considering Lemmas 2.1 and 2.3 together with (1.1), for any real k , $|\rho_2^*| \geq k \geq \sqrt{\frac{2R_{-1}(G)-1}{n-1}}$, we have that

$$|\rho_2^*| \geq k \geq \sqrt{\frac{2R_{-1}(G)-1}{n-1}} \geq \frac{RE(G)-1}{n-1} = \frac{\sum_{i=2}^n |\rho_i^*|}{n-1} \geq \left(\prod_{i=2}^n |\rho_i^*| \right)^{\frac{1}{n-1}} = \left(\frac{|\det A|}{\prod_{i=1}^n d_i} \right)^{\frac{1}{n-1}}.$$

Then, we deduce that $f(|\rho_2^*|) \geq f(k)$. Combining this with (3.2), the inequality (3.1) is obtained. The equality in (3.1) holds if and only if

$$|\rho_2^*| = k \quad \text{and} \quad |\rho_3^*| = \cdots = |\rho_n^*|.$$

Since $\sum_{i=2}^n |\rho_i^*|^2 = 2R_{-1}(G) - 1$, the above conditions imply that $|\rho_3^*| = \cdots = |\rho_n^*| = \sqrt{\frac{2R_{-1}(G)-1-k^2}{n-2}}$. This completes the proof. \square

Corollary 3.1. *Let G be a connected graph of order n . Then*

$$(3.3) \quad RE(G) \geq 1 + \sqrt{\frac{2R_{-1}(G) - 1}{n - 1}} + (n - 2) \left(\frac{|\det A|}{\sqrt{\frac{2R_{-1}(G)-1}{n-1}} \prod_{i=1}^n d_i} \right)^{\frac{1}{n-2}}.$$

If the maximum vertex degree Δ is equal to $n - 1$, the equality in (3.3) holds if and only if $G \cong K_n$, or $G \cong K_1 \vee r K_2$, with $n = 2r + 1$ ($r \geq 2$).

Proof. The inequality (3.3) is obtained from (3.1) for $k = \sqrt{\frac{2R_{-1}(G)-1}{n-1}}$. Now, assume that equality in (3.3) holds. Then

$$|\rho_2^*| = \sqrt{\frac{2R_{-1}(G) - 1}{n - 1}} \quad \text{and} \quad |\rho_3^*| = \cdots = |\rho_n^*|.$$

Since $\sum_{i=2}^n |\rho_i^*| = 2R_{-1}(G) - 1$, we get

$$|\rho_3^*| = \cdots = |\rho_n^*| = \sqrt{\frac{2R_{-1}(G) - 1}{n - 1}}.$$

The above results state that $|\rho_2^*| = |\rho_3^*| = \cdots = |\rho_n^*|$, that is $|\rho_2| = |\rho_3| = \cdots = |\rho_n|$. Then, by Lemma 2.2 if $\Delta = n - 1$, the equality in (3.3) holds if and only if $G \cong K_n$, or $G \cong K_1 \vee r K_2$, with $n = 2r + 1$ ($r \geq 2$). \square

Remark 3.1. The lower bounds (3.1) and (3.3) are stronger than the lower bound (2.6). Moreover, by Theorem 3.1, it is possible to derive stronger lower bound than (3.3) using any real k such that $|\rho_2^*| \geq k \geq \sqrt{\frac{2R_{-1}(G)-1}{n-1}}$.

In the next theorem we establish a relationship between Randić energy and general Randić index R_{-1} .

Theorem 3.2. *Let G be a connected graph of order $n \geq 3$. Then, for any real k , such that $|\rho_2^*| \geq k \geq \sqrt{\frac{2R_{-1}(G)-1}{n-1}}$, we have*

$$(3.4) \quad RE(G) \leq 1 + k + \sqrt{(n - 2)(2R_{-1}(G) - 1 - k^2)}.$$

Equality holds if and only if G is a graph with the property $\rho_1 = |\rho_1^| = 1$, $|\rho_2^*| = k$ and $|\rho_i^*| = \sqrt{\frac{2R_{-1}(G)-1-k^2}{n-2}}$, for $i = 3, 4, \dots, n$.*

Proof. By the Cauchy-Schwarz inequality (see, e.g., [23]), we have that

$$\sum_{i=3}^n |\rho_i^*| \leq \left(\sum_{i=3}^n 1 \right)^{1/2} \left(\sum_{i=3}^n |\rho_i^*|^2 \right)^{1/2},$$

that is

$$RE(G) \leq |\rho_1^*| + |\rho_2^*| + \sqrt{(n-2)(2R_{-1}(G) - |\rho_1^*|^2 - |\rho_2^*|^2)}.$$

By Lemma 2.1, we have that $\rho_1 = |\rho_1^*| = 1$. Considering this fact with the above inequality, we get

$$(3.5) \quad RE(G) \leq 1 + |\rho_2^*| + \sqrt{(n-2)(2R_{-1}(G) - 1 - |\rho_2^*|^2)}.$$

Now, observe the function

$$f(x) = x + \sqrt{(n-2)(2R_{-1}(G) - 1 - x^2)}, \quad x \geq 0.$$

This function is monotone decreasing for $x \geq \sqrt{\frac{2R_{-1}(G)-1}{n-1}}$. Therefore for any $k \geq 0$ with the property $|\rho_2^*| \geq k \geq \sqrt{\frac{2R_{-1}(G)-1}{n-1}}$, holds that $f(|\rho_2^*|) \leq f(k)$. From this inequality and (3.5) we obtain (3.4).

The equality case for (3.4) can be proved similarly as in case of Theorem 3.1. \square

Remark 3.2. When $k = \sqrt{\frac{2R_{-1}(G)-1}{n-1}}$, from (3.4) the inequality (2.1) is obtained, which means that inequality (3.4) is stronger than (2.1).

Remark 3.3. Recall that the Randić spectrum of a bipartite graph is symmetric with respect to the origin, that is, $\rho_i = -\rho_{n-i+1}$, for $i = 1, 2, \dots, n$ [10]. In this case, $|\rho_1^*| = \rho_1 = 1 = |\rho_n| = |\rho_2^*|$. On the other hand, $\rho = |\rho_3^*| = |\rho_4^*|$.

Having in mind the above remark, by a similar procedure as in Theorem 3.2, the following result can be proven.

Theorem 3.3. *Let G be a connected bipartite graph of order $n \geq 5$. Then, for any real k such that $|\rho_3^*| \geq k \geq \sqrt{\frac{2R_{-1}(G)-2}{n-2}}$, we have*

$$(3.6) \quad RE(G) \leq 2 + 2k + \sqrt{(n-4)(2R_{-1}(G) - 2 - 2k^2)}.$$

Equality holds if and only if G is a graph with the property $\rho_1 = |\rho_1^| = |\rho_2^*| = 1$, $|\rho_3^*| = |\rho_4^*| = k$ and $|\rho_i^*| = \sqrt{\frac{2R_{-1}(G)-2-2k^2}{n-4}}$, for $i = 5, \dots, n$.*

Remark 3.4. When $k = \sqrt{\frac{2R_{-1}(G)-2}{n-2}}$, from (3.6) the inequality (2.2) is obtained. Furthermore, the inequality (3.6) is stronger than (2.2) and (2.3).

Theorem 3.4. *Let G be a connected graph of order $n \geq 2$. Then*

$$(3.7) \quad RE(G) \leq 1 + \sqrt{(n-1) \left(2R_{-1}(G) - 1 - \frac{1}{2}(|\rho_2^*| - |\rho_n^*|)^2 \right)}.$$

Equality holds if and only if G is a graph with the property $\rho_1 = |\rho_1^| = 1$ and $|\rho_3^*| = \dots = |\rho_{n-1}^*| = \frac{|\rho_2^*| + |\rho_n^*|}{2}$.*

Proof. Based on the Lagrange's identity (see e.g. [23]), we have that

$$\begin{aligned}
 (n-1) \sum_{i=2}^n |\rho_i^*|^2 - \left(\sum_{i=2}^n |\rho_i^*| \right)^2 &= \sum_{2 \leq i < j \leq n} (|\rho_i^*| - |\rho_j^*|)^2 \\
 &\geq (|\rho_2^*| - |\rho_n^*|)^2 + \sum_{i=3}^{n-1} \left((|\rho_i^*| - |\rho_n^*|)^2 + (|\rho_2^*| - |\rho_i^*|)^2 \right) \\
 (3.8) \quad &\geq (|\rho_2^*| - |\rho_n^*|)^2 + \frac{1}{2} \sum_{i=3}^{n-1} (|\rho_2^*| - |\rho_n^*|)^2 \\
 &= \frac{n-1}{2} (|\rho_2^*| - |\rho_n^*|)^2.
 \end{aligned}$$

Since

$$(n-1) \sum_{i=2}^n |\rho_i^*|^2 - \left(\sum_{i=2}^n |\rho_i^*| \right)^2 = (n-1)(2R_{-1}(G) - 1) - (RE(G) - 1)^2,$$

from (3.8) the inequality (3.7) is obtained.

Equality in (3.8) holds if and only if $|\rho_3^*| = \dots = |\rho_{n-1}^*|$ and $|\rho_i^*| - |\rho_n^*| = |\rho_2^*| - |\rho_i^*|$, for $i = 3, \dots, n-1$, which implies that equality in (3.7) holds if and only if $\rho_1 = |\rho_1^*| = 1$ and $|\rho_3^*| = \dots = |\rho_{n-1}^*| = \frac{|\rho_2^*| + |\rho_n^*|}{2}$. \square

Remark 3.5. Let us note that the inequality (3.7) is stronger than (2.1).

The proof of the next theorem is analogous to that of Theorem 3.4, hence omitted.

Theorem 3.5. *Let G be a connected bipartite graph of order $n \geq 4$. Then*

$$(3.9) \quad RE(G) \leq 2 + \sqrt{(n-2) \left(2R_{-1}(G) - 2 - \frac{1}{2}(|\rho_3^*| - |\rho_n^*|)^2 \right)}.$$

Equality holds if and only if G is a graph with the property $\rho_1 = |\rho_1^| = |\rho_2^*| = 1$ and $|\rho_4^*| = \dots = |\rho_{n-1}^*| = \frac{|\rho_3^*| + |\rho_n^*|}{2}$.*

Remark 3.6. Notice that the inequality (3.9) is stronger than (2.2).

We now give a relationship between $E(G)$ and $RE(G)$.

Theorem 3.6. *Let G be a graph of order $n \geq 2$ and size m , without isolated vertices. Then we have*

$$(3.10) \quad E(G)RE(G) \leq 2n\sqrt{mR_{-1}(G)}.$$

Equality holds if and only if $G \cong \frac{n}{2}K_2$, for even n .

Proof. For $p_i = 1$, $a_i = |\lambda_i^*|$, $b_i = |\rho_i^*|$, $i = 1, 2, \dots, n$, the inequality (2.7) becomes

$$\sum_{i=1}^n 1 \sum_{i=1}^n |\lambda_i^*| |\rho_i^*| \geq \sum_{i=1}^n |\lambda_i^*| \sum_{i=1}^n |\rho_i^*|,$$

that is

$$(3.11) \quad E(G)RE(G) \leq n \sum_{i=1}^n |\lambda_i^*| |\rho_i^*|.$$

On the other hand, having in mind Cauchy–Schwarz inequality, we have that

$$n \sum_{i=1}^n |\lambda_i^*| |\rho_i^*| \leq n \left(\sum_{i=1}^n |\lambda_i^*|^2 \right)^{1/2} \left(\sum_{i=1}^n |\rho_i^*|^2 \right)^{1/2},$$

that is

$$(3.12) \quad n \sum_{i=1}^n |\lambda_i^*| |\rho_i^*| \leq 2n\sqrt{mR_{-1}(G)}.$$

Now, from (3.11) and (3.12) we arrive at (3.10).

Equality in (3.11) holds if and only if $|\lambda_1^*| = \dots = |\lambda_n^*|$, or $|\rho_1^*| = \dots = |\rho_n^*|$. Equality in (3.12) holds if and only if $|\lambda_i^*| = C|\rho_i^*|$, $C = \text{Const}$, for $i = 1, 2, \dots, n$. Thus, equalities in both (3.11) and (3.12) hold if and only if $|\lambda_1^*| = \dots = |\lambda_n^*|$ and $|\rho_1^*| = \dots = |\rho_n^*|$. Since G has no isolated vertices, equality in (3.10) holds if and only if $G \cong \frac{n}{2}K_2$, for even n . \square

Recall that the Sombor energy of a graph G , denoted by $E_{SO}(G)$, is introduced as the sum of the absolute values of the eigenvalues of its Sombor matrix [13]. The following relationship between graph energy and Sombor energy can be found in [15].

Theorem 3.7 ([15]). *If G is a bipartite graph whose all cycles (if any) have size not divisible by 4, then*

$$E_{SO}(G) \leq \sqrt{2}\Delta E(G).$$

From Theorem 3.6 and Theorem 3.7, we directly have the following.

Corollary 3.2. *If G is a bipartite graph whose all cycles (if any) have size not divisible by 4, then*

$$E_{SO}(G)RE(G) \leq 2\Delta n\sqrt{2mR_{-1}(G)}.$$

Corollary 3.3. *Let G be a graph of order $n \geq 2$ and size m , without isolated vertices. Then we have*

$$E(G) \leq n\sqrt{mR_{-1}(G)}.$$

Proof. In [3] it was proven that $RE(G) \geq 2$. Considering this with inequality (3.10) we obtain the required result. \square

Corollary 3.4. *Let G be a graph of order $n \geq 2$ and size m , without isolated vertices. Then we have*

$$(3.13) \quad E(G) \leq n\sqrt{\frac{M_2(G)}{m}},$$

where $M_2(G) = \sum_{i \sim j} d_i d_j$ is the second Zagreb index [16].

Proof. In [3] it was also proven that

$$RE(G) \geq 2R_{-1}(G).$$

From the above and inequality (3.10) we obtain

$$(3.14) \quad E(G) \leq n \sqrt{\frac{m}{R_{-1}(G)}}.$$

On the other hand, by the inequality between arithmetic and harmonic means (see, e.g., [23]), we have that

$$M_2(G)R_{-1}(G) \geq m^2.$$

Combining the above and inequality (3.14) we arrive at (3.13). \square

Acknowledgement. This research was partly supported by the Serbian Ministry of Education, Science and Technological Development, grant No. 451-03-68/2022-14/200102.

REFERENCES

- [1] Ş. B. Bozkurt and D. Bozkurt, *Sharp upper bounds for energy and Randić energy*, MATCH Commun. Math. Comput. Chem. **70**(2) (2013), 669–680.
- [2] Ş. B. Bozkurt, A. D. Gungor, I. Gutman and A. S. Cevik, *Randić matrix and Randić energy*, MATCH Commun. Math. Comput. Chem. **64** (1) (2010), 239–250.
- [3] M. Cavers, S. Fallat and S. Kirkland, *On the normalized Laplacian energy and general Randić index R_{-1} of graphs*, Linear Algebra Appl. **433** (2010), 172–190. <https://doi.org/10.1016/j.laa.2010.02.002>
- [4] D. Cvetković, M. Doob and H. Sachs, *Spectra of Graph-Theory and Application*, Academic Press, New York, 1980.
- [5] K. C. Das, I. Gutman, I. Milovanović, E. Milovanović and B. Furtula, *Degree-based energies of graphs*, Linear Algebra Appl. **554** (2018), 185–204. <https://doi.org/10.1016/j.laa.2018.05.027>
- [6] K. C. Das and S. Sorgun, *On Randić energy of graphs*, MATCH Commun. Math. Comput. Chem. **72**(1) (2014), 227–238.
- [7] K. C. Das, S. Sorgun and K. Xu, *On Randić energy of graphs*, in: I. Gutman and X. Li, (Eds.), *Energies of Graphs - Theory and Applications* University of Kragujevac, Kragujevac, 2016, 111–122.
- [8] K. C. Das and S. Sun, *Extremal graphs for Randić energy*, MATCH Commun. Math. Comput. Chem. **77**(1) (2017), 77–84.
- [9] S. Fajtlowicz, *On conjectures of graffiti-II*, Congr. Numer. **60** (1987) 187–197.
- [10] B. Furtula and I. Gutman, *Comparing energy and Randić energy*, Maced. J. Chem. Chem. Eng. **32** (2013), 117–123.
- [11] E. Glogić, E. Zogić and N. Glišović, *Remarks on the upper bound for the Randić energy of bipartite graphs*, Discrete Appl. Math. **221** (2017), 67–70. <https://doi.org/10.1016/j.dam.2016.12.005>
- [12] I. Gutman, *The energy of a graph*, Ber. Math.-Statist. Sect. Forschungszentrum Graz **103** (1978), 1–22.
- [13] I. Gutman, *Spectrum and energy of the Sombor matrix*, Vojnotehnički preglod **69** (2021), 551–561.
- [14] I. Gutman, B. Furtula and Ş. B. Bozkurt, *On Randić energy*, Linear Algebra App. **442** (2014), 50–57. <https://doi.org/10.1016/j.laa.2013.06.010>

- [15] I. Gutman, I. Redžepović and J. Rada, *Relating energy and Sombor energy*, Contrib. Math. **4** (2021), 41–44. <https://doi.org/10.47443/cm.2021.0054>
- [16] I. Gutman, B. Ruščić, N. Trinajstić and C. F. Wilcox, *Graph theory and molecular orbitals. XII Acyclic polyenes*, J. Chem. Phys. **62** (1975), 3399–3405.
- [17] I. Gutman and N. Trinajstić, *Graph theory and molecular orbitals. Total π -electron energy of alternant hydrocarbons*, Chem. Phys. Lett. **17** (1972), 535–538.
- [18] J. He, Y. Liu and J. Tian, *Note on the Randić energy of graphs*, Kragujevac J. Math. **42** (2018), 209–215.
- [19] J. Li, J. M. Guo and W. C. Shiu, *A note on Randić energy*, MATCH Commun. Math. Comput. Chem. **74**(2) (2015), 389–398.
- [20] B. Liu, Y. Huang and J. Feng, *A note on the Randić spectral radius*, MATCH Commun. Math. Comput. Chem. **68**(3) (2012), 913–916.
- [21] A. D. Maden, *New bounds on the incidence energy, Randić energy and Randić Estrada index*, MATCH Commun. Math. Comput. Chem. **74**(2) (2015), 367–387.
- [22] E. I. Milovanović, M. R. Popović, R. M. Stanković and I. Ž. Milovanović, *Remark on ordinary and Randić energy of graphs*, J. Math. Inequal. **10** (2016), 687–692. <https://dx.doi.org/10.7153/jmi-10-55>
- [23] D. S. Mitrinović and P. M. Vasić, *Analytic Inequalities*, Springer Verlag, Berlin, Heidelberg, New York, 1970.
- [24] S. Nikolić, G. Kovačević, A. Milićević and N. Trinajstić, *Modified Zagreb indices*, Croat. Chem. Acta **76** (2003), 113–124.

¹YENIKENT KARDELEN KONUTLARI,
SELÇUKLU, 42070 KONYA, TURKEY
Email address: srf_burcu_bozkurt@hotmail.com

²FACULTY OF ELECTRONIC ENGINEERING,
UNIVERSITY OF NIŠ,
NIŠ, SERBIA
Email address: igor@elfak.ni.ac.rs
Email address: ema@elfak.ni.ac.rs