

NEW UPPER AND LOWER BOUNDS FOR SOME DEGREE-BASED GRAPH INVARIANTS

A. GHALAVAND¹, A. ASHRAFI¹, AND I. GUTMAN²

ABSTRACT. For a simple graph G with vertex set $V(G)$ and edge set $E(G)$, let $\deg(u)$ be the degree of the vertex $u \in V(G)$. The forgotten index of G and its coindex are defined as $F(G) = \sum_{v \in V(G)} \deg^3(v)$ and $\bar{F}(G) = \sum_{uv \notin E(G)} [\deg^2(u) + \deg^2(v)]$. New bounds for the first Zagreb index $M_1(G) = \sum_{v \in V(G)} \deg(v)^2$, forgotten index, and its coindex are obtained.

1. INTRODUCTION

Throughout this paper, all graphs considered are assumed to be simple, i.e., without directed, weighted, or multiple edges, without self-loops and with a finite number of vertices. Let G be such a graph with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$ and edge set $E(G)$. A graph with n vertices and m edges will be referred to as an (n, m) -graph.

By $\deg(v)$ or $\deg_G(v)$ is denoted the degree of the vertex $v \in V(G)$. Let $D(G) = \{\deg(v_1), \deg(v_2), \dots, \deg(v_n)\}$. If $D(G) = \{r\}$, then G is said to be r -regular. If $D(G) = \{r, s\}$, then we say that G is (r, s) -biregular. This includes the case of regular graphs if $r = s$. Analogously, if $D(G) = \{r, s, t\}$, then the graph G will be said to be (r, s, t) -triregular. Let, in addition, $\Delta = \max_{v \in V(G)} \deg(v)$ and $\delta = \min_{v \in V(G)} \deg(v)$.

The *first Zagreb index* $M_1(G)$ is defined as [13]

$$M_1 = M_1(G) = \sum_{v \in V(G)} \deg^2(v) = \sum_{uv \in E(G)} [\deg(u) + \deg(v)].$$

It is the oldest and most studied degree-based graph invariant; details of its mathematical theory and chemical applications can be found in the surveys [5, 11, 17].

Key words and phrases. Degree (of vertex), coindex, forgotten index, F -index, Zagreb index.
2010 *Mathematics Subject Classification.* Primary: 05C07. Secondary: 05C90.

Received: March 6, 2018.

Accepted: March 22, 2018.

In the paper [13], M_1 was used for designing approximate expressions for total π -electron energy. In the same paper, also the sum of cubes of vertex degrees (F) was used for the same purpose. However, whereas M_1 eventually gained much popularity [5, 11, 17], no attention was paid to F . Only more than forty years later, the invariant F attracted some interest, thanks to the discovery of its applicability in physical chemistry [4]. For this reason it was named *forgotten index* and is defined as [4]:

$$F = F(G) = \sum_{v \in V(G)} \deg(v)^3 = \sum_{uv \in E(G)} [\deg(u)^2 + \deg(v)^2].$$

In the last few years, numerous mathematical studies of the forgotten index have been published, see [1–3, 6, 7, 10, 12, 16].

Some of pharmacological applications of the F -index were also attempted [15].

Both M_1 and F are special cases of the so-called *first general Zagreb index*, defined as

$$M_1^\alpha = M_1^\alpha(G) = \sum_{u \in V(G)} \deg(u)^\alpha = \sum_{uv \in E(G)} [\deg(u)^{\alpha-1} + \deg(v)^{\alpha-1}],$$

where α is an arbitrary real number [15, 18].

The coindex of M_1^α is defined as [18]

$$\overline{M}_1^\alpha(G) = \sum_{\substack{uv \notin E(G) \\ u \neq v}} [\deg(u)^{\alpha-1} + \deg(v)^{\alpha-1}].$$

The special case of this expressions for $\alpha = 3$ is the coindex of the forgotten index [8, 14]

$$\overline{F}(G) = \sum_{\substack{uv \notin E(G) \\ u \neq v}} [\deg(u)^2 + \deg(v)^2].$$

2. MAIN RESULTS

We first state results that improve those reported in [12]. Denote by \overline{G} the complement of the graph G .

Theorem 2.1. *Let G be an (n, m) -graph. Then*

$$F(G) + F(\overline{G}) = n^4 + M_1(G)(3n - 3) - 2m(3n^2 - 6n + 3) - n(3n^2 - 3n + 1)$$

and

$$\begin{aligned} F(G) \times F(\overline{G}) = & n^4 F(G) + (3n - 3)F(G) M_1(G) - 2m(3n^2 - 6n + 3)F(G) \\ & - n(3n^2 - 3n + 1)F(G) - F(G)^2. \end{aligned}$$

Proof. By definition of a graph complement, we have

$$\begin{aligned} F(\overline{G}) &= \sum_{u \in V(G)} \deg_{\overline{G}}(u)^3 = \sum_{u \in V(G)} [n - 1 - \deg_G(u)]^3 \\ &= \sum_{u \in V(G)} [n^3 + \deg_G(u)^2(3n - 3) - \deg_G(u)(3n^2 - 6n + 3) \\ &\quad - 3n^2 + 3n - 1 - \deg_G(u)^3] \\ &= n^4 + M_1(G)(3n - 3) - 2m(3n^2 - 6n + 3) - n(3n^2 - 3n + 1) - F(G). \quad \square \end{aligned}$$

Theorem 2.2. *Let G be an (n, m) -graph. Then*

$$F(G) \leq n\Delta^3 + 3\Delta M_1(G) - 6m\Delta^2 \text{ and } F(G) \geq n\delta^3 + 3\delta M_1(G) - 6m\delta^2,$$

with equalities if and only if G is regular.

Proof. Define an auxiliary function $Y_1(G) = \sum_{u \in V(G)} [\deg(u) - k]^3$, where k is a real number. Then,

$$\begin{aligned} Y_1(G) &= \sum_{u \in V(G)} [\deg(u)^3 - k^3 - 3\deg(u)^2k + 3\deg(u)k^2] \\ &= F(G) - nk^3 - 3kM_1(G) + 6mk^2. \end{aligned}$$

If $k = \Delta$, then $Y_1(G) \leq 0$ and $F(G) \leq n\Delta^3 + 3\Delta M_1(G) - 6m\Delta^2$. For $k = \delta$, $Y_1(G) \geq 0$ and $F(G) \geq n\delta^3 + 3\delta M_1(G) - 6m\delta^2$. The equalities hold if and only if G is regular. \square

Theorem 2.3. *Let G be an (n, m) -graph. Then*

$$F(G) \geq M_1(G)(\delta + 2\Delta) - \Delta^2(2m - n\delta) - 4m\Delta\delta$$

and

$$F(G) \leq M_1(G)(\Delta + 2\delta) - \delta^2(2m - n\Delta) - 4m\delta\Delta$$

with equalities if and only if G is (Δ, δ) -biregular.

Proof. Define $Y_2(G) = \sum_{u \in V(G)} [\deg(u) - k]^2 [\deg(u) - h]$, where k and h are real numbers. Then,

$$\begin{aligned} Y_2(G) &= \sum_{u \in V(G)} [\deg(u)^2 + k^2 - 2\deg(u)k] [\deg(u) - h] \\ &= \sum_{u \in V(G)} [\deg(u)^3 - \deg(u)^2h + \deg(u)k^2 - k^2h - 2\deg(u)^2k + 2\deg(u)kh] \\ &= F(G) - M_1(G)(h + 2k) + k^2(2m - nh) + 4mkh. \end{aligned}$$

If $k = \Delta$ and $h = \delta$, then $Y_2(G) \geq 0$ and $F(G) \geq M_1(G)(\delta + 2\Delta) - \Delta^2(2m - n\delta) - 4m\Delta\delta$. For $k = \delta$ and $h = \Delta$, we have $Y_2(G) \leq 0$ and $F(G) \leq M_1(G)(\Delta + 2\delta) - \delta^2(2m - n\Delta) - 4m\delta\Delta$. The equalities hold if and only if G is (Δ, δ) -biregular. \square

Theorem 2.4. *Let G be an (n, m) -graph. Then $F(G) \geq 2[M_1(G) + m - n]$. If G is connected, then equality holds if and only if $G \cong P_n$ or $G \cong C_n$.*

Proof. Define the auxiliary function $Y_3(G) = \sum_{u \in V(G)} [\deg(u)^2 - 1][\deg(u) - 2]$ and note that $Y_3(G) = 0$ if and only if $\Delta(G) \leq 2$. In case of connected graphs, this will occur if either $G \cong P_n$ or $G \cong C_n$.

Now,

$$\begin{aligned} Y_3(G) &= \sum_{u \in V(G)} [\deg(u)^3 - 2\deg(u)^2 - \deg(u) + 2] \\ &= F(G) - 2M_1(G) - 2m + 2n. \end{aligned}$$

Since $Y_3(G) \geq 0$, $F(G) \geq 2[M_1(G) + m - n]$ with equality for connected graphs if and only if $G \cong P_n$ or $G \cong C_n$. \square

Theorem 2.5. *Let G be an (n, m) -graphs. Then*

$$F(G) \leq (3\Delta - 3)M_1(G) - 2m(3\Delta^2 - 6\Delta + 2) + n\Delta(\Delta - 1)(\Delta - 2)$$

and

$$F(G) \geq (3\delta + 3)M_1(G) - 2m(3\delta^2 + 6\delta + 2) + n\delta(\delta + 1)(\delta + 2).$$

The equalities holds if and only if G is $(\delta, \delta + 1, \delta + 2)$ -triregular.

Proof. Define $Y_4(G) = \sum_{u \in V(G)} [\deg(u) - a][\deg(u) - b][\deg(u) - c]$, where a, b , and c are real numbers. Then,

$$\begin{aligned} Y_4(G) &= \sum_{u \in V(G)} [\deg(u)^3 - \deg(u)^2(a + b + c) + \deg(u)(ab + ac + bc) - abc] \\ &= F(G) - (a + b + c)M_1(G) + 2m(ab + ac + bc) - nabc. \end{aligned}$$

If $a = \Delta$, $b = \Delta - 1$ and $c = \Delta - 2$, then $Y_4(G) \leq 0$ and $F(G) \leq (3\Delta - 3)M_1(G) - 2m(3\Delta^2 - 6\Delta + 2) + n\Delta(\Delta - 1)(\Delta - 2)$. For $a = \delta$, $b = \delta + 1$ and $c = \delta + 2$, $Y_4(G) \geq 0$ and $F(G) \geq (3\delta + 3)M_1(G) - 2m(3\delta^2 + 6\delta + 2) + n\delta(\delta + 1)(\delta + 2)$. The equalities hold if and only if G is $(\delta, \delta + 1, \delta + 2)$ -triregular. \square

For the sake of completeness, we mention here a result from [18].

Theorem 2.6. [18] *Let G be an (n, m) -graph. Then for $\alpha \geq 1$,*

$$\overline{M_1^{\alpha+1}}(G) = (n - 1)M_1^\alpha(G) - M_1^{\alpha+1}(G).$$

Theorem 2.7. *Let G be an (n, m) -graph. Then*

$$\overline{F}(G) \geq 2m[2\Delta(n - 1) + 3\Delta^2] - n[(n - 1)\Delta^2 + \Delta^3] - 3\Delta M_1(G).$$

The equality holds if and only if G is regular.

Proof. Define

$$Y_5(G) = (n - 1) \sum_{u \in V(G)} [\deg(u) - \Delta]^2 - \sum_{u \in V(G)} [\deg(u) - \Delta]^3.$$

Then,

$$\begin{aligned} Y_5(G) &= (n-1) \sum_{u \in V(G)} [\deg(u)^2 + \Delta^2 - 2\Delta \deg(u)] \\ &\quad - \sum_{u \in V(G)} [\deg(u)^3 - \Delta^3 - 3\Delta \deg(u)^2 + 3\Delta^2 \deg(u)] \\ &= (n-1)M_1(G) - F(G) + n[(n-1)\Delta^2 + \Delta^3] \\ &\quad - 2m[2\Delta(n-1) + 3\Delta^2] + 3\Delta M_1(G). \end{aligned}$$

Since $Y_5(G) \geq 0$, one can see that

$$(n-1)M_1(G) - F(G) \geq 2m[2\Delta(n-1) + 3\Delta^2] - n[(n-1)\Delta^2 + \Delta^3] - 3\Delta M_1(G).$$

The equality holds if and only if G is a regular graph. Therefore, by Theorem 2.6,

$$\bar{F}(G) \geq 2m[2\Delta(n-1) + 3\Delta^2] - n[(n-1)\Delta^2 + \Delta^3] - 3\Delta M_1(G)$$

with equality if and only if G is regular. □

Theorem 2.8. *Let G be an (n, m) -graph. Then*

$$\begin{aligned} \bar{F}(G) &\geq 2m[(n-1)(2\Delta-1) + \Delta^2 + 2\Delta(\Delta-1)] - M_1(G)(3\Delta-1) \\ &\quad - n[(n-1)\Delta(\Delta-1) + \Delta^2(\Delta-1)]. \end{aligned}$$

The equality holds if and only if G is $(\Delta, \Delta-1)$ -biregular.

Proof. We define the auxiliary function

$$\begin{aligned} Y_6(G) &= (n-1) \sum_{u \in V(G)} [\deg(u) - \Delta][\deg(u) - (\Delta-1)] \\ &\quad - \sum_{u \in V(G)} [\deg(u) - \Delta]^2 [\deg(u) - (\Delta-1)]. \end{aligned}$$

Then,

$$\begin{aligned} Y_6(G) &= (n-1) \sum_{u \in V(G)} [\deg(u)^2 - \deg(u)(2\Delta-1) + \Delta(\Delta-1)] \\ &\quad - \sum_{u \in V(G)} [\deg(u)^3 - \deg(u)^2(3\Delta-1) + \deg(u)\Delta^2 \\ &\quad \quad - \Delta^2(\Delta-1) + 2\deg(u)\Delta(\Delta-1)] \\ &= (n-1)M_1(G) - 2m(n-1)(2\Delta-1) + n(n-1)\Delta(\Delta-1) \\ &\quad - F(G) + M_1(G)(3\Delta-1) - 2m\Delta^2 + n\Delta^2(\Delta-1) - 4m\Delta(\Delta-1) \\ &= (n-1)M_1(G) - F(G) - 2m[(n-1)(2\Delta-1) + \Delta^2 + 2\Delta(\Delta-1)] \\ &\quad + n[(n-1)\Delta(\Delta-1) + \Delta^2(\Delta-1)] + M_1(G)(3\Delta-1). \end{aligned}$$

Since $Y_6(G) \geq 0$,

$$(n-1)M_1(G) - F(G) \geq 2m \left[(n-1)(2\Delta-1) + \Delta^2 + 2\Delta(\Delta-1) \right] \\ - n \left[(n-1)\Delta(\Delta-1) + \Delta^2(\Delta-1) \right] - (3\Delta-1)M_1(G),$$

with equality if and only if G is a $(\Delta, \Delta-1)$ -biregular graph. We now apply Theorem 2.6 to show that

$$\overline{F}(G) \geq 2m \left[(n-1)(2\Delta-1) + \Delta^2 + 2\Delta(\Delta-1) \right] \\ - n \left[(n-1)\Delta(\Delta-1) + \Delta^2(\Delta-1) \right] - (3\Delta-1)M_1(G)$$

with equality if and only if G is $(\Delta, \Delta-1)$ -biregular. \square

Theorem 2.9. *Let G be an (n, m) -graph. Then*

$$\overline{F}(G) \leq 2m \left[(n-1)(\delta + \Delta) + \Delta^2 + 2\Delta\delta \right] - n \left[(n-1)\Delta\delta + \Delta^2\delta \right] - (\delta + 2\Delta)M_1(G).$$

The equality holds if and only if G is (Δ, δ) -biregular.

Proof. Define the function

$$Y_7(G) = (n-1) \sum_{u \in V(G)} [\deg(u) - \Delta][\deg(u) - \delta] - \sum_{u \in V(G)} [\deg(u) - \Delta]^2 [\deg(u) - \delta].$$

Then,

$$Y_7(G) = (n-1) \sum_{u \in V(G)} \left[\deg(u)^2 - \deg(u)(\delta + \Delta) + \Delta\delta \right] \\ - \sum_{u \in V(G)} \left[\deg(u)^3 - \deg(u)^2(\delta + 2\Delta) + \deg(u)\Delta^2 - \Delta^2\delta + 2\deg(u)\Delta\delta \right] \\ = (n-1)M_1(G) - 2m(n-1)(\delta + \Delta) + n(n-1)\Delta\delta \\ - F(G) + M_1(G)(\delta + 2\Delta) - 2m\Delta^2 + n\Delta^2\delta - 4m\Delta\delta \\ = (n-1)M_1(G) - F(G) - 2m \left[(n-1)(\delta + \Delta) + \Delta^2 + 2\Delta\delta \right] \\ + n \left[(n-1)\Delta\delta + \Delta^2\delta \right] + (\delta + 2\Delta)M_1(G).$$

Since $Y_7(G) \leq 0$,

$$(n-1)M_1(G) - F(G) \leq 2m \left[(n-1)(\delta + \Delta) + \Delta^2 + 2\Delta\delta \right] \\ - n \left[(n-1)\Delta\delta + \Delta^2\delta \right] - (\delta + 2\Delta)M_1(G),$$

and the equality holds if and only if G is a (Δ, δ) -biregular graph. We now apply Theorem 2.6 to show that,

$$\overline{F}(G) \leq 2m \left[(n-1)(\delta + \Delta) + \Delta^2 + 2\Delta\delta \right] - n \left[(n-1)\Delta\delta + \Delta^2\delta \right] - (\delta + 2\Delta)M_1(G),$$

with equality holding if and only if G is (Δ, δ) -biregular. \square

Theorem 2.10. *Let G be an (n, m) -graph. Then the following holds.*

- (a) $M_1(G) \leq 2m(\delta + \Delta) - n\Delta\delta$, with equality if and only if G is (Δ, δ) -biregular.
 (b) $M_1(G) \geq 2m(2\Delta - 1) - n\Delta(\Delta - 1)$ and $M_1(G) \geq 2m(2\delta + 1) - n\delta(\delta + 1)$. The equalities holds if and only if G is $(\delta, \delta + 1)$ -biregular.
 (c) Let r be a real number. Then $M_1(G) \geq 4ma - nr^2$, with equality if and only if G is an r -regular graph.

Proof. Consider the function $Y_8(G) = \sum_{u \in V(G)} [\deg(u) - a][\deg(u) - b]$, where a and b are real numbers. Then we have,

$$\begin{aligned} Y_8(G) &= \sum_{u \in V(G)} [\deg(u)^2 - \deg(u)b - \deg(u)a + ab] \\ &= M_1(G) - 2m(a + b) + nab. \end{aligned}$$

If $a = \Delta$ and $b = \delta$, then $Y_8(G) \leq 0$ and $M_1(G) \leq 2m(\delta + \Delta) - n\Delta\delta$. Now the equality holds if and only if G is a (Δ, δ) -biregular graph. This completes the part (a).

Suppose that $a = \Delta$ and $b = \Delta - 1$. Then $Y_8(G) \geq 0$ and $M_1(G) \geq 2m(2\Delta - 1) - n\Delta(\Delta - 1)$. For $a = \delta$ and $b = \delta + 1$, $Y_8(G) \geq 0$ and $M_1(G) \geq 2m(2\delta + 1) - n\delta(\delta + 1)$. The equalities hold if and only if G is $(\delta, \delta + 1)$ -biregular, which completes the proof of part (b).

Finally, assume that $a = b = r$. Then $Y_8(G) \geq 0$ and $M_1(G) \geq 4ma - nr^2$. The equality holds if and only if G is r -regular. \square

Acknowledgements. The research of the first two authors was partially supported by the University of Kashan under grant no 364988/111.

REFERENCES

- [1] H. Abdo, D. Dimitrov and I. Gutman, *On extremal trees with respect to the F-index*, Kuwait J. Sci. **44**(4) (2017), 1–8.
- [2] S. Akhter and M. Imran, *Computing the forgotten topological index of four operations on graphs*, AKCE Int. J. Graphs Comb. **14** (2017), 70–79.
- [3] S. Akhter, M. Imran and M. R. Farahani, *Extremal unicyclic and bicyclic graphs with respect to the F-index*, AKCE Int. J. Graphs Comb. **14** (2017), 80–91.
- [4] B. Furtula and I. Gutman, *A forgotten topological index*, J. Math. Chem. **53** (2015), 1184–1190.
- [5] B. Borovićanin, K. C. Das, B. Furtula and I. Gutman, *Bounds for Zagreb indices*, MATCH Commun. Math. Comput. Chem. **78** (2017), 17–100.
- [6] Z. Che and Z. Chen, *Lower and upper bounds of the forgotten topological index*, MATCH Commun. Math. Comput. Chem. **76** (2016), 635–648.
- [7] N. De, S. M. Abu Nayeem and A. Pal, *F-index of some graph operations*, Discrete Math. Algorithms Appl. **8** (2016), ID 1650025.
- [8] N. De, S. M. Abu Nayeem and A. Pal, *The F coindex of some graph operations*, Springer Plus **5** (2016), Paper ID 221.
- [9] W. Gao, M. K. Siddiqui, M. Imran, M. K. Jamil and M. R. Farahani, *Forgotten topological index of chemical structure in drugs*, Saudi Pharma. J. **24** (2016), 258–264.
- [10] S. Ghobadi and M. Ghorbaninejad, *The forgotten topological index of four operations on some special graphs*, Bulletin of Mathematical Sciences and Applications **16** (2016), 89–95.

- [11] I. Gutman and K. C. Das, *The first Zagreb index 30 years after*, MATCH Commun. Math. Comput. Chem. **50** (2004), 83–92.
- [12] I. Gutman, A. Ghalavand, T. Dehghan-Zadeh and A. R. Ashrafi, *Graphs with smallest forgotten index*, Iranian Journal of Mathematical Chemistry **8** (2017), 259–273.
- [13] I. Gutman and N. Trinajstić, *Graph theory and molecular orbitals. Total π -electron energy of alternant hydrocarbons*, Chemical Physics Letters **17** (1972), 535–538.
- [14] A. Khaksari and M. Ghorbani, *On the forgotten topological index*, Iranian Journal of Mathematical Chemistry **8** (2017), 327–338.
- [15] X. Li and H. Zhao, *Trees with the first three smallest and largest generalized topological indices*, MATCH Commun. Math. Comput. Chem. **50** (2004), 57–62.
- [16] I. Ž. Milovanović, E. I. Milovanović, I. Gutman and B. Furtula, *Some inequalities for the forgotten topological index*, International Journal of Applied Graph Theory **1** (2017), 1–15.
- [17] S. Nikolić, G. Kovačević, A. Miličević and N. Trinajstić, *The Zagreb indices 30 years after*, Croatica Chemica Acta **76** (2003), 113–124.
- [18] S. Zhang and H. Zhang, *Unicyclic graphs with the first three smallest and largest first general Zagreb index*, MATCH Commun. Math. Comput. Chem. **55** (2006), 427–438.

¹DEPARTMENT OF PURE MATHEMATICS,
FACULTY OF MATHEMATICAL SCIENCE,
UNIVERSITY OF KASHAN,
KASHAN 87317-51167, I. R. IRAN
Email address: ali.ghalavand.kh@gmail.com
Email address: ashrafi@kashanu.ac.ir

²FACULTY OF SCIENCE,
UNIVERSITY OF KRAGUJEVAC,
34000 KRAGUJEVAC, SERBIA
Email address: gutman@kg.ac.rs