

SEVERAL THEOREMS OF APPROXIMATION THEORY FOR THE q -BESSEL TRANSFORM

OTHMAN TYR¹ AND ABDELAALI DADES²

ABSTRACT. In this paper, we prove several theorems in approximation theory concerning the q -Bessel transform. A new estimate for the q -Bessel function is obtained. By applying the generalized modulus of continuity, we establish new Jackson-type inverse estimates for the generalized q -Bessel transform in the space $\mathcal{L}_{q,\alpha}^2(\mathbb{R}_q^+)$, where $q \in (0, 1)$ and $\alpha > -1/2$.

1. INTRODUCTION

Let $f : \mathbb{R} \rightarrow \mathbb{C}$ a Lebesgue square-integrable function on \mathbb{R} , i.e., $f \in L^2(\mathbb{R})$. We define the finite differences of order $k \in \mathbb{N}$ by

$$\Delta_h^k(f; x) = (F_h - I)^k f(x) = \sum_{i=0}^k (-1)^{k-i} \binom{k}{i} F_h^i f(x), \quad x \in \mathbb{R},$$

where F_h is the Steklov operator defined by

$$(1.1) \quad F_h f(x) = 2^{-1} h^{-1} \int_{x-h}^{x+h} f(t) dt, \quad h > 0,$$

and I is the unit operator in $L^2(\mathbb{R})$.

For a given positive real number δ , the k^{th} -order generalized modulus of continuity of a function f is defined by

$$\Omega_k(f, \delta) = \sup_{0 < h \leq \delta} \|\Delta_h^k(f; x)\|_{L^2(\mathbb{R})}.$$

Key words and phrases. q -Bessel transform, generalized q -Bessel translation, modulus of continuity.
2020 Mathematics Subject Classification. Primary: 26B10. Secondary: 33D15.

DOI

Received: February 14, 2025.

Accepted: August 03, 2025.

Let $W_{2,\Phi}^{r,k}(D)$, where $r = 0, 1, \dots$ and $k = 1, 2, \dots$, denote the class of functions $f \in L^2(\mathbb{R})$ for which the generalized partial derivatives exist in the sense of Levi:

$$\frac{\partial f}{\partial x}, \frac{\partial^2 f}{\partial x^2}, \dots, \frac{\partial^r f}{\partial x^r},$$

which all belong to $L^2(\mathbb{R})$ (see [21, p. 172]). These functions satisfy the estimate

$$\Omega_k(D^r f, \delta) = O(\Phi(\delta^k)), \quad \text{as } \delta \rightarrow 0,$$

where $D = \frac{\partial}{\partial x}$, and the iterated derivatives $D^i f$ are defined recursively by

$$D^0 f = f, \quad D^i f = D(D^{i-1} f), \quad i = 1, 2, \dots, r.$$

Here, Φ is a continuous, strictly increasing function on $[0, +\infty)$ with $\Phi(0) = 0$.

The following theorem is an analogue of Jackson's direct theorem from the classical theory of function approximation [21, Ch. 5] (see also [1, Theorem 1]).

Theorem 1.1. *It holds that*

$$\sup_{f \in W_{2,\Phi}^{r,k}(D)} \sqrt{\int_{|\lambda| \geq N} |\hat{f}(\lambda)|^2 d\lambda} = O\left(N^{-r} \Phi\left[\left(\frac{2}{N}\right)^k\right]\right),$$

as $N \rightarrow +\infty$, where $r = 0, 1, \dots$, $k = 1, 2, \dots$, and \hat{f} stands for the Fourier transform of f .

In the case where $\Phi(t) = t^\nu$ with $\nu > 0$, Abilov et al. characterized the functions $f \in L^2(\mathbb{R})$ by the following equivalence (see [1, Theorem 2]).

Theorem 1.2. *Let $\Phi(t) = t^\nu$ with $\nu > 0$. Then, the assertions $f \in W_{2,t^\nu}^{r,k}(D)$ and*

$$\sqrt{\int_{|\lambda| \geq N} |\hat{f}(\lambda)|^2 d\lambda} = O(N^{-r-k\nu}), \quad \text{as } N \rightarrow +\infty,$$

are equivalent, where $r = 0, 1, \dots$, $k = 1, 2, \dots$, and $0 < \nu < 2$.

Recently, considerable attention has been paid to the development of various q -analogues of Fourier analysis using tools from quantum calculus (see [4, 5, 8, 9, 12] and references therein). In the context of q -Bessel analysis, many works have focussed on the construction of q -Fourier analysis associated with the q -Hankel transform. This theory was first introduced by Koornwinder and Swarttouw [19], and later extended by Fitouhi et al. [9, 10, 13]. It is therefore natural to investigate q -analogues of classical theorems in this framework.

In this paper, building on the work of Abilov, we establish new Jackson-type inverse estimates for the q -Bessel transform for certain classes of functions characterised by a modulus of continuity and associated with the q -Bessel operator. Similar results were obtained in [2, 6, 7, 25–30]. To achieve these results, we use a generalised q -Bessel translation operator rather than the Steklov operator defined in (1.1).

2. HARMONIC ANALYSIS ASSOCIATED WITH THE q -BESSEL OPERATOR

This section provides the essential background to q -harmonic analysis related to the q -Bessel transform. We give a brief overview of the standard notions and notations of q -theory, while further details and results on q -Bessel analysis can be found in [5, 9–11, 13, 17, 18].

Throughout this paper, we assume $q \in]0, 1[$ and $\alpha > -1/2$. We introduce the following set

$$\mathbb{R}_q^+ = \{q^n, n \in \mathbb{Z}\}.$$

Let $a \in \mathbb{C}$, the q -shifted factorials are defined by:

$$(2.1) \quad (a; q)_0 = 1, \quad (a; q)_n = \prod_{l=0}^{n-1} (1 - aq^l), \quad n = 1, 2, \dots, \quad (a; q)_\infty = \prod_{l=0}^{+\infty} (1 - aq^l).$$

The q -derivative of a function f is given by

$$\mathcal{D}_q f(x) = \frac{f(x) - f(qx)}{(1 - q)x}, \quad \text{if } x \neq 0,$$

where $\mathcal{D}_q f(0) = f'(0)$ provided $f'(0)$ exists. Note that when f is differentiable at x , then $\mathcal{D}_q f(x)$ tends to $f'(x)$ as q tends to 1^- .

The general q -hypergeometric series is defined by

$$(2.2) \quad {}_r\phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r \\ b_1, b_2, \dots, b_s \end{matrix}; q, z \right] = {}_r\phi_s (a_1, a_2, \dots, a_r; b_1, b_2, \dots, b_s; q, z) \\ = \sum_{n=0}^{+\infty} \frac{(a_1, a_2, \dots, a_r; q)_n}{(b_1, b_2, \dots, b_s; q)_n (q; q)_n} \left[(-1)^n q^{n(n-1)/2} \right]^{1+s-r} z^n,$$

where the q -shifted factorial is defined by (2.1) and

$$(a_1, a_2, \dots, a_r; q)_n = (a_1; q)_n (a_2; q)_n \cdots (a_r; q)_n.$$

The q -Jackson integrals from 0 to a and from 0 to $+\infty$ are defined by [14, 17]

$$\int_0^a f(x) d_q x = (1 - q)a \sum_{n=0}^{+\infty} q^n f(aq^n), \\ \int_0^{+\infty} f(x) d_q x = (1 - q) \sum_{n=-\infty}^{+\infty} q^n f(q^n),$$

provided the sums converge absolutely. The q -Jackson integral in a generic interval $[a, b]$ is given by

$$\int_a^b f(x) d_q x := \int_0^b f(x) d_q x - \int_0^a f(x) d_q x.$$

We denote by $\mathcal{L}_{q,\alpha}^p(\mathbb{R}_q^+)$, $p \geq 1$, the set of all real functions f defined on \mathbb{R}_q^+ for which

$$\|f\|_{q,p,\alpha} = \left[\int_0^{+\infty} |f(x)|^p x^{2\alpha+1} d_q x \right]^{1/p} < +\infty.$$

We denote by $\mathcal{C}_{q,0}(\mathbb{R}_q^+)$, for the space of functions defined on \mathbb{R}_q^+ tending to 0 as $x \rightarrow +\infty$ and continuous at 0. The space $\mathcal{C}_{q,0}(\mathbb{R}_q^+)$, when equipped with the topology of uniform convergence, is a complete normed linear space with norm

$$\|f\|_{q,\infty} = \sup_{x \in \mathbb{R}_q^+} |f(x)|.$$

In [16], Ismail introduced the third q -Bessel function, defined as follows

$$\begin{aligned} J_\alpha(x; q^2) &= \frac{(q^{2\alpha+2}; q^2)_\infty}{(q^2; q^2)_\infty} x^\alpha {}_1\phi_1 \left(0; q^{2\alpha+2}; q^2, q^2 x^2 \right) \\ (2.3) \quad &= \frac{(q^{2\alpha+2}; q^2)_\infty}{(q^2; q^2)_\infty} \sum_{n=0}^{+\infty} (-1)^n \frac{q^{n(n+1)}}{(q^{2\alpha+2}; q^2)_n (q^2; q^2)_n} x^{2n+\alpha}. \end{aligned}$$

It satisfies (see [5]) the following.

- For all $n \in \mathbb{Z}$, $\alpha \geq -1/2$, we have

$$(2.4) \quad |J_\alpha(q^n; q^2)| \leq \frac{(-q^2; q^2)_\infty (-q^{2\alpha+2}; q^2)_\infty}{(q^2; q^2)_\infty} \begin{cases} q^{n\alpha}, & \text{if } n \geq 0, \\ q^{n(n-\alpha-1)}, & \text{if } n < 0. \end{cases}$$

- For all $\sigma \in \mathbb{R}$, $x \in \mathbb{R}_q^+$, we have

$$J_\alpha(x; q^2) = o(x^{-\sigma}), \quad \text{as } x \rightarrow +\infty.$$

In particular

$$J_\alpha(x; q^2) \rightarrow 0, \quad \text{as } x \rightarrow +\infty.$$

The function $J_\alpha(\cdot; q^2)$ is given in normalized form by

$$(2.5) \quad j_\alpha(x; q^2) = \sum_{n=0}^{+\infty} (-1)^n \frac{q^{n(n+1)}}{(q^{2\alpha+2}; q^2)_n (q^2; q^2)_n} x^{2n}.$$

The following lemma was proved in [12, Lemma 3.1].

Lemma 2.1. *We have the following inequalities for the q -Bessel function:*

$$(2.6) \quad j_\alpha(x; q^2) = O(1), \quad \text{if } x \geq 0 \text{ and } x \in \mathbb{R}_q^+,$$

$$(2.7) \quad 1 - j_\alpha(x; q^2) = O(1), \quad \text{if } x \geq 1 \text{ and } x \in \mathbb{R}_q^+,$$

$$(2.8) \quad 1 - j_\alpha(x; q^2) = O(x^2), \quad \text{if } x \leq 1 \text{ and } x \in \mathbb{R}_q^+.$$

The q -integration theorem by a change of variable can be stated as follows

$$\int_a^b H \left(\frac{\lambda}{s} \right) \lambda^{2\alpha+1} d_q \lambda = s^{2\alpha+2} \int_{\frac{a}{s}}^{\frac{b}{s}} H(x) x^{2\alpha+1} d_q x, \quad \text{for all } s \in \mathbb{R}_q^+.$$

For $\lambda \in \mathbb{C}$, the function $x \mapsto j_\alpha(\lambda x, q^2)$ is a solution of the q -differential equation

$$\begin{cases} \Lambda_{q,\alpha} f(x) = -\lambda^2 f(x), \\ f(0) = 1, \end{cases}$$

where $\Lambda_{q,\alpha}$ denotes the q -Bessel operator, defined by

$$\Lambda_{q,\alpha}f(x) = \frac{f(q^{-1}x) - (1 + q^{2\alpha})f(x) + q^{2\alpha}f(qx)}{x^2}, \quad x \in \mathbb{R}_q^+.$$

We now give the definition of the q -Bessel transform associated with the q -Bessel operator.

Definition 2.1. The q -Bessel transform $\mathcal{F}_{q,\alpha}$, is defined for every function $f \in \mathcal{L}_{q,\alpha}^1(\mathbb{R}_q^+)$ by

$$\mathcal{F}_{q,\alpha}(f)(\lambda) = c_{q,\alpha} \int_0^{+\infty} f(x) j_\alpha(\lambda x; q^2) x^{2\alpha+1} d_q x, \quad \text{for all } \lambda \in \mathbb{R}_q^+,$$

where

$$c_{q,\alpha} = \frac{1}{1-q} \cdot \frac{(q^{2\alpha+2}; q^2)_\infty}{(q^2; q^2)_\infty}.$$

It is well-known that the q -Bessel transform satisfies the following properties (see [10, 13, 15]).

- If $f \in \mathcal{L}_{q,\alpha}^1(\mathbb{R}_q^+)$, then $\mathcal{F}_{q,\alpha}(f) \in \mathcal{C}_{q,0}(\mathbb{R}_q^+)$ and $\|\mathcal{F}_{q,\alpha}(f)\|_{q,\infty} \leq \mathcal{B}_{q,\alpha} \|f\|_{q,1,\alpha}$, where

$$\mathcal{B}_{q,\alpha} = \frac{1}{1-q} \cdot \frac{(-q^2; q^2)_\infty (-q^{2\alpha+2}; q^2)_\infty}{(q^2; q^2)_\infty}.$$

- For every $f \in \mathcal{L}_{q,\alpha}^2(\mathbb{R}_q^+)$, one has

$$(2.9) \quad \mathcal{F}_{q,\alpha}(\Lambda_{q,\alpha}f)(\lambda) = -\lambda^2 \mathcal{F}_{q,\alpha}(f)(\lambda), \quad \text{for all } \lambda \in \mathbb{R}_q^+.$$

Theorem 2.1 gives the Plancherel formula and inversion formula for the q -Bessel transform.

Theorem 2.1. *i) The q -Bessel transform $\mathcal{F}_{q,\alpha}$ is an isomorphism from $\mathcal{L}_{q,\alpha}^2(\mathbb{R}_q^+)$ onto itself and satisfies the q -Plancherel formula*

$$(2.10) \quad \|\mathcal{F}_{q,\alpha}(f)\|_{q,2,\alpha} = \|f\|_{q,2,\alpha}, \quad \text{for all } f \in \mathcal{L}_{q,\alpha}^2(\mathbb{R}_q^+).$$

ii) If $f \in \mathcal{L}_{q,\alpha}^1(\mathbb{R}_q^+)$ such that $\mathcal{F}_{q,\alpha}(f) \in \mathcal{L}_{q,\alpha}^1(\mathbb{R}_q^+)$, then the q -inversion formula holds and we have

$$(2.11) \quad f(x) = c_{q,\alpha} \int_0^{+\infty} \mathcal{F}_{q,\alpha}(f)(\lambda) j_\alpha(\lambda x; q^2) \lambda^{2\alpha+1} d_q \lambda, \\ \text{a.e. on } \mathbb{R}_q^+.$$

The q -generalized translation operator associated with the q -Bessel transform, denoted by $\mathcal{T}_{q,h}^\alpha$ for $h \in \mathbb{R}_q^+$, was introduced in [13] and later corrected in [10]. It is defined using Jackson's q -integral and the q -shifted factorial as

$$\mathcal{T}_{q,h}^\alpha f(x) = \int_0^{+\infty} f(t) \mathcal{K}_{q,\alpha}(h, x, t) t^{2\alpha+1} d_q t,$$

where

$$\mathcal{K}_{q,\alpha}(h, x, y) = c_{q,\alpha}^2 \int_0^{+\infty} j_\alpha(ht; q^2) j_\alpha(xt; q^2) j_\alpha(yt; q^2) t^{2\alpha+1} d_q t.$$

In particular, the product formula

$$\mathcal{T}_{q,h}^\alpha j_\alpha(x; q^2) = j_\alpha(h; q^2) j_\alpha(x; q^2)$$

holds.

The q -generalized translation operator has the following properties (see [10, 11]).

Theorem 2.2. *i) For $f \in \mathcal{L}_{q,\alpha}^p(\mathbb{R}_q^+)$, $p \geq 1$, we have $\mathcal{T}_{q,h}^\alpha f \in \mathcal{L}_{q,\alpha}^p(\mathbb{R}_q^+)$ and*

$$\|\mathcal{T}_{q,h}^\alpha f\|_{q,p,\alpha} \leq \|f\|_{q,p,\alpha}.$$

ii) For $f \in \mathcal{L}_{q,\alpha}^1(\mathbb{R}_q^+)$, we have

$$(2.12) \quad \mathcal{F}_{q,\alpha}(\mathcal{T}_{q,h}^\alpha f)(\lambda) = j_\alpha(\lambda h; q^2) \mathcal{F}_{q,\alpha}(f)(\lambda).$$

For every $f \in \mathcal{L}_{q,\alpha}^2(\mathbb{R}_q^+)$, we define the differences $\Delta_h^m f$ of order m , $m = 1, 2, \dots$, with step $h > 0$, $h \in \mathbb{R}_q^+$ by:

$$\begin{aligned} \Delta_h^1 f(x) &= \Delta_h f(x) := \mathcal{T}_{q,h}^\alpha f(x) - f(x), \\ \Delta_h^m f(x) &= \Delta_h(\Delta_h^{m-1} f(x)), \quad \text{for } m \geq 2. \end{aligned}$$

The m^{th} -order generalized modulus of continuity of $f \in \mathcal{L}_{q,\alpha}^2(\mathbb{R}_q^+)$ is defined by

$$\omega_m(f, \delta)_{q,2,\alpha} = \sup_{0 < h \leq \delta} \|\Delta_h^m f\|_{q,2,\alpha}, \quad \delta > 0.$$

Let $W_{q,2}^r(\Lambda_{q,\alpha})$, $r = 0, 1, \dots$, denote the class of functions $f \in \mathcal{L}_{q,\alpha}^2(\mathbb{R}_q^+)$ that have generalized derivatives in the sense of distributions satisfying

$$\Lambda_{q,\alpha}^r f \in \mathcal{L}_{q,\alpha}^2(\mathbb{R}_q^+), \quad r = 1, 2, \dots,$$

i.e.,

$$W_{q,2}^r(\Lambda_{q,\alpha}) := \left\{ f \in \mathcal{L}_{q,\alpha}^2(\mathbb{R}_q^+) : \Lambda_{q,\alpha}^r f \in \mathcal{L}_{q,\alpha}^2(\mathbb{R}_q^+), \quad r = 1, 2, \dots \right\},$$

where

$$\Lambda_{q,\alpha}^0 f = f, \quad \Lambda_{q,\alpha}^r f = \Lambda_{q,\alpha}(\Lambda_{q,\alpha}^{r-1} f), \quad r = 1, 2, \dots$$

From (2.9), for all $f \in W_{q,2}^r(\Lambda_{q,\alpha})$, we have

$$(2.13) \quad \mathcal{F}_{q,\alpha}(\Lambda_{q,\alpha}^r f)(\lambda) = (-1)^r \lambda^{2r} \mathcal{F}_{q,\alpha}(f)(\lambda), \quad r = 1, 2, \dots$$

3. MAIN RESULTS

We now present the main results of this paper. Their proofs rely on several preliminary lemmas.

Lemma 3.1. *For $\alpha \geq -1/2$ and $x \in \mathbb{R}_q^+$, we have*

$$(3.1) \quad \sqrt{x} J_\alpha(x; q^2) = \mathcal{O}(1), \quad \text{for } x \geq 0.$$

Moreover, we have

$$(3.2) \quad j_\alpha(x; q^2) = \mathcal{O}(x^{-\alpha - \frac{1}{2}}), \quad \text{for } x \geq 0.$$

Proof. First, note that

$$x \in \mathbb{R}_q^+, x \geq 1 \quad \text{if and only if} \quad x = q^n, n \leq 0,$$

and

$$x \in \mathbb{R}_q^+, 0 \leq x \leq 1 \quad \text{if and only if} \quad x = q^n, n \geq 0.$$

From this and (2.4), we distinguish two cases.

For $x = q^n \in \mathbb{R}_q^+$ with $n \geq 0$, we obtain

$$|q^{\frac{n}{2}} J_\alpha(q^n; q^2)| \leq \frac{(-q^2; q^2)_\infty (-q^{2\alpha+2}; q^2)_\infty}{(q^2; q^2)_\infty} q^{n(\alpha+\frac{1}{2})}.$$

Since $n(\alpha + \frac{1}{2}) \geq 0$, it follows that $q^{n(\alpha+\frac{1}{2})} \leq 1$, and therefore we obtain

$$|q^{\frac{n}{2}} J_\alpha(q^n; q^2)| \leq \frac{(-q^2; q^2)_\infty (-q^{2\alpha+2}; q^2)_\infty}{(q^2; q^2)_\infty}.$$

Therefore, equivalently,

$$(3.3) \quad \sqrt{x} J_\alpha(x; q^2) = \mathcal{O}(1), \quad \text{for } x \in \mathbb{R}_q^+, 0 \leq x \leq 1.$$

For the case $x = q^n \in \mathbb{R}_q^+$ with $n \leq 0$, we get

$$|q^{\frac{n}{2}} J_\alpha(q^n; q^2)| \leq \frac{(-q^2; q^2)_\infty (-q^{2\alpha+2}; q^2)_\infty}{(q^2; q^2)_\infty} q^{n(n-\alpha-\frac{1}{2})}.$$

Since $n(n - \alpha - \frac{1}{2}) \geq 0$, it follows that

$$q^{n(n-\alpha-\frac{1}{2})} \leq 1,$$

and thus we again obtain

$$|q^{\frac{n}{2}} J_\alpha(q^n; q^2)| \leq \frac{(-q^2; q^2)_\infty (-q^{2\alpha+2}; q^2)_\infty}{(q^2; q^2)_\infty}.$$

Consequently,

$$(3.4) \quad \sqrt{x} J_\alpha(x; q^2) = \mathcal{O}(1), \quad \text{for } x \in \mathbb{R}_q^+, x \geq 1.$$

Now, combining (3.3) and (3.4), we obtain

$$\sqrt{x} J_\alpha(x; q^2) = \mathcal{O}(1), \quad \text{for } x \geq 0, x \in \mathbb{R}_q^+,$$

which proves (3.1). To prove (3.2), we use formulas (2.3) and (2.5), which yield

$$j_\alpha(x; q^2) = \frac{(q^2; q^2)_\infty}{(q^{2\alpha+2}; q^2)_\infty} x^{-\alpha} J_\alpha(x; q^2).$$

Thus, taking into account the formula above and (3.1), we obtain the result. \square

We now present an important lemma that will lead us to the main result.

Lemma 3.2. *Let $h > 0$ with $h \in \mathbb{R}_q^+$. If $f \in W_{q,2}^r(\Lambda_{q,\alpha})$, then*

$$\|\Delta_h^m(\Lambda_{q,\alpha}^r f)\|_{q,2,\alpha}^2 = \int_0^{+\infty} \lambda^{4r} |1 - j_\alpha(\lambda h, q^2)|^{2m} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda,$$

where $r = 0, 1, 2, \dots$

Proof. According to the formula (2.12), we obtain

$$\mathcal{F}_{q,\alpha}(\Delta_h f)(\lambda) = \mathcal{F}_{q,\alpha}(\mathcal{T}_{q,h}^\alpha f)(\lambda) - \mathcal{F}_{q,\alpha}(f)(\lambda) = (j_\alpha(\lambda h, q^2) - 1) \mathcal{F}_{q,\alpha}(f)(\lambda).$$

Using the recurrence proof with respect to m , we obtain

$$\mathcal{F}_{q,\alpha}(\Delta_h^m f)(\lambda) = (j_\alpha(\lambda h, q^2) - 1)^m \mathcal{F}_{q,\alpha}(f)(\lambda).$$

In view of formula (2.13), we get

$$\mathcal{F}_{q,\alpha}(\Delta_h^m(\Lambda_{q,\alpha}^r f))(\lambda) = (-1)^r \lambda^{2r} (j_\alpha(\lambda h, q^2) - 1)^m \mathcal{F}_{q,\alpha}(f)(\lambda).$$

Now, by appealing the q -Plancherel formula (2.10), we have the desired result. \square

The following theorem is an analogue of Abilov's theorem [1, Theorem 1]. It represents a Jackson-type direct theorem from the classical theory of function approximation [21, Chapter 5].

Theorem 3.1. *For a function $f \in \mathcal{L}_{q,\alpha}^2(\mathbb{R}_q^+)$ in the class $W_{q,2}^r(\Lambda_{q,\alpha})$ there exists a fixed constant $c > 0$ such that for all $N > 0$, we have*

$$\mathcal{J}_N(f) = \mathcal{O}\left(N^{-2r} \omega_m\left(\Lambda_{q,\alpha}^r f, \frac{c}{N}\right)_{q,2,\alpha}\right),$$

where $r = 0, 1, 2, \dots$, $m = 1, 2, \dots$

Proof. Firstly, for all $h \in \mathbb{R}_q^+$, we can see that

$$\begin{aligned} \mathcal{J}_N^2(f) &= \int_N^{+\infty} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda \\ &\leq \int_N^{+\infty} |j_\alpha(\lambda h, q^2)| \cdot |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda \\ (3.5) \quad &\quad + \int_N^{+\infty} |1 - j_\alpha(\lambda h, q^2)| \cdot |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda. \end{aligned}$$

According to the formula (3.2), there exists a constant $c_1 > 0$ such that

$$|j_\alpha(\lambda h, q^2)| \leq c_1(\lambda h)^{-\alpha-\frac{1}{2}}.$$

Hence,

$$\begin{aligned} \int_N^{+\infty} |j_\alpha(\lambda h, q^2)| \cdot |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda &\leq c_1 h^{-\alpha-\frac{1}{2}} \int_N^{+\infty} \lambda^{-\alpha-\frac{1}{2}} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda \\ &\leq c_1 (hN)^{-\alpha-\frac{1}{2}} \mathcal{J}_N^2(f). \end{aligned}$$

Choosing $c > 0$ such that the constant $c_2 = 1 - c_1 c^{-\alpha - \frac{1}{2}}$ is positive and setting $h = c/N$ in inequality (3.5). Then, we have

$$(3.6) \quad c_2 \mathcal{J}_N^2(f) \leq \int_N^{+\infty} |1 - j_\alpha(\lambda h, q^2)| \cdot |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda.$$

Furthermore, by Hölder's inequality, the second term in (3.6) satisfies

$$\begin{aligned} & \int_N^{+\infty} |1 - j_\alpha(\lambda h, q^2)| \cdot |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda \\ &= \int_N^{+\infty} |1 - j_\alpha(\lambda h, q^2)| \left(|\mathcal{F}_{q,\alpha}(f)(\lambda)| \lambda^{\alpha+\frac{1}{2}} \right)^{2-\frac{1}{m}} \left(|\mathcal{F}_{q,\alpha}(f)(\lambda)| \lambda^{\alpha+\frac{1}{2}} \right)^{\frac{1}{m}} d_q \lambda \\ &\leq \left(\int_N^{+\infty} |1 - j_\alpha(\lambda h, q^2)|^{2m} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda \right)^{\frac{1}{2m}} (\mathcal{J}_N(f))^{\frac{2m-1}{m}} \\ &= \left(\int_N^{+\infty} \frac{1}{\lambda^{4r}} \lambda^{4r} |1 - j_\alpha(\lambda h, q^2)|^{2m} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda \right)^{\frac{1}{2m}} (\mathcal{J}_N(f))^{\frac{2m-1}{m}} \\ &\leq N^{\frac{-2r}{m}} (\mathcal{J}_N(f))^{\frac{2m-1}{m}} \left(\int_N^{+\infty} \lambda^{4r} |1 - j_\alpha(\lambda h, q^2)|^{2m} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda \right)^{\frac{1}{2m}}. \end{aligned}$$

From Lemma 3.2, we conclude that

$$\int_N^{+\infty} \lambda^{4r} |1 - j_\alpha(\lambda h, q^2)|^{2m} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda \leq \|\Delta_h^m(\Lambda_{q,\alpha}^r f)\|_{q,2,\alpha}^2.$$

Therefore,

$$\int_N^{+\infty} |1 - j_\alpha(\lambda h, q^2)| \cdot |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda \leq N^{\frac{-2r}{m}} (\mathcal{J}_N(f))^{\frac{2m-1}{m}} \|\Delta_h^m(\Lambda_{q,\alpha}^r f)\|_{q,2,\alpha}^{\frac{1}{m}}.$$

For $h = c/N$, we obtain

$$c_2 \mathcal{J}_N^2(f) \leq N^{\frac{-2r}{m}} (\mathcal{J}_N(f))^{\frac{2m-1}{m}} \omega_m^{1/m} \left(\Lambda_{q,\alpha}^r f, \frac{c}{N} \right)_{q,2,\alpha}^{\frac{1}{m}}.$$

Consequently, by raising both sides to the power m and simplifying by $(\mathcal{J}_N(f))^{2m}$, we finally obtain

$$\mathcal{J}_N(f) \leq c_2^{-m} N^{-2r} \omega_m \left(\Lambda_{q,\alpha}^r f, \frac{c}{N} \right)_{q,2,\alpha}.$$

Therefore,

$$\mathcal{J}_N(f) = \mathcal{O} \left(N^{-2r} \omega_m \left(\Lambda_{q,\alpha}^r f, \frac{c}{N} \right)_{q,2,\alpha} \right).$$

This completes the proof of Theorem 3.1. \square

Theorems 3.2 and 3.3 are analogues of the classical inverse theorems of approximation theory due to Stechkin and Timan (see [22, 23]).

Theorem 3.2. *Let $f \in \mathcal{L}_{q,\alpha}^2(\mathbb{R}_q^+)$. Then, for all $N > 0$,*

$$\omega_m \left(f, \frac{1}{N} \right)_{q,2,\alpha} = \mathcal{O} \left(N^{-2m} \left(\sum_{l=0}^{N-1} (l+1)^{4m-1} \mathcal{J}_l^2(f) \right)^{\frac{1}{2}} \right).$$

Proof. From Lemma 3.2, we have

$$\|\Delta_h^m f\|_{q,2,\alpha}^2 = \int_0^{+\infty} |1 - j_\alpha(\lambda h, q^2)|^{2m} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda.$$

This integral is divided into two:

$$\int_0^{+\infty} = \int_0^N + \int_N^{+\infty} = I_1 + I_2,$$

where $N = [\frac{1}{h}]$. We estimate them separately. From (2.6), we have the estimate

$$(3.7) \quad I_2 \leq c_3 \int_N^{+\infty} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda = c_3 \mathcal{J}_N^2(f).$$

Now, we estimate I_1 . From relation (2.8), we have

$$\begin{aligned} I_1 &\leq c_4 h^{4m} \int_0^N \lambda^{4m} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda \\ &= c_4 h^{4m} \sum_{l=0}^{N-1} \int_l^{l+1} \lambda^{4m} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda \\ &\leq c_4 h^{4m} \sum_{l=0}^{N-1} a_l (\mathcal{J}_l^2(f) - \mathcal{J}_{l+1}^2(f)), \end{aligned}$$

where $a_l = (l+1)^{4m}$ and

$$\mathcal{J}_l^2(f) = \int_l^{+\infty} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda.$$

For all integers $M \geq 1$, the Abel transformation shows

$$\begin{aligned} \sum_{l=0}^M a_l (\mathcal{J}_l^2(f) - \mathcal{J}_{l+1}^2(f)) &= a_0 \mathcal{J}_0^2(f) - a_M \mathcal{J}_{M+1}^2(f) + \sum_{l=1}^M (a_l - a_{l-1}) \mathcal{J}_l^2(f) \\ &\leq \mathcal{J}_0^2(f) + \sum_{l=1}^M (a_l - a_{l-1}) \mathcal{J}_l^2(f), \end{aligned}$$

because $a_M \mathcal{J}_{M+1}^2(f) \geq 0$. Hence,

$$I_1 \leq c_4 h^{4m} \left(\mathcal{J}_0^2(f) + \sum_{l=1}^{N-1} (a_l - a_{l-1}) \mathcal{J}_l^2(f) \right).$$

Moreover by the finite increments theorem, we have

$$a_l - a_{l-1} \leq 4m(l+1)^{4m-1}.$$

Then,

$$I_1 \leq c_4 N^{-4m} \left(\mathcal{J}_0^2(f) + 4m \sum_{l=1}^{N-1} (l+1)^{4m-1} \mathcal{J}_l^2(f) \right),$$

since $N \leq \frac{1}{h}$. Combining the estimates for I_1 and I_2 gives

$$\|\Delta_h^m f\|_{q,2,\alpha}^2 = \mathcal{O} \left(N^{-4m} \sum_{l=0}^{N-1} (l+1)^{4m-1} \mathcal{J}_l^2(f) \right),$$

which implies

$$\omega_m \left(f, \frac{1}{N} \right)_{q,2,\alpha} = \mathcal{O} \left(N^{-2m} \left(\sum_{l=0}^{N-1} (l+1)^{4m-1} \mathcal{J}_l^2(f) \right)^{\frac{1}{2}} \right),$$

and this completes the proof. \square

Theorem 3.3. *Let $f \in \mathcal{L}_{q,\alpha}^2(\mathbb{R}_q^+)$. If the series*

$$\sum_{l=1}^{+\infty} l^{2r-1} \mathcal{J}_l(f), \quad r = 1, 2, \dots$$

converges, then $f \in W_{q,2}^r(\Lambda_{q,\alpha})$ and, for all $N > 0$,

$$\omega_m \left(\Lambda_{q,\alpha}^r f, \frac{1}{N} \right)_{q,2,\alpha} = \mathcal{O} \left(N^{-4m} \sum_{l=0}^{N-1} (l+1)^{4r+4m-1} \mathcal{J}_l^2(f) \right)^{\frac{1}{2}} + \mathcal{O} \left(\sum_{l=\lceil \frac{N}{2} \rceil}^{+\infty} l^{2r-1} \mathcal{J}_l(f) \right).$$

Proof. Let $f \in \mathcal{L}_{q,\alpha}^2(\mathbb{R}_q^+)$. By (2.13) and Plancherel formula (2.10), we have

$$\begin{aligned} \|\Lambda_{q,\alpha}^r f\|_{q,2,\alpha}^2 &= \int_0^{+\infty} \lambda^{4r} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda \\ &= \sum_{l=0}^{+\infty} \int_l^{l+1} \lambda^{4r} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda \\ &= \sum_{l=0}^{+\infty} (l+1)^{4r} (\mathcal{J}_l^2(f) - \mathcal{J}_{l+1}^2(f)). \end{aligned}$$

Using an Abel transformation, we obtain

$$\|\Lambda_{q,\alpha}^r f\|_{q,2,\alpha}^2 \leq \mathcal{J}_0^2(f) + 4r \sum_{l=1}^{+\infty} (l+1)^{4r-1} \mathcal{J}_l^2(f).$$

From the inequality $l+1 \leq 2l$, we conclude

$$\|\Lambda_{q,\alpha}^r f\|_{q,2,\alpha}^2 \leq c_5 \left(\mathcal{J}_0^2(f) + \sum_{l=1}^{+\infty} l^{4r-1} \mathcal{J}_l^2(f) \right).$$

Hence,

$$\|\Lambda_{q,\alpha}^r f\|_{q,2,\alpha} = \mathcal{O} \left(\sum_{l=1}^{+\infty} l^{2r-1} \mathcal{J}_l(f) \right).$$

Since the series

$$\sum_{l=1}^{+\infty} l^{2r-1} \mathcal{J}_l(f), \quad r = 1, 2, \dots$$

converges, we see that $f \in W_{q,2}^r(\Lambda_{q,\alpha})$.

On the other hand, it follows from Lemma 3.2 that

$$\|\Delta_h^m(\Lambda_{q,\alpha}^r f)\|_{q,2,\alpha}^2 = \int_0^{+\infty} \lambda^{4r} |1 - j_\alpha(\lambda h, q^2)|^{2m} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda.$$

This integral is divided into two:

$$\int_0^{+\infty} = \int_0^N + \int_N^{+\infty} = K_1 + K_2,$$

where $N = [\frac{1}{h}]$. We estimate them separately.

By rearranging terms analogous to summation by parts and proceeding as with I_1 , we obtain

$$\begin{aligned} K_1 &\leq c_6 h^{4m} \int_0^N \lambda^{4(r+m)} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda \\ &= c_6 h^{4m} \sum_{l=0}^{N-1} \int_l^{l+1} \lambda^{4(r+m)} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda \\ &\leq c_6 h^{4m} \sum_{l=0}^{N-1} (l+1)^{4(r+m)} (\mathcal{J}_l^2(f) - \mathcal{J}_{l+1}^2(f)) \\ &\leq c_6 N^{-4m} \sum_{l=0}^{N-1} (l+1)^{4r+4m-1} \mathcal{J}_l^2(f). \end{aligned}$$

Now we estimate K_2 , by relation (2.6), we obtain

$$\begin{aligned} K_2 &= \int_N^{+\infty} \lambda^{4r} |1 - j_\alpha(\lambda h, q^2)|^{2m} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda \\ &= \mathcal{O} \left(\int_N^{+\infty} \lambda^{4r} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda \right) \\ &= \mathcal{O} \left(\sum_{m=1}^{+\infty} \int_{2^{m-1}N}^{2^m N} \lambda^{4r} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda \right) \\ &= \mathcal{O} \left(\sum_{m=1}^{+\infty} (2^m N)^{4r} \int_{2^{m-1}N}^{2^m N} |\mathcal{F}_{q,\alpha}(f)(\lambda)|^2 \lambda^{2\alpha+1} d_q \lambda \right) \\ &= \mathcal{O} \left(\sum_{m=1}^{+\infty} (2^m N)^{4r} \mathcal{J}_{2^{m-1}N}^2(f) \right), \end{aligned}$$

i.e.,

$$(K_2)^{\frac{1}{2}} = \mathcal{O} \left(\sum_{m=1}^{+\infty} (2^m N)^{2r} \mathcal{J}_{2^{m-1}N}^2(f) \right).$$

Taking account of the fact that

$$\begin{aligned} 2^{4r} \sum_{l=2^{m-2}N+1}^{2^{m-1}N} l^{2r-1} \mathcal{J}_l(f) &\geq 2^{4r} (2^{m-2}N)^{2r-1} \mathcal{J}_{2^{m-1}N}(f) 2^{m-2}N \\ &= (2^m N)^{2r} \mathcal{J}_{2^{m-1}N}(f), \end{aligned}$$

we obtain the estimate

$$(K_2)^{\frac{1}{2}} = \mathcal{O} \left(\sum_{m=1}^{+\infty} \sum_{l=2^{m-2}N+1}^{2^{m-1}N} l^{2r-1} \mathcal{J}_l(f) \right) = \mathcal{O} \left(\sum_{l=[\frac{N}{2}]}^{+\infty} l^{2r-1} \mathcal{J}_l(f) \right).$$

Combining the estimates for K_1 and K_2 gives

$$\|\Delta_h^m(\Lambda_{q,\alpha}^r f)\|_{q,2,\alpha} = \mathcal{O} \left(N^{-4m} \sum_{l=0}^{N-1} (l+1)^{4r+4m-1} \mathcal{J}_l^2(f) \right)^{\frac{1}{2}} + \mathcal{O} \left(\sum_{l=\lceil \frac{N}{2} \rceil}^{+\infty} l^{2r-1} \mathcal{J}_l(f) \right),$$

which implies that

$$\omega_m \left(\Lambda_{q,\alpha}^r f, \frac{1}{N} \right)_{q,2,\alpha} = \mathcal{O} \left(N^{-4m} \sum_{l=0}^{N-1} (l+1)^{4r+4m-1} \mathcal{J}_l^2(f) \right)^{\frac{1}{2}} + \mathcal{O} \left(\sum_{l=\lceil \frac{N}{2} \rceil}^{+\infty} l^{2r-1} \mathcal{J}_l(f) \right).$$

□

REFERENCES

- [1] V. A. Abilov, F. V. Abilova and M. K. Kerimov, *Some remarks concerning the Fourier transform in the space $L_2(\mathbb{R})$* , Comput. Math. Math. Phys. **48**(6) (2008), 885–891.
- [2] V. A. Abilov, F. V. Abilova and M. K. Kerimov, *On estimates for the Fourier-Bessel integral transform in the space $L^2(\mathbb{R}^+)$* , Zh. Vychisl. Mat. Mat. Fiz. **49**(7) (2009), 1158–1166.
- [3] V. A. Abilov, F. V. Abilova and M. K. Kerimov, *Some remarks concerning the Fourier transform in the space $L_2(\mathbb{R}^n)$* , Zh. Vychisl. Mat. Mat. Fiz. **48** (2008), 939–945. [Comput. Math. Math. Phys. **48** (2008), 885–891].
- [4] A. Achak, R. Daher, L. Dhaouadi and E. Loualid, *An analog of Titchmarsh's theorem for the q -Bessel transform*, Ann. Univ. Ferrara **65**(1) (2019). <https://doi.org/10.1007/s11565-018-0309-3>
- [5] N. Bettaibi, F. Bouzeffour, H. Ben Elmonser and W. Binous, *Elements of harmonic analysis related to the third basic zero order Bessel function*, J. Math. Anal. Appl. **342** (2008), 1203–1219.
- [6] R. Daher and M. El Hamma, *On estimates for the Bessel transform in the space $L_{p,\alpha}(\mathbb{R}^+)$* , Thai J. Math. **11**(3) (2013).
- [7] R. Daher and M. El Hamma, *On estimates for the Fourier transform in the space $L^2(\mathbb{R}^n)$* , C. R. Math. **352** (2014), 235–240. <https://doi.org/10.1016/j.crma.2013.12.016>
- [8] R. Daher and O. Tyr, *An analog of Titchmarsh's theorem for the q -Dunkl transform in the space $L_{q,\alpha}^2(\mathbb{R}_q)$* , J. Pseudo-Differ. Oper. Appl. (2020). <https://doi.org/10.1007/s11868-020-00330-6>
- [9] L. Dhaouadi, *On the q -Bessel Fourier transform*, Bull. Math. Anal. Appl. **5**(2) (2013), 42–60.
- [10] L. Dhaouadi, A. Fitouhi and J. El Kamel, *Inequalities in q -Fourier analysis*, J. Inequal. Pure Appl. Math. **7**(5) (2006), Article 171.
- [11] L. Dhaouadi, *Positivity of the generalized translation associated with the q -Hankel transform and applications*, Integr. Transf. Spec. Funct. **26**(2) (2015), 102–117.
- [12] S. El Ouadih, R. Daher and M. El Hamma, *Moduli of smoothness and K -functional in $L_2(\mathbb{R}_q^+)$ space with power weight*, J. Math. Anal. Appl. **45**(3) (2019), 491–503.
- [13] A. Fitouhi, M. Hamza and F. Bouzeffour, *The q - j_α -Bessel function*, J. Approx. Theory **115** (2002), 144–166.
- [14] F. H. Jackson, *On a q -definite integrals*, J. Pure Appl. Math. **41** (1910), 193–203.
- [15] M. Haddad, *Hankel transform in quantum calculus and applications*, Fract. Calc. Appl. Anal. **9** (2006), 371–384.
- [16] M. E. H. Ismail, *The zeros of basic Bessel functions, the function $J_{v+ax}(x)$, and associated orthogonal polynomials*, J. Math. Anal. Appl. **86** (1982), 1–19.
- [17] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, Encyclopedia of Mathematics and its Applications, Vol. 35, Cambridge Univ. Press, Cambridge, 1990.
- [18] V. G. Kac and P. Cheung, *Quantum Calculus*, Universitext, Springer-Verlag, New York, 2002.

- [19] H. T. Koelink and R. F. Swarttouw, *On the zeros of the Hahn-Exton q -Bessel function and associated q -Lommel polynomials*, J. Math. Anal. Appl. **186** (1994), 690–710.
- [20] T. H. Koornwinder and R. F. Swarttouw, *On q -analogues of the Hankel and Fourier transforms*, Trans. Amer. Math. Soc. **333** (1992), 445–461.
- [21] S. M. Nikol'skii, *Approximation of Functions of Several Variables and Embedding Theorems*, 2nd Edn., Nauka, Moscow, 1977; English transl. of 1st Ed., Grundlehren Math. Wiss., Vol. 205, Springer-Verlag, New York, Heidelberg, 1975.
- [22] S. B. Stechkin, *On the order of best approximation of continuous functions*, Izv. Akad. Nauk SSSR Ser. Mat. **15**(3) (1951), 219–242.
- [23] A. F. Timan and M. F. Timan, *Generalized modulus of continuity and best approximation in the mean*, Dokl. Akad. Nauk SSSR **71**(1) (1950), 17–20. (in Russian)
- [24] E. C. Titchmarsh, *Introduction to the Theory of Fourier Integrals*, Oxford University Press, Oxford, 1937.
- [25] O. Tyr and A. Dades, *Results concerning the Dunkl transform in the space $L^p(\mathbb{R}^d, w_\kappa(x) dx)$* , Asian-European J. Math. (2025), Article ID 2550076.
- [26] O. Tyr and R. Daher, *Abilov's estimates for the Clifford-Fourier transform in real Clifford algebras analysis*, Ann. Univ. Ferrara **69** (2023), 227–243.
- [27] O. Tyr and R. Daher, *Abilov's inequalities in the Laguerre hypergroup*, Filomat **39**(1) (2025), 313–324.
- [28] O. Tyr and R. Daher, *On the approximation of functions by Jacobi-Dunkl expansions in the weighted space $\mathbb{L}_2^{\alpha, \beta}$* , Ukr. Math. J. **74** (2023), 1629–1644.
- [29] O. Tyr, *On Abilov's inequalities in the sphere*, J. Math. Sci. (2025). <https://doi.org/10.1007/s10958-025-07656-w>
- [30] O. Tyr, R. Daher, S. E. Ouadiah and O. El Fourchi, *On the Jackson-type inequalities in approximation theory connected to the q -Dunkl operators in the weighted space* Bol. Soc. Mat. Mex. **27**(51) (2021).

¹LRST LABORATORY, THE HIGHER SCHOOL OF EDUCATION AND TRAINING OF AGADIR
 IBN ZOHR UNIVERSITY,
 AGADIR, MOROCCO
E-mail address: o.tyr@uiz.ac.ma
ORCID iD: <https://orcid.org/0000-0002-7261-8809>

²LABORATORY OF FUNDAMENTAL AND APPLIED MATHEMATICS,
 DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES AÏN CHOCK,
 HASSAN II UNIVERSITY,
 CASABLANCA, MOROCCO
E-mail address: dadesabdelali@gmail.com
ORCID iD: <https://orcid.org/0000-0003-1649-2364>