

APPROXIMATE SOLUTION OF BRATU DIFFERENTIAL EQUATIONS USING TRIGONOMETRIC BASIC FUNCTIONS

BAHRAM AGHELI

ABSTRACT. In this paper, I have proposed a method for finding an approximate function for Bratu differential equations (BDEs), in which trigonometric basic functions are used. First, by defining trigonometric basic functions, I define the values of the transformation function in relation to trigonometric basis functions (TBFs). Following that, the approximate function is defined as a linear combination of trigonometric base functions and values of transform function which is named trigonometric transform method (TTM), and the convergence of the method is also presented. To get an approximate solution function with discrete derivatives of the solution function, we have determined the approximate solution function which satisfies in the Bratu differential equations (BDEs). In the end, the algorithm of the method is elaborated with several examples. In one example, I have presented an absolute error comparison of some approximate methods.

1. INTRODUCTION

A problem of the non-linear eigenvalue problem in n dimensions is the Bratu differential equations (BDEs) as follows [13]

$$(1.1) \quad \sum_{i=1}^n \left(\frac{\partial}{\partial t_i} \right)^2 \Phi(t_1, t_2, \dots, t_n) + \lambda \exp(\Phi(t_1, t_2, \dots, t_n)) = 0,$$

in which $|x_i| \leq 1$ for $i = 1, 2, \dots, n$, with the following boundary conditions as $|x_i| = 1$,

$$(1.2) \quad \Phi(t_1, t_2, \dots, t_n) = 0.$$

The main objective in this paper is to offer a simple method in which it is possible to apply trigonometric transform method (TTM) to tackle with the one-dimensional

Key words and phrases. Trigonometric transform, Bratu differential equations, basic functions.
2010 Mathematics Subject Classification. Primary: 33D52. Secondary: 35A24, 65C30.
Received: May 29, 2018.
Accepted: November 06, 2018.

(1D) BDEs of the following form

$$(1.3) \quad u''(t) + \lambda \exp(u(t)) = 0, \quad 0 < t \leq T,$$

$$(1.4) \quad u(0) = u_0, \quad u_t(0) = u'_0,$$

where $\lambda > 0$ and $t \in \mathbb{R}$ are constant functions (see [12, 23]).

The analytic solution for BDEs is presented as follows:

$$u(t) = \log \left(\frac{\cosh \left(\frac{\phi}{2} \left(t - \frac{1}{2} \right) \right)}{\cosh \left(\frac{\phi}{4} \right)} \right)^{-2},$$

in which ϕ is the solution of $\phi = \sqrt{2\lambda} \cosh \left(\frac{\phi}{4} \right)$ (see [12, 23]). Whereas $\lambda_\epsilon = 3.513830719$, the BDEs has

- one solutions when $\lambda = \lambda_\epsilon$;
- two solutions if $\lambda < \lambda_\epsilon$;
- no solution when $\lambda > \lambda_\epsilon$.

Researchers and scholars are requested to check papers that have been introduced to get a better grasp of thoroughgoing introduction about BDEs and its history in [10, 18].

On the importance and motivation for Bratu differential equation, it should noted that it has a key role in many of the physical phenomena, chemical models and other sciences. Such applications include the model of thermal reaction process, the fuel ignition model of the thermal combustion theory, the Chandrasekhar model of the expansion of the universe, the radiative heat transfer nanotechnology and the chemical reaction theory (see [9, 10, 12, 18]).

As another instance, mathematical modeling in chemistry for the electro-spinning process is related to BDEs via thermo-electro-hydrodynamics balance equations. Colantoni and his co-author in [5] represented a model that is the mono-dimensional Bratu equation as follows:

$$(1.5) \quad u''(t) - \lambda \exp(u(t)) = 0,$$

featuring $\lambda = \frac{18 E^2 (I - r^2 k E)^2}{\rho^2 r^4}$, in which

- r is the radius of the jet at axial coordinate X in the Figure 1;
- I is the electrical current intensity;
- E is the electric area in the axial direction;
- ρ is the material density;
- k is a fixed value which is only dependent on temperature with regard to incompressible polymer.

Many researchers have used numerical methods for the purpose of solving the BDEs. We can refer to a number of familiar methods, including Homotopic perturbation method [8], Finite difference [19], Optimal homotopy asymptotic method [6], Wavelet method [17], Laplace transform decomposition method [15], B-splines method [4], Variational iteration technique [7], Adomian decomposition method [23], Differential




FIGURE 1. Electro-spinning process setup.

quadrature method [21], Lie-group shooting method [1], Reproducing kernel Hilbert space method [2], Pseudo-spectral collocation method [3] and [11, 12, 14, 16, 22].

This paper is organized as what follows: in Section 2, discretization of the derivative is given. In Section 3, we have expressed the trigonometric Basic functions (TBFs). In Section 4, a description of the new approach that is named trigonometric transform method (TTM) is presented. Some numerical examples are offered in Section 5. And conclusions are drawn in Section 6.

2. DISCRETIZATION OF THE DERIVATIVE

In this section, we introduce discretization of the derivative of a function. The approximation of derivatives by forward differences is one of the most basic tools in finite difference methods for the approximate solution of differential equations, especially initial value problems. The n -th order forward difference is given by

$$u^{(n)}(t) \approx \frac{1}{h^n} \sum_{i=0}^n (-1)^i \binom{n}{i} u((n-i)h + t), \quad n \in \mathbb{N}.$$

Depending on the application, the spacing h may be a variable or a fixed. In this paper, we consider $\tau = t_{j+1} - t_j$ and $t_j = a + j\tau$ for $j = 0, 1, 2, \dots$. For second order derivative we have:

$$(2.1) \quad u''(t_{k+1}) \approx \frac{1}{h^2} (u(t_{j+1}) - 2u(t_j) + u(t_{j-1})),$$

in which $u(t_0)$ and $u'(t_0)$ are known and $u(t_{-1}) = u(0) - \tau u_t(0)$.

3. TRIGONOMETRIC BASIC FUNCTIONS (TBFs)

In this section, we introduce the trigonometric basis functions and properties that are used in the main sections of the paper to approximate the function of the solution.

Definition 3.1. Presuming that for $n \geq 1$, $a = t_0 < t_1 < \dots < t_{n-1} < t_n = b$ be specified nodes, we express that basic functions T_0, T_1, \dots, T_n are defined on $[a, b]$

with their trigonometric functions $T_0(t), T_1(t), \dots, T_n(t)$, as follows:

$$(3.1) \quad \begin{aligned} T_0(t) &= \begin{cases} 0.5 \left(1 + \cos \frac{\pi}{h_0}(t - t_0)\right), & t_0 \leq t \leq t_1, \\ 0, & \text{otherwise,} \end{cases} \\ T_k(t) &= \begin{cases} 0.5 \left(1 + \cos \frac{\pi}{h_{k-1}}(t - t_k)\right), & t_{k-1} \leq t \leq t_k, \\ 0.5 \left(1 + \cos \frac{\pi}{h_k}(t - t_k)\right), & t_k \leq t \leq t_{k+1}, \\ 0, & \text{otherwise,} \end{cases} \quad k = 1, 2, 3, \dots, n-1, \\ T_n(t) &= \begin{cases} 0.5 \left(1 + \cos \frac{\pi}{h_{n-1}}(t - t_n)\right), & t_{n-1} \leq t \leq t_n, \\ 0, & \text{otherwise,} \end{cases} \end{aligned}$$

in which $h_k = t_{k+1} - t_k$ for $k = 0, 1, \dots, n-1$.

Remark 3.1. The trigonometric functions introduced in Definition 3.1 are the trigonometric basis functions (TBFs) in which the following properties are satisfied.

- (1) T_k of $[a, b]$ to $[0, 1]$ is continuous, $\sum_{k=0}^n T_k(t) = 1$ for all $t \in [a, b]$ and $T_k(t_k) = 1$, $k = 0, 1, 2, \dots, n$.
- (2) $T_k(t) = 0$ if $t \notin (t_{k-1}, t_{k+1})$, for $k = 1, 2, \dots, n-1$, $T_0(t) = 0$ if $t \notin (t_0, t_1)$ and $T_n(t) = 0$ if $t \notin (t_{n-1}, t_n)$.
- (3) On subinterval $[t_{k-1}, t_{k+1}]$ for $k = 1, 2, \dots, n-1$, $T_k(t)$, certainly is an increasing function on $[t_{k-1}, t_k]$ and decreasing function on $[t_k, t_{k+1}]$. Basic functions are called uniform as long as $t_{k+1} - t_k = h = \frac{b-a}{n}$ and two additional properties coincide.
- (4) $T_k(t_k - t) = T_k(t_k + t)$, for all $t \in [0, h]$ and $k = 1, 2, \dots, n-1$;
- (5) $T_k(t) = T_{k-1}(t - h)$ and $T_{k+1}(t) = T_k(t - h)$, for $k = 1, 2, \dots, n-1$ and $t \in [t_k, t_{k+1}]$.

Lemma 3.1 ([20]). Consider $n \geq 2$, T_0, T_1, \dots, T_n , be the TBFs which builds on $[a, b]$. Therefore,

$$(3.2) \quad \int_{t_0}^{t_1} T_1(t) dt = \int_{t_{n-1}}^{t_n} T_n(t) dt = \frac{h}{2}$$

and

$$(3.3) \quad \int_{t_{k-1}}^{t_{k+1}} T_k(t) dt = h,$$

for $k = 1, 2, \dots, n-1$, in which h is the distance between each of the two neighboring nodes.

Definition 3.2. Let f be a function belonging to $C[a, b]$ and T_i , $i = 0, 1, \dots, n$, be the TBFs which buildup on $[a, b]$. We define the F_k that is the transform of function f on $[a, b]$ with respect to basic functions T_k given by

$$(3.4) \quad F_k = \frac{\int_a^b f(t) T_k(t) dt}{\int_a^b T_k(t) dt}, \quad k = 0, 1, 2, \dots, n.$$

Definition 3.3. Let f be a function belonging to $C[a, b]$ and T_i , $i = 0, 1, \dots, n$, be the TBFs which buildup on $[a, b]$ and F_k be transform of function f on $[a, b]$ with respect to basic functions T_k . Then

$$f_n(t) = \sum_{k=0}^n F_k T_k(t)$$

is approximate of function f on $[a, b]$ with respect to TBFs.

Theorem 3.1 (Convergence). *Let f be a uniformly continuous function on $[a, b]$. Thus, for any $\epsilon > 0$, there exists n_ϵ such that for all $n \geq n_\epsilon$*

$$(3.5) \quad |f(t) - f_{n_\epsilon}(t)| < \epsilon.$$

Proof. f is a uniformly continuous function on $[a, b]$. Therefore,

$$(\forall \epsilon > 0) (\exists \delta = \delta(\epsilon)) (|x - t| < \delta \Rightarrow |f(x) - f(t)| < \epsilon \quad (0 < \delta < \epsilon)).$$

For all $\epsilon > 0$, we have

$$|f(t) - f_n(t)| = \left| \sum_{i=0}^n T_i(t)f(t) - \sum_{i=0}^n F_i T_i(t) \right| \leq \sum_{i=0}^n T_i(t) |f(t) - F_i| < \epsilon.$$

It is sufficient to show that $|f(t) - F_i| < \epsilon$. Let $x, t \in [x_{i-1}, x_{i+1}]$, $i = 1, 2, \dots, n-1$, so that we can evaluate

$$|f(x) - F_i| = \left| f(x) - \frac{\int_a^b f(t)T_i(t)dt}{\int_a^b T_i(t)dt} \right| \leq \frac{\int_{x_{i-1}}^{x_{i+1}} T_i(t) |f(x) - f(t)| dt}{\int_{x_{i-1}}^{x_{i+1}} T_i(t)dt} < \epsilon,$$

if and only if

$$\delta < 2h < \epsilon \quad \text{or} \quad h < \frac{\epsilon}{2}.$$

Regarding $h = \frac{b-a}{n}$, it is sufficient that $n_\epsilon > \frac{2(b-a)}{\epsilon}$. □

For description of fractional derivative, we have the following proposition.

Proposition 3.1. *With substituting $f_n(t) = \sum_{k=0}^n F_k T_k(t)$ in (2.1), we will have the next equation for $k = 0, 1, 2, \dots, n-1$:*

$$(3.6) \quad f_n''(t_{k+1}) \approx \frac{1}{h^2}(F_{j+1} - 2F_j + F_{j-1}).$$

4. DESCRIPTION OF THE NEW APPROACH

Let solution of (1.3) be continuous on $[0, b]$. To gain approximate solution of $u(x)$, we divide $[0, b]$ to n equal partition with step length τ :

$$(4.1) \quad t_0 = 0, \quad t_i = t_0 + i\tau, \quad i = 0, 1, \dots, n, \tau = \frac{b}{n}.$$

Considering the trigonometric functions with regard to Definition 3.1 on $[0, b]$ and Definition 3.3, we can gain approximate function $u(x)$ by $u_n(x) = \sum_{k=0}^n U_k T_k(t)$. It is

evident that for calculating $u_n(t)$, $t \in [0, b]$, we should calculate U_k , $k = 0, 1, 2, \dots, n$. In order to gain the approximate solution of the problem (1.3), $u_n(t)$ for points t_0, t_1, \dots, t_n must be satisfied in (1.3). Due to the boundary conditions (1.4), $u_n(t_0) := u(t_0) = u_0$ and for other points t_1, t_2, \dots, t_n , we have

$$(4.2) \quad u_n''(t_{k+1}) + \lambda \exp(u_n(t_{k+1})) = 0, \quad k = 0, 1, 2, \dots, n-1,$$

in which $m-1 < \rho \leq m$ and $m \in \mathbb{Z}^+$.

Using (3.6) and (4.2) converts to the following form for $k = 0, 1, 2, \dots, n-1$:

$$(4.3) \quad \frac{1}{h^2}(U_{k+1} - 2U_k + U_{k-1}) + \lambda \exp(U_n(t_{k+1})) = 0,$$

where $U_0 = u(0)$ and $U_{-1} = u(0) - u'(0)$ are known initial conditions.

Now, using the boundary condition, we can calculate U_1, U_2, \dots, U_n by the obtained recursive equation (4.3) and then gain the approximate solution $u(t) \approx u_n(t)$ for (1.3).

In order to gain approximation of BDEs, an algorithm by this method is offered in the subsequent algorithm.

Algorithm 1: An algorithm for approximation of BDEs

Step 1: Input n and b .

Step 2: Set $\tau \leftarrow \frac{b}{n}$.

Step 3: Locate $t_k \leftarrow k\tau$, $k = 0, 1, 2, \dots, n$.

Step 4: Choose TBFs $T_k(t)$ toward $k = 0, 1, 2, \dots, n$.

Step 5: Set recursive equations

$$\frac{1}{h^2}(U_{k+1} - 2U_k + U_{k-1}) + \lambda \exp(U_n(t_{k+1})) = 0,$$

where $U_0 = u(0)$ and $U_{-1} = u(0) - \tau u'(0)$.

Step 6: Calculate every U_k , $k = 1, 2, \dots, n$, of an equation of degree one.

Step 7: The approximate solution is

$$u_n(t) \approx \sum_{k=0}^n U_k T_k(t).$$

5. EXAMPLES

Now that it is easier to understand trigonometric transform, a number of examples will be given in this section and then will be calculated. These examples include BDEs. In all these examples, software *Mathematica* 11 has been used for calculations and graphs.

Example 5.1. We propose the BDEs for the first example [23]:

$$(5.1) \quad u''(t) - 2 \exp(u(t)) = 0, \quad 0 \leq t \leq 1,$$

with the precise solution $u(t) = \log((\cos t)^{-2})$ and the primary conditions:

$$(5.2) \quad u(0) = 0, \quad u'(0) = 0.$$

Following the TTM, according to what was formulated and presented in section 4 for (5.1)–(5.2), we can calculate U_1, U_2, \dots, U_n , and then gain the approximate solution $u_n(t)$ of (5.1).

In Table 1, we can see the estimated solutions for Eq.(5.1), which is derived for various values of n applying TTM. Also, the estimated and approximate solutions are illustrate in Figure 2.

TABLE 1. Approximate result of Example 5.1 with various values of n .

t	TTM				Exact
	$n = 50$	$n = 500$	$n = 1000$	$n = 1500$	
0.0	0.0	0.0	0.0	0.0	0.0
0.2	0.0543317	0.0407728	0.0404703	0.0402949	0.0402695
0.4	0.193714	0.165493	0.164871	0.164416	0.164458
0.6	0.42896	0.385508	0.384559	0.383323	0.38393
0.8	0.799043	0.725417	0.723832	0.722438	0.722781




FIGURE 2. Figure for Example 5.1 exact and the approximation solutions.

Noteworthy in the values obtained in the Table 1 is that by increasing the amount n , a more accurate answer for (5.1) can be achieved.

Example 5.2. Consider the BDEs for the second example [23]:

$$(5.3) \quad u''(t) + \pi^2 \exp(-u(t)) = 0, \quad 0 \leq t \leq 1,$$

given that the primary conditions:

$$(5.4) \quad u(0) = 0, \quad u'(0) = \pi.$$

The unknown coefficient U_1, U_2, \dots, U_n with due attention to the TTM, according to Section 4 for (5.3)–(5.4) are calculated.


In Table 2 and in Figure 3, we can view the precise and approximate answers for $n = 1500$ through applying TTM.

The approximate solution obtained by the proposed method corresponds to the precise solution $u(t) = \log(1 + \sin(\pi t))$.

In Figure 3, we can see the estimated solutions toward $n = 1500$, which is derived for various value of t applying TTM.

TABLE 2. Approximate result of example 5.2.

t	TTM	Exact	Absolute Error	Relative Error
0.2	0.462127	0.46234	212.789×10^{-6}	460.455×10^{-6}
0.4	0.66794	0.668371	430.849×10^{-6}	645.042×10^{-6}
0.6	0.667754	0.668371	616.549×10^{-6}	923.317×10^{-6}
0.8	0.46142	0.46234	920.306×10^{-6}	1.99451×10^{-3}

FIGURE 3. Comparison of the approximate solution (5.3) with exact solution for $n = 1500$.

Example 5.3. We offer the BDEs for the third example [23]:

$$(5.5) \quad u''(t) - \pi^2 \exp(u(t)) = 0, \quad 0 \leq t \leq 1,$$

including the primary conditions:

$$(5.6) \quad u(0) = 0, \quad u'(0) = -\pi.$$

It can be seen in Table 3 and Figure 4 that solving equations with approximate expression is calculated and displayed for $n = 1500$ and various values of t .

TABLE 3. Approximate result of Example 5.3.

t	TTM	Exact	Absolute Error	Relative Error
0.2	0.451242	0.451272	30.7122×10^{-6}	68.0615×10^{-6}
0.4	-0.227657	-0.226202	1.45505×10^{-3}	6.39141×10^{-3}
0.6	-0.576992	-0.573173	3.81849×10^{-3}	6.61792×10^{-3}
0.8	-0.699629	-0.69232	7.30951×10^{-3}	10.4477×10^{-3}

In Table 4, we can see the estimated solutions toward $n = 1500$, which is derived for various values of t applying TTM.

Toward $n = 1500$, the solution that we have gained is in accordance with the precise solution $u(t) = \log\left(\frac{1}{1 - \sin(1 - \pi t)}\right)$.

Example 5.4. Consider the BDEs [1]:

$$(5.7) \quad u''(t) + 2 \exp(u(t)) = 0, \quad 0 \leq t \leq 1,$$

supposing that the primary conditions:

$$(5.8) \quad u(0) = 0, \quad u'(0) = 0.$$

TABLE 4. Absolute error comparison of Example 5.4.

t	BNM	NPS	LTM	DM	BSM	LGSM	SCM	TTM	
								n=1500	n=1500
0.1	1.91×10^{-14}	9.71×10^{-9}	2.13×10^{-3}	1.52×10^{-2}	1.72×10^{-5}	4.03416×10^{-6}	6.88×10^{-4}	0	0
0.3	1.17×10^{-13}	1.98×10^{-8}	6.19×10^{-3}	5.89×10^{-3}	4.49×10^{-5}	5.22122×10^{-6}	8.21×10^{-4}	169.624×10^{-6}	169.624×10^{-6}
0.5	1.88×10^{-13}	2.60×10^{-8}	9.60×10^{-3}	6.98×10^{-3}	5.56×10^{-5}	1.4554×10^{-8}	8.60×10^{-4}	341.417×10^{-6}	341.417×10^{-6}
0.7	1.16×10^{-13}	1.98×10^{-8}	1.19×10^{-3}	5.89×10^{-3}	4.49×10^{-5}	5.19455×10^{-6}	8.21×10^{-4}	424.587×10^{-6}	424.587×10^{-6}
0.9	1.90×10^{-14}	9.71×10^{-9}	1.09×10^{-3}	1.52×10^{-3}	1.72×10^{-5}	4.01345×10^{-6}	6.88×10^{-4}	445.899×10^{-6}	445.899×10^{-6}




FIGURE 4. Comparison of the approximate solution (5.5) with exact solution for $n = 1500$.

The unknown coefficient U_1, U_2, \dots, U_n , with due attention to the TTM, according to Section 4 for (5.7)–(5.8) are calculated.

Table 4 illustrates an absolute error comparison of the TTM and approximate methods: Block Nyström method (BNM) [12], Non-polynomial spline (NPS) [11], Laplace transform method (LTM) [18], Decomposition method (DM) [16], B-splines method (BSM) [4], Lie-group shooting method (LGSM) [1] and Sinc-collocation method (SCM).

In Figure 5, we can see the estimated solutions toward $n = 1500$, which is derived for various value of t applying TTM.




FIGURE 5. Comparison of the approximate solution (5.7) with exact solution for $n = 1500$.

Noteworthy in the values obtained in the last column Table 4 is that by increasing the amount n , a more accurate answer for (5.7) can be achieved.

6. CONCLUSION

I have proposed a method for finding an approximate function of Bratu differential equations (BDEs), in which TTM are used. All examples with absolute and relative errors show that we have favorably applied trigonometric transform method TTM to obtain approximate solution of the BDEs. The obtained solutions that are very close analytical solutions indicate that a little iteration of TTM will result in some useful solutions. As the result seems necessary from the authors' point of view, the suggested technique has the potentials to be practical in solving other similar ordinary

differential equations of integer orders and partial differential equations of non integer orders.

REFERENCES

- [1] S. Abbasbandy, M. Hashemi and C.-S. Liu, *The lie-group shooting method for solving the Bratu equation*, Commun. Nonlinear Sci. Numer. Simul. **16** (2011), 4238–4249.
- [2] A. Akgül, F. Geng et al., *Reproducing kernel Hilbert space method for solving Bratu's problem*, Bull. Malays. Math. Sci. Soc. **38** (2015), 271–287.
- [3] J. P. Boyd, *One-point pseudospectral collocation for the one-dimensional Bratu equation*, Appl. Math. Comput. **217** (2011), 5553–5565.
- [4] H. Caglar, N. Caglar, M. Özer, A. Valaristos and A. N. Anagnostopoulos, *B-spline method for solving Bratu's problem*, Int. J. Comput. Math. **87** (2010), 1885–1891.
- [5] A. Colantoni and K. Boubaker, *Electro-spun organic nanofibers elaboration process investigations using comparative analytical solutions*, Carbohydrate Polymers **101** (2014), 307–312.
- [6] M. A. Darwish and B. S. Kashkari, *Numerical solutions of second order initial value problems of Bratu-type via optimal homotopy asymptotic method*, American Journal of Computational Mathematics **4** (2014), 47–54.
- [7] N. Das, R. Singh, A.-M. Wazwaz and J. Kumar, *An algorithm based on the variational iteration technique for the Bratu-type and the lane-Emden problems*, J. Math. Chem. **54** (2016), 527–551.
- [8] X. Feng, Y. He and J. Meng, *Application of homotopy perturbation method to the Bratu-type equations*, Topol. Methods Nonlinear Anal. **31** (2008), 243–252.
- [9] D. A. Frank-Kamenetskii, *Diffusion and Heat Exchange in Chemical Kinetics*, Princeton University Press, Princeton, New Jersey, 2015.
- [10] J. Jacobsen and K. Schmitt, *The Liouville-Bratu-Gelfand problem for radial operators*, J. Differential Equations **184** (2002), 283–298.
- [11] R. Jalilian, *Non-polynomial spline method for solving Bratu's problem*, Comput. Phys. Commun. **181** (2010), 1868–1872.
- [12] S. Jator and V. Manathunga, *Block Nyström type integrator for Bratu's equation*, J. Comput. Appl. Math. **327** (2018), 341–349.
- [13] J. Karkowski, *Numerical experiments with the Bratu equation in one, two and three dimensions*, Comput. Appl. Math. **32** (2013), 231–244.
- [14] E. Keshavarz, Y. Ordokhani and M. Razzaghi, *The Taylor wavelets method for solving the initial and boundary value problems of Bratu-type equations*, Appl. Numer. Math. **128** (2018), 205–216.
- [15] S. A. Khuri, *A new approach to Bratu's problem*, Appl. Math. Comput. **147** (2004), 131–136.
- [16] S. Liao and Y. Tan, *A general approach to obtain series solutions of nonlinear differential equations*, Stud. Appl. Math. **119** (2007), 297–354.
- [17] X. Liu, Y. Zhou, X. Wang and J. Wang, *A wavelet method for solving a class of nonlinear boundary value problems*, Commun. Nonlinear Sci. Numer. Simul. **18** (2013), 1939–1948.
- [18] J. S. McGough, *Numerical continuation and the Gelfand problem*, Appl. Math. Comput. **89** (1998), 225–239.
- [19] A. Mohsen, *A simple solution of the Bratu problem*, Comput. Math. Appl. **67** (2014), 26–33.
- [20] I. Perfilieva, *Fuzzy transforms: theory and applications*, Fuzzy Sets and Systems **157** (2006), 993–1023.
- [21] O. Ragb, L. Seddek and M. Matbulu, *Iterative differential quadrature solutions for Bratu problem*, Comput. Math. Appl. **74** (2017), 249–257.
- [22] H. Temimi and M. Ben-Romdhane, *An iterative finite difference method for solving Bratu's problem*, J. Comput. Appl. Math. **292** (2016), 76–82.
- [23] A.-M. Wazwaz, *Adomian decomposition method for a reliable treatment of the Bratu-type equations*, Appl. Math. Comput. **166** (2005), 652–663.

DEPARTMENT OF MATHEMATICS,
QAEMSHAHR BRANCH,
ISLAMIC AZAD UNIVERSITY,
QAEMSHAHR, IRAN
Email address: `b.agheli@qaemiau.ac.ir`