

THE MINIMUM EDGE COVERING ENERGY OF A GRAPH

SAMIRA SABETI¹, AKRAM BANIHASHEMI DEHKORDI¹,
AND SAEED MOHAMMADIAN SEMNANI¹

ABSTRACT. In this paper, we introduce a new kind of graph energy, the minimum edge covering energy, $E_{C_E}(G)$. It depends both on the underlying graph G , and on its particular minimum edge covering C_E . Upper and lower bounds for $E_{C_E}(G)$ are established. The minimum edge covering energy and some of the coefficients of the polynomial of well-known families of graphs like Star, Path and Cycle Graphs are computed.

1. INTRODUCTION

In the study of spectral graph theory, we use the spectra of the certain matrix associated with the graph, such as the adjacency matrix, the Laplacian matrix, and other related matrices. Some useful information about the graph can be obtained from the spectra of these various matrices. Let G be a simple graph and let its vertex set be $V(G) = \{v_1, v_2, \dots, v_n\}$. The adjacency matrix $A(G)$ of the graph G is a square matrix of order n whose (i, j) -entry is equal to unity if the vertices v_i and v_j are adjacent and it is equal to zero otherwise. The eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ of $A(G)$, assumed in nonincreasing order, are the eigenvalues of the graph G . The concept of energy of a graph was introduced by I. Gutman [1] in the year 1978 as the sum of the absolute values of its eigenvalues:

$$E(G) = \sum_{i=1}^n |\lambda_i|.$$

One of the remarkable chemical applications of spectral graph theory is based on the close correspondence between the graph eigenvalues and the molecular orbital energy

Key words and phrases. Minimum edge covering set, minimum edge covering matrix, graph energy, minimum edge covering eigenvalues.

2018 *Mathematics Subject Classification.* Primary: 05C50.

Received: November 19, 2018.

Accepted: July 15, 2019.

levels of π -electron in conjugated hydrocarbons. An interesting quantity in Huckel theory is the sum of the energies of all the electrons in a molecule, so-called total π -electron energy E . For more details on the mathematical aspects of the theory of graph energy see [2–5]. The basic properties including various upper and lower bounds for energy of a graph have been established [6,7], and it has found remarkable chemical applications in the molecular orbital theory of conjugated molecules [8]. Recently C. Adiga et al. [8] defined the minimum covering energy, $E_C(G)$ of a graph which depends on its particular minimum cover C . Further, incidence energy, matching energy, minimum dominating energy, Laplacian minimum dominating energy and minimum dominating distance energy of a graph G can be found in [9–16]. Motivated by these papers, we study the minimum edge covering E_{C_E} of a graph G . We compute some properties of the characteristic polynomial of a minimum edge covering matrix of a graph G . Upper and lower bounds for $E_{C_E}(G)$ are established. It is possible that the minimum edge covering energy that we are considering in this paper may have some applications in chemistry as well as in other areas. Let $G = (V, E)$ be a simple finite graph that is has no loops no multiple and directed edges. Graph G has n vertices, m edges with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$ and edge set $E(G) = \{e_1, e_2, \dots, e_m\}$.

Definition 1.1. A subset C of V is called a covering set of G if every edge of G is incident to at least one vertex of C . Any covering set with minimum cardinality is called a minimum covering set.

Definition 1.2. A subset of E is called an edge covering set of G if every vertex of G is incident to at least one edge of it. Any edge covering set with minimum cardinality is called a minimum edge covering set. Let C_E be a minimum edge covering set of a graph G . The minimum edge covering matrix of G is a $m \times m$ matrix $A_{C_E}(G) = (e_{ij})_{m \times m}$, where

$$e_{ij} = \begin{cases} 1, & \text{if } e_i, e_j \text{ are adjacent,} \\ 1, & \text{if } i = j \text{ and } e_i \in C_E, \\ 0, & \text{otherwise.} \end{cases}$$

The characteristic polynomial of $A_{C_E}(G)$ is denoted by $f_m(G, \lambda) = \det(\lambda I - A_{C_E}(G)) = c_0\lambda^m + c_1\lambda^{m-1} + c_2\lambda^{m-2} + \dots + c_m$. The matrix $A_{C_E}(G)$ is real and symmetric. Then the eigenvalues of $A_{C_E}(G)$ are real numbers and are labeled in non-increasing order $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_m$. The minimum edge covering energy of G is defined as

$$E_{C_E}(G) = \sum_{i=1}^m |\lambda_i|.$$

2. PROBLEM FORMULATION AND SOME BASIC PROPERTIES OF MINIMUM EDGE COVERING ENERGY

Following theorem obtains the coefficients of polynomial without applying determinant expansion.

Theorem 2.1. *Let G be a graph with vertex set V , edge set E , and the minimum edge covering set C_E . Let $f_m(G, \lambda) = \det(\lambda I - A_{C_E}(G)) = c_0\lambda^m + c_1\lambda^{m-1} + c_2\lambda^{m-2} + \dots + c_m$ be the characteristic polynomial of G . Then*

- (i) $c_0 = 1$;
- (ii) $c_1 = -|C_E|$;
- (iii) $c_2 = \binom{|C_E|}{2} - \left(\frac{1}{2} \sum_{i=1}^m \deg(e_i)\right)$;
- (iv) $c_3 = -\binom{|C_E|}{3} + |C_E| \left(\frac{1}{2} \sum_{i=1}^m \deg(e_i)\right) - \sum_{i \in C_E} \deg(e_i) - 2(K_{1,3} + C_3)$, where C_3 is the number of triangles and $K_{1,3}$ is the number of star graphs with four vertices in G .

Proof. (i) Directly from the definition of $f_m(G, \lambda)$, it follows that $c_0 = 1$.

(ii) Since the sum of diagonal elements of $A_{C_E}(G)$ is equal to $|C_E|$, sum of determinants of all 1×1 principal submatrices of $A_{C_E}(G)$ is the trace of $A_{C_E}(G)$ which evidently equal to $|C_E|$. Thus, $(-1)^1 c_1 = |C_E|$.

(iii) $(-1)^2 c_2$ is equal to sum of determinats of all the 2×2 principal submatrices of $A_{C_E}(G)$, that is

$$\begin{aligned} c_2 &= \sum_{1 \leq i < j \leq m} \begin{vmatrix} e_{ii} & e_{ij} \\ e_{ji} & e_{jj} \end{vmatrix} = \sum_{1 \leq i < j \leq m} (e_{ii}e_{jj} - e_{ij}e_{ji}) \\ &= \sum_{1 \leq i < j \leq m} e_{ii}e_{jj} - \sum_{1 \leq i < j \leq m} e_{ij}^2 = \binom{|C_E|}{2} - \frac{1}{2} \sum_{i=1}^m \deg(e_i). \end{aligned}$$

(iv)

$$\begin{aligned} c_3 &= (-1)^3 \sum_{1 \leq i < j < k \leq m} \begin{vmatrix} e_{ii} & e_{ij} & e_{ik} \\ e_{ji} & e_{jj} & e_{jk} \\ e_{ki} & e_{kj} & e_{kk} \end{vmatrix} \\ &= - \sum_{1 \leq i < j < k \leq m} [e_{ii}(e_{jj}e_{kk} - e_{kj}e_{jk}) - e_{ij}(e_{ji}e_{kk} - e_{ki}e_{jk}) \\ &\quad + e_{ik}(e_{ji}e_{kj} - e_{ki}e_{jj})] \\ &= - \sum_{1 \leq i < j < k \leq m} e_{ii}e_{jj}e_{kk} + \sum_{1 \leq i < j < k \leq m} [e_{ii}e_{jk} + e_{jj}e_{ik} + e_{kk}e_{ij}] \\ &\quad - \sum_{1 \leq i < j < k \leq m} e_{ij}e_{jk}e_{ki} - \sum_{1 \leq i < j < k \leq m} e_{ik}e_{kj}e_{ji} \\ &= \binom{|C_E|}{3} + |C_E| \left(\frac{1}{2} \sum_{i=1}^m \deg(e_i)\right) - \sum_{i \in C_E} \deg(e_i) - 2(k_{1,3} + C_3). \end{aligned}$$

Thus,

$$c_3 = \binom{|C_E|}{3} + |C_E| \left(\frac{1}{2} \sum_{i=1}^m \deg(e_i)\right) - \sum_{i \in C_E} \deg(e_i) - 2(k_{1,3} + C_3). \quad \square$$

Corollary 2.1. *Let G be a path P_n with n vertices and m edges, then*

- (i) $c_0 = 1$;
- (ii) $c_1 = -\left\lceil \frac{m+1}{2} \right\rceil$, $\lceil x \rceil$ is the smallest integer number greater than or equals to x ;
- (iii) $c_2 = \binom{\left\lceil \frac{m+1}{2} \right\rceil}{2} - (m - 1)$;
- (iv) $c_3 = \binom{\left\lceil \frac{m+1}{2} \right\rceil}{3} + \left\lceil \frac{m+1}{2} \right\rceil (m - 1) - \sum_{i \in C_E} \deg(e_i)$.

Corollary 2.2. *Let G be a cycle C_n with n vertices and m edges, then*

- (i) $c_0 = 1$;
- (ii) $c_1 = -\left\lceil \frac{m}{2} \right\rceil$;
- (iii) $c_2 = \binom{\left\lceil \frac{m}{2} \right\rceil}{2} - m$;
- (iv) $c_3 = -\binom{\left\lceil \frac{m}{2} \right\rceil}{3} + \left\lceil \frac{m}{2} \right\rceil (m - 2)$, $m \geq 4$.

Remark 2.1. Let G be a path P_n with n vertices and m edges, then

$$\sum_{i=1}^m \deg(e_i) = 2(m - 1).$$

Theorem 2.2. *If $\lambda_1, \lambda_2, \dots, \lambda_m$ are eigenvalues of $A_{C_E}(G)$, then*

$$\sum_{i=1}^m \lambda_i^2 = |C_E| + \sum_{i=1}^m \deg(e_i).$$

Proof. The sum of squares of the eigenvalues of $A_{C_E}(G)$ is just the trace of $(A_{C_E}(G))^2$. Therefore,

$$\sum_{i=1}^m \lambda_i^2 = \sum_{i=1}^m \sum_{j=1}^m e_{ij}e_{ji} = 2 \sum_{i < j} (e_{ij})^2 + \sum_{i=1}^m (e_{ii})^2 = |C_E| + \sum_{i=1}^m \deg(e_i). \quad \square$$

Theorem 2.3 (Parity theorem). *Let G be a graph with a minimum edge covering set C_E . If the minimum edge covering energy $E_{C_E}(G)$ of G is a rational number, then*

$$(2.1) \quad E_{C_E} \equiv |C_E| \pmod{2}.$$

Proof. Let $\lambda_1, \lambda_2, \dots, \lambda_r$ be positive, and the rest of the minimum edge covering eigenvalues non-positive. Thus,

$$E_{C_E}(G) = \sum_{i=1}^m |\lambda_i| = (\lambda_1 + \lambda_2 + \dots + \lambda_r) - (\lambda_{r+1} + \lambda_{r+2} + \dots + \lambda_m),$$

implies $E_{C_E}(G) = 2(\lambda_1 + \lambda_2 + \dots + \lambda_r) - |C_E|$. Since $\lambda_1, \lambda_2, \dots, \lambda_r$ are algebraic integers, so is their sum. Hence, $\lambda_1 + \lambda_2 + \dots + \lambda_r$ must be an integer if $E_{C_E}(G)$ is rational. \square

3. THE MINIMUM EDGE COVERING ENERGY OF SOME GRAPHS

Theorem 3.1. *The minimum edge covering energy of a star graph $K_{1,n-1}$ is m for $n \geq 2$. The polynomial of it is $\lambda^m - m\lambda^{m-1}$.*

Proof. Let $K_{1,n-1}$ be a star graph with vertex set $V = \{v_0, v_1, v_2, \dots, v_{n-1}\}$ and center v_0 , then

$$A_{C_E}(G) = \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 \\ 1 & 1 & 1 & \cdots & 1 \\ 1 & 1 & 1 & \cdots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & 1 & \cdots & 1 \end{pmatrix},$$

that $A_{C_E}(G)$ is $m \times m$. Then its characteristic polynomial is

$$(3.1) \quad f_m(k_{1,n-1}, \lambda) = \det \begin{pmatrix} \lambda - 1 & -1 & \cdots & -1 \\ -1 & \lambda - 1 & \cdots & -1 \\ \vdots & \vdots & \ddots & \vdots \\ -1 & -1 & \cdots & \lambda - 1 \end{pmatrix}.$$

By computing determinant of upper triangular of matrix (3.1) we will have

$$f_m(k_{1,n-1}, \lambda) = \lambda^m - m\lambda^{m-1}.$$

Then

$$\text{Spec}(k_{1,n-1}) = \begin{pmatrix} m & 0 \\ 1 & m - 1 \end{pmatrix},$$

$$E_{C_E}(G) = \sum_{i=1}^m |\lambda_i| = m. \quad \square$$

Theorem 3.2. *Let G be a path or a cycle graph with a minimum edge covering set C_E . Then $E_{C_E}(G) \simeq 2m - |C_E|$ or $|E_{C_E}(G) - (2m - |C_E|)| \leq 1$.*

Proof. Let $\lambda_1, \lambda_2, \dots, \lambda_r$ be positive eigenvalues, and the rest of the minimum edge covering eigenvalues non-positive. Thus,

$$E_{C_E}(G) = \sum_{i=1}^m |\lambda_i| = (\lambda_1 + \lambda_2 + \cdots + \lambda_r) - (\lambda_{r+1} + \lambda_{r+2} + \cdots + \lambda_m),$$

implies $E_{C_E}(G) = 2(\lambda_1 + \lambda_2 + \cdots + \lambda_r) - |C_E|$. Since $\lambda_1, \lambda_2, \dots, \lambda_r$ are algebraic integers, so sum $(\lambda_1 + \lambda_2 + \cdots + \lambda_r) \simeq m$. Hence, $E_{C_E}(G) \simeq 2m - |C_E|$. \square

Theorem 3.3 (Upper bound). *Let G be a graph with n vertices, m edges, and let C_E be a minimum edge covering set of G . Then*

$$E_{C_E}(G) \leq \sqrt{m \left(\sum_{i=1}^m \deg(e_i) + |C_E| \right)}.$$

Proof. Let $\lambda_1 \geq \lambda_2 \geq \lambda_3 \cdots \geq \lambda_m$ be the eigenvalues of $A_{C_E}(G)$. Bearing in mind the Cauchy-Schwarz inequality

$$\left(\sum_{i=1}^m a_i b_j\right)^2 \leq \left(\sum_{i=1}^m a_i^2\right) \left(\sum_{i=1}^m b_i^2\right),$$

we put $a_i = 1$ and $b_i = |\lambda_i|$ and we have

$$[E_{C_E}(G)]^2 = \left(\sum_{i=1}^m |\lambda_i|\right)^2 \leq m \left(\sum_{i=1}^m |\lambda_i|^2\right) = m \sum_{i=1}^m \lambda_i^2 = m \left(\sum_{i=1}^m \deg(e_i) + |C_E|\right). \square$$

Theorem 3.4 (Lower bound). *Let G be a graph with n vertices and m edges, and let C_E be a minimum edge covering set of G . If $D = |\det A_{C_E}(G)|$, then*

$$(3.2) \quad [E_{C_E}(G)]^2 \geq \sum_{i=1}^m \deg(e_i) + |C_E| + m(m-1) \sqrt[m]{\left(\prod_{i=1}^m \lambda_i\right)^2},$$

with equality if G is a star of order n .

Proof. We have

$$[E_{C_E}(G)]^2 = \left(\sum_{i=1}^m |\lambda_i|\right)^2 = \left(\sum_{i=1}^m |\lambda_i|\right) \left(\sum_{i=1}^m |\lambda_i|\right) = \sum_{i=1}^m |\lambda_i|^2 + \sum_{i \neq j} |\lambda_i| |\lambda_j|.$$

Now, by inequality between the arithmetic mean and geometric mean, we have

$$\frac{1}{m(m-1)} \sum_{i \neq j} |\lambda_i| |\lambda_j| \geq \left(\prod_{i \neq j} |\lambda_i| |\lambda_j|\right)^{\frac{1}{m(m-1)}}.$$

Thus,

$$\begin{aligned} [E_{C_E}(G)]^2 &\geq \sum_{i=1}^m |\lambda_i|^2 + m(m-1) \left(\prod_{i \neq j} |\lambda_i| |\lambda_j|\right)^{\frac{1}{m(m-1)}} \\ &\geq \sum_{i=1}^m |\lambda_i|^2 + m(m-1) \left(\prod_{i=1}^m |\lambda_i|^{2(m-1)}\right)^{\frac{1}{m(m-1)}} \\ &= \sum_{i=1}^m |\lambda_i|^2 + m(m-1) \left|\prod_{i=1}^m \lambda_i\right|^{\frac{2}{m}} \\ &= \sum_{i=1}^m \deg(e_i) + |C_E| + m(m-1) D^{\frac{2}{m}}. \end{aligned}$$

Since in the star graphs the multiplicity of $\lambda = 0$ is $m-1$, so $\prod_{i=1}^m \lambda_i = \det A_{C_E}(G) = 0$, $|C_E| = m$ and $\sum_{i=1}^m \deg(e_i) = m(m-1)$. Then, by placing the above values in (3.2), equality cases hold

$$m^2 = [E_{C_E}(K_{1,n-1})]^2 \geq m(m-1) + m + m(m-1) \sqrt[m]{\left(\prod_{i=1}^m \lambda_i\right)^2} = m^2. \quad \square$$

ACKNOWLEDGEMENTS

This research was supported by Research Council of Semnan University. The authors are grateful to Dr. Tatjana Aleksic Lampert and the anonymous reviewer for his/her valuable comments and suggestions helped to improve the quality of this work.

REFERENCES

- [1] I. Gutman, *The energy of a graph*, Ber. Math-Statist. Sect. Forschungsz. Graz **103** (1978), 1–22.
- [2] I. Gutman, B. Furtula, E. Zogić and E. Glogić, *Resolvent energy of graphs*, MATCH Commun. Math. Comput. Chem. **75** (2016), 279–290.
- [3] R. B. Bapat, *Graphs and Matrices*, Hindustan Book Agency, Springer, London, 2011.
- [4] S. K. Vaidya and K. M. Popat, *Some new results on energy of graphs*, MATCH Commun. Math. Comput. Chem. **77** (2017), 589–594.
- [5] V. Nikiforov, *The energy of graphs and matrices*, J. Math. Anal. Appl. **326** (2007), 1472–1475.
- [6] A. Jahanbani, *Lower bounds for the energy of graphs*, AKCE Int. J. Graphs Comb. **5** (2018), 88–96.
- [7] K. C. Das and I. Gutman, *Bounds for the energy of graphs*, Hacet. J. Math. Stat. **45**(3) (2016), 695–703.
- [8] C. Adiga, A. Bayad, I. Gutman and S. A. Srinivas, *The minimum covering energy of graph*, Kragujevac Journal of Science **34** (2012), 39–56.
- [9] I. Gutman and S. Wagner, *The matching energy of a graph*, Applied Mathematics **160** (2012), 2177–2187.
- [10] J. Zhang, H. Kan and X. Liu, *Graphs with extremal incident energy*, Filomat **29**(6) (2015), 1251–1258.
- [11] M. Jooyandeh, D. Kiani and M. Mirzakhah, *Incidence energy of a graph*, MATCH Commun. Math. Comput. Chem. **62** (2009), 561–572.
- [12] I. Gutman, D. Kiani and M. Mirzakhah, *On incidence energy of a graph*, MATCH Commun. Math. Comput. Chem. **62** (2009), 573–580.
- [13] R. Kanna, B. N. Dharmendra and G. Sridhara, *Laplacian minimum dominating energy of a graph*, International Journal of Pure and Applied Mathematics **89**(4) (2013), 565–581.
- [14] R. Kanna, B. N. Dharmendra and G. Sridhara, *Minimum dominating distance energy of a graph*, J. Indian Math. Soc. (N.S.) **20** (2014), 19–29.
- [15] R. Kanna, B. N. Dharmendra and G. Sridhara, *Minimum dominating energy of a graph*, International Journal of Pure and Applied Mathematics **85**(4) (2013), 707–718.
- [16] S. K. Vaidya and R. M. Pandit, *Edge domination in some path and cycle related graphs*, Hindawi Publishing Corporation, ISRN Discrete Mathematics (2014), Article ID 975812, 5 pages.

¹DEPARTMENT OF MATHEMATICS, STATISTICS AND COMPUTER SCIENCE,
SEM NAN UNIVERSITY,

ADDRESS SEM NAN UNIVERSITY, SEM NAN, IRAN

Email address: sabeti.samira@semnan.ac.ir

Email address: banihashemi.akram@semnan.ac.ir

Email address: s_mohammadian@semnan.ac.ir