

CONVERGENCE OF DOUBLE COSINE SERIES

KARANVIR SINGH¹ AND KANAK MODI²

ABSTRACT. In this paper we consider double cosine series whose coefficients form a null sequence of bounded variation of order $(p, 0)$, $(0, p)$ and (p, p) with the weight $(jk)^{p-1}$ for some $p > 1$. We study pointwise convergence, uniform convergence and convergence in L^r -norm of the series under consideration. In a certain sense our results extend the results of Young [7], Kolmogorov [3] and Móricz [4, 5].

1. INTRODUCTION

Consider the double cosine series

$$(1.1) \quad \sum_{j=0}^{\infty} \sum_{k=0}^{\infty} \lambda_j \lambda_k a_{jk} \cos jx \cos ky,$$

on positive quadrant $T = [0, \pi] \times [0, \pi]$ of the two dimensional torus where $\lambda_0 = \frac{1}{2}$ and $\lambda_j = 1$ for $j = 1, 2, 3, \dots$

The rectangular partial sums $S_{mn}(x, y)$ and the *Cesàro* means $\sigma_{mn}(x, y)$ of the series (1.1) are defined as

$$S_{mn}(x, y) = \sum_{j=0}^m \sum_{k=0}^n \lambda_j \lambda_k a_{jk} \cos jx \cos ky,$$
$$\sigma_{mn}(x, y) = \frac{1}{(m+1)(n+1)} \sum_{j=0}^m \sum_{k=0}^n S_{jk}(x, y), \quad m, n > 0,$$

Key words and phrases. Rectangular partial sums, L^r -convergence, *Cesàro* means, monotone sequences.

2010 *Mathematics Subject Classification.* Primary: 42A20, 42A32.

Received: September 17, 2017.

Accepted: June 05, 2018.

and for $\lambda > 1$, the truncated Cesàro means are defined by

$$V_{mn}^\lambda(x, y) = \frac{1}{([\lambda m] - m)([\lambda n] - n)} \sum_{j=m+1}^{[\lambda m]} \sum_{k=n+1}^{[\lambda n]} S_{jk}(x, y).$$

Now assuming the coefficients $\{a_{jk} : j, k \geq 0\}$ in (1.1) be a double sequence of real numbers which satisfy the following conditions for some positive integer p :

$$(1.2) \quad |a_{jk}|(jk)^{p-1} \rightarrow 0 \text{ as } \max\{j, k\} \rightarrow \infty,$$

$$(1.3) \quad \lim_{k \rightarrow \infty} \sum_{j=0}^{\infty} |\Delta_{p0} a_{jk}|(jk)^{p-1} = 0,$$

$$(1.4) \quad \lim_{j \rightarrow \infty} \sum_{k=0}^{\infty} |\Delta_{0p} a_{jk}|(jk)^{p-1} = 0,$$

$$(1.5) \quad \sum_{j=0}^{\infty} \sum_{k=0}^{\infty} |\Delta_{pp} a_{jk}|(jk)^{p-1} < \infty.$$

The finite order differences $\Delta_{pq} a_{jk}$ are defined by

$$\begin{aligned} \Delta_{00} a_{jk} &= a_{jk}, \\ \Delta_{pq} a_{jk} &= \Delta_{p-1,q} a_{jk} - \Delta_{p-1,q} a_{j+1,k}, \quad p \geq 1, q \geq 0, \\ \Delta_{pq} a_{jk} &= \Delta_{p,q-1} a_{jk} - \Delta_{p,q-1} a_{j,k+1}, \quad p \geq 0, q \geq 1. \end{aligned}$$

Also a double induction argument gives

$$\Delta_{pq} a_{jk} = \sum_{s=0}^p \sum_{t=0}^q (-1)^{s+t} \binom{p}{s} \binom{q}{t} a_{j+s, k+t}.$$

We can call the above mentioned conditions (1.2)-(1.5) as conditions of bounded variation of order $(p, 0)$, $(0, p)$ and (p, p) respectively with the weight $(jk)^{p-1}$. Obviously these conditions generalise the concept of monotone sequences. Also any sequence satisfying (1.5) with $p = 2$ is called a quasi-convex sequence [3, 5]. Clearly the conditions (1.3) and (1.4) can be derived from (1.2) and (1.5) for $p = 1$ and moreover for $p = 1$, the conditions (1.2) and (1.5) reduce to $|a_{jk}| \rightarrow 0$ as $\max\{j, k\} \rightarrow \infty$ and

$$\sum_{j=0}^{\infty} \sum_{k=0}^{\infty} |\Delta_{11} a_{jk}| < \infty.$$

Generally the pointwise convergence of the series (1.1) is defined in Pringsheim's sense ([8], Vol. 2, Ch. 17) which means that the rectangular partial sums of the type

$$S_{mn}(x, y) = \sum_{j=0}^m \sum_{k=0}^n \lambda_j \lambda_k a_{jk} \cos jx \cos ky, \quad m, n \geq 0,$$

are formed and then by taking both m, n tend to ∞ (independently of one another) the limit $f(x, y)$ (provided it exists) is assigned to the series (1.1) as its sum.

Also let $\|f\|_r$ denotes the $L^r(T^2)$ -norm, i.e,

$$\|f\|_r = \left(\int_0^\pi \int_0^\pi |f(x, y)|^r dx dy \right)^{1/r}, \quad 1 \leq r < \infty$$

and $\|f\|$ denotes $L^1(T^2)$ -norm, i.e,

$$\|f\| = \int_0^\pi \int_0^\pi |f(x, y)| dx dy.$$

In this paper, we will investigate the validity of the following statements:

- (a) $S_{mn}(x, y)$ converges pointwise to $f(x, y)$ for every $(x, y) \in T^2$;
- (b) $S_{mn}(x, y)$ converges uniformly to $f(x, y)$ on T^2 ;
- (c) $\|S_{mn}(x, y) - f(x, y)\|_r = o(1)$ as $\min\{m, n\} \rightarrow 0$.

Such type of problems have been studied by Young [7] and Kolmogorov [3] for one-dimensional case (single trigonometric series especially cosine series) and by Móricz [4, 5] and K. Kaur, Bhatia and Ram [2] for double trigonometric series. In [5], Móricz studied both double cosine series and double sine series as far as their integrability and convergence in L^1 -norm is concerned where as in [4] he studied double trigonometric series of the form

$$\sum_{-\infty}^{\infty} \sum_{-\infty}^{\infty} c_{jk} e^{i(jx+ky)},$$

under coefficients of bounded variation. All of them discussed the case for $p = 1$ or $p = 2$ only. Our aim in this paper is to extend the above results from $p = 1$ to general cases for double cosine series.

In the results, C_p and C_{pr} denote constants which may not be the same at each occurrence. Also we write $\lambda_n = [\lambda n]$ where n is a positive integer, $\lambda > 1$ is a real number and $[\cdot]$ means greatest integral part.

The first main result reads as follows.

Theorem 1.1. *Assume that conditions (1.2)-(1.5) are satisfied for some $p \geq 1$, then*

- (i) $S_{mn}(x, y)$ converges pointwise to $f(x, y)$ for every $(x, y) \in T^2$ such that $x, y > 0$;
- (ii) $\|S_{mn}(x, y) - f(x, y)\|_r = o(1)$ as $\min\{m, n\} \rightarrow \infty$, $1 \leq r < \infty$.

The above theorem has been proved by Móricz [4, 5] for $p = 1$ and $p = 2$ using suitable estimates for Dirichlet's kernel $D_j(x)$ and Fejér kernel $K_j(x)$. In the case of a single series for $p = 2$, the results regarding convergence have been proved by Kolmogorov [3].

Obviously, condition (1.5) implies any of the following conditions:

$$(1.6) \quad \lim_{\lambda \downarrow 1} \overline{\lim}_{n \rightarrow \infty} \sum_{j=0}^{\infty} \sum_{k=n+1}^{\lambda_n} \frac{\lambda_n - k + 1}{\lambda_n - n} |\Delta_{pp} a_{jk}| (jk)^{p-1} = 0,$$

$$(1.7) \quad \lim_{\lambda \downarrow 1} \overline{\lim}_{m \rightarrow \infty} \sum_{j=m+1}^{\lambda_m} \sum_{k=0}^{\infty} \frac{\lambda_m - j + 1}{\lambda_m - m} |\Delta_{pp} a_{jk}| (jk)^{p-1} = 0.$$

We introduce the following three sums for $m, n \geq 0$ and $\lambda > 1$:

$$\begin{aligned} \sum_{10}^\lambda(m, n, x, y) &= \sum_{j=m+1}^{\lambda_m} \sum_{k=0}^n \frac{\lambda_m - j + 1}{\lambda_m - m} a_{jk} \cos jx \cos ky, \\ \sum_{01}^\lambda(m, n, x, y) &= \sum_{j=0}^m \sum_{k=n+1}^{\lambda_n} \frac{\lambda_n - k + 1}{\lambda_n - n} a_{jk} \cos jx \cos ky, \\ \sum_{11}^\lambda(m, n, x, y) &= \sum_{j=m+1}^{\lambda_m} \sum_{k=n+1}^{\lambda_n} \frac{\lambda_m - j + 1}{\lambda_m - m} \frac{\lambda_n - k + 1}{\lambda_n - n} a_{jk} \cos jx \cos ky \end{aligned}$$

and we have

$$\begin{aligned} \sum_{11}^\lambda(m, n; x, y) &= \frac{1}{(\lambda_m - m)} \sum_{u=m+1}^{\lambda_m} \left(\sum_{01}^\lambda(u, n; x, y) - \sum_{01}^\lambda(m, n; x, y) \right), \\ \sum_{11}^\lambda(m, n; x, y) &= \frac{1}{(\lambda_n - n)} \sum_{v=n+1}^{\lambda_n} \left(\sum_{10}^\lambda(m, v; x, y) - \sum_{10}^\lambda(m, n; x, y) \right). \end{aligned}$$

This implies

$$(1.8) \quad \sum_{11}^\lambda(m, n; x, y) \leq \left\{ \begin{array}{l} 2 \sup_{m \leq u \leq \lambda_m} \left(|\sum_{01}^\lambda(u, n; x, y)| \right) \\ 2 \sup_{n \leq v \leq \lambda_n} \left(|\sum_{10}^\lambda(m, v; x, y)| \right) \end{array} \right\}.$$

The second result of this paper is the following theorem.

Theorem 1.2. (i) *Let $E \subset T^2$. Assume that the following conditions are satisfied:*

$$(1.9) \quad \lim_{\lambda \downarrow 1} \overline{\lim}_{m, n \rightarrow \infty} \left(\sup_{(x, y) \in E} |\sum_{10}^\lambda(m, n; x, y)| \right) = 0,$$

$$(1.10) \quad \lim_{\lambda \downarrow 1} \overline{\lim}_{m, n \rightarrow \infty} \left(\sup_{(x, y) \in E} |\sum_{01}^\lambda(m, n; x, y)| \right) = 0.$$

If $V_{mn}^\lambda(x, y)$ converges uniformly to $f(x, y)$ on $E \subset T^2$ as $\min\{m, n\} \rightarrow \infty$ (that is, in the unrestricted sense), then so does S_{mn} .

(ii) *Assume that the following conditions are satisfied for some $r \geq 1$:*

$$(1.11) \quad \begin{aligned} \lim_{\lambda \downarrow 1} \overline{\lim}_{m, n \rightarrow \infty} \left(\|\sum_{10}^\lambda(m, n; x, y)\|_r \right) &= 0, \\ \lim_{\lambda \downarrow 1} \overline{\lim}_{m, n \rightarrow \infty} \left(\|\sum_{01}^\lambda(m, n; x, y)\|_r \right) &= 0. \end{aligned}$$

If $\|V_{mn}^\lambda - f\|_r \rightarrow 0$ unrestrictedly then $\|S_{mn} - f\|_r \rightarrow 0$ as $\min\{m, n\} \rightarrow \infty$.

We will also prove the following theorem.

Theorem 1.3. *Assume that the conditions (1.2)-(1.4) and (1.6)-(1.7) are satisfied for some $p \geq 1$, then*

- (i) *if $V_{mn}^\lambda(x, y)$ converges uniformly to $f(x, y)$ as $\min\{m, n\} \rightarrow \infty$, then so does S_{mn} ;*
- (ii) *if $\|V_{mn}^\lambda - f\|_r \rightarrow 0$ unrestrictedly for some r with $1 \leq r < \infty$, then $\|S_{mn} - f\|_r \rightarrow 0$ as $\min\{m, n\} \rightarrow \infty$.*

2. NOTATION AND FORMULAS

We define for every $\alpha = 0, 1, 2, \dots$ the sequence $S_0^\alpha, S_1^\alpha, S_2^\alpha, \dots$ by the conditions

$$S_n^0 = S_n, \quad S_n^\alpha = \sum_{u=0}^n S_u^{\alpha-1}, \quad \alpha \geq 1$$

and

$$A_n^0 = 1, \quad A_n^\alpha = \sum_{u=0}^n A_u^{\alpha-1}, \quad \alpha \geq 1,$$

denotes binomial coefficients. Also

$$A_n^\alpha = \binom{n + \alpha}{n} \simeq \frac{n^\alpha}{\Gamma(\alpha + 1)}, \quad \alpha \neq -1, -2, -3, \dots$$

The Cesàro means T_n^α of order α of $\sum a_n$ will be defined by $T_n^\alpha = \frac{S_n^\alpha}{A_n^\alpha}$ and also it is known [8] that $\int_0^\pi |T_n^\alpha(x)| dx, \alpha > 0$, is bounded for all n .

3. LEMMAS

We require the following lemmas for the proof of our results.

Lemma 3.1. *For $m, n \geq 0$ and $p > 1$, the following representation holds:*

$$\begin{aligned} S_{mn}(x, y) &= \sum_{j=0}^m \sum_{k=0}^n \lambda_j \lambda_k a_{jk} \cos jx \cos ky \\ &= \sum_{j=0}^m \sum_{k=0}^n \Delta_{pp} a_{jk} S_j^{p-1}(x) S_k^{p-1}(y) + \sum_{j=0}^m \sum_{t=0}^{p-1} \Delta_{pt} a_{j,n+1} S_j^{p-1}(x) S_n^t(y) \\ &\quad + \sum_{k=0}^n \sum_{s=0}^{p-1} \Delta_{sp} a_{m+1,k} S_m^s(x) S_k^{p-1}(y) + \sum_{s=0}^{p-1} \sum_{t=0}^{p-1} \Delta_{st} a_{m+1,n+1} S_m^s(x) S_n^t(y). \end{aligned}$$

Lemma 3.2 ([1]). *For $m, n \geq 0$ and $\lambda > 1$, the following representation holds:*

$$\begin{aligned} S_{mn} - \sigma_{mn} &= \frac{\lambda_m + 1}{\lambda_m - m} \frac{\lambda_n + 1}{\lambda_n - n} (\sigma_{\lambda_m, \lambda_n} - \sigma_{\lambda_m, n} - \sigma_{m, \lambda_n} + \sigma_{mn}) \\ &\quad + \frac{\lambda_m + 1}{\lambda_m - m} (\sigma_{\lambda_m, n} - \sigma_{mn}) + \frac{\lambda_n + 1}{\lambda_n - n} (\sigma_{m, \lambda_n} - \sigma_{mn}) \\ &\quad - \sum_{11}^\lambda(m, n, x, y) - \sum_{10}^\lambda(m, n, x, y) - \sum_{01}^\lambda(m, n, x, y). \end{aligned}$$

Lemma 3.3. For $m, n \geq 0$ and $\lambda > 1$, we have the following representation:

$$V_{mn}^\lambda - S_{mn} = \sum_{11}^\lambda(m, n, x, y) + \sum_{10}^\lambda(m, n, x, y) + \sum_{01}^\lambda(m, n, x, y).$$

Proof. We have

$$V_{mn}^\lambda(x, y) = \frac{1}{(\lambda_m - m)(\lambda_n - n)} \sum_{j=m+1}^{\lambda_m} \sum_{k=n+1}^{\lambda_n} S_{jk}(x, y).$$

Performing double summation by parts, we have

$$\begin{aligned} V_{mn}^\lambda &= \frac{\lambda_m + 1}{\lambda_m - m} \frac{\lambda_n + 1}{\lambda_n - n} \sigma_{\lambda_m, \lambda_n} - \frac{\lambda_m + 1}{\lambda_m - m} \frac{n + 1}{\lambda_n - n} \sigma_{\lambda_m, n} \\ &\quad - \frac{m + 1}{\lambda_m - m} \frac{\lambda_n + 1}{\lambda_n - n} \sigma_{m, \lambda_n} + \frac{m + 1}{\lambda_m - m} \frac{n + 1}{\lambda_n - n} \sigma_{mn} \\ &= \frac{\lambda_m + 1}{\lambda_m - m} \frac{\lambda_n + 1}{\lambda_n - n} (\sigma_{\lambda_m, \lambda_n} - \sigma_{\lambda_m, n} - \sigma_{m, \lambda_n} + \sigma_{mn}) \\ &\quad + \frac{\lambda_m + 1}{\lambda_m - m} (\sigma_{\lambda_m, n} - \sigma_{mn}) + \frac{\lambda_n + 1}{\lambda_n - n} (\sigma_{m, \lambda_n} - \sigma_{mn}) + \sigma_{mn}. \end{aligned}$$

The use of Lemma 3.2, gives

$$\begin{aligned} V_{mn}^\lambda - S_{mn} &= \sum_{j=m+1}^{\lambda_m} \sum_{k=n+1}^{\lambda_n} \frac{\lambda_m - j + 1}{\lambda_m - m} \frac{\lambda_n - k + 1}{\lambda_n - n} a_{jk} \cos jx \cos ky \\ &\quad + \sum_{j=m+1}^{\lambda_m} \sum_{k=0}^n \frac{\lambda_m - j + 1}{\lambda_m - m} a_{jk} \cos jx \cos ky \\ &\quad + \sum_{j=0}^m \sum_{k=n+1}^{\lambda_n} \frac{\lambda_n - k + 1}{\lambda_n - n} a_{jk} \cos jx \cos ky. \end{aligned} \quad \square$$

Lemma 3.4. For $m, n \geq 0$ and $\lambda > 1$, we have the following: representation

$$\begin{aligned} \sum_{10}^\lambda(m, n; x, y) &= \sum_{j=m+1}^{\lambda_m} \sum_{k=0}^n \frac{\lambda_m - j + 1}{\lambda_m - m} a_{jk} \cos jx \cos ky \\ &= \sum_{j=m+1}^{\lambda_m} \sum_{k=0}^n \frac{\lambda_m - j + 1}{\lambda_m - m} \Delta_{pp} a_{jk} S_j^{p-1}(x) S_k^{p-1}(y) \\ &\quad + \sum_{j=m+1}^{\lambda_m} \sum_{t=0}^{p-1} \frac{\lambda_m - j + 1}{\lambda_m - m} \Delta_{pt} a_{j, n+1} S_j^{p-1}(x) S_n^t(y) \\ &\quad + \frac{1}{\lambda_m - m} \sum_{j=m+1}^{\lambda_m} \sum_{s=0}^{p-1} \sum_{k=0}^n \Delta_{sp} a_{j+1, k} S_j^s(x) S_k^{p-1}(y) \\ &\quad + \frac{1}{\lambda_m - m} \sum_{j=m+1}^{\lambda_m} \sum_{s=0}^{p-1} \sum_{t=0}^{p-1} \Delta_{st} a_{j+1, n+1} S_j^s(x) S_n^t(y) \end{aligned}$$

$$\begin{aligned}
 & - \sum_{s=0}^{p-1} \sum_{k=0}^n \Delta_{sp} a_{m+1,k} S_m^s(x) S_k^{p-1}(y) \\
 & - \sum_{s=0}^{p-1} \sum_{t=0}^{p-1} \Delta_{st} a_{m+1,n+1} S_m^s(x) S_n^t(y).
 \end{aligned}$$

Proof. We have by summation by parts,

$$\begin{aligned}
 & \sum_{10}^\lambda(m, n; x, y) \\
 & = \sum_{k=0}^n \cos ky \left(\sum_{j=m+1}^{\lambda_m} \frac{\lambda_m - j + 1}{\lambda_m - m} a_{jk} \cos jx \right) \\
 & = \sum_{k=0}^n \cos ky \left(\sum_{j=m+1}^{\lambda_m} \frac{\lambda_m - j + 1}{\lambda_m - m} \Delta_{p0} a_{jk} S_j^{p-1}(x) \right. \\
 & \quad \left. + \frac{1}{\lambda_m - m} \sum_{j=m+1}^{\lambda_m} \sum_{s=0}^{p-1} \Delta_{s0} a_{j+1,k} S_j^s(x) - \sum_{s=0}^{p-1} \Delta_{s0} a_{m+1,k} S_m^s(x) \right) \\
 & = \sum_{j=m+1}^{\lambda_m} \frac{\lambda_m - j + 1}{\lambda_m - m} S_j^{p-1}(x) \left(\sum_{k=0}^n \Delta_{p0} a_{jk} \cos ky \right) \\
 & \quad + \frac{1}{\lambda_m - m} \sum_{j=m+1}^{\lambda_m} \sum_{s=0}^{p-1} \left(\sum_{k=0}^n \Delta_{s0} a_{j+1,k} \cos ky \right) S_j^s(x) \\
 & \quad - \sum_{s=0}^{p-1} \left(\sum_{k=0}^n \Delta_{s0} a_{m+1,k} \cos ky \right) S_m^s(x) \\
 & = \sum_{j=m+1}^{\lambda_m} \frac{\lambda_m - j + 1}{\lambda_m - m} S_j^{p-1}(x) \left(\sum_{k=0}^n \Delta_{pp} a_{jk} S_k^{p-1}(y) + \sum_{t=0}^{p-1} \Delta_{pt} a_{j,n+1} S_n^t(y) \right) \\
 & \quad + \frac{1}{\lambda_m - m} \sum_{j=m+1}^{\lambda_m} \sum_{s=0}^{p-1} \left(\sum_{k=0}^n \Delta_{sp} a_{j+1,k} S_k^{p-1}(y) + \sum_{t=0}^{p-1} \Delta_{st} a_{j+1,n+1} S_n^t(y) \right) S_j^s(x) \\
 & \quad - \sum_{s=0}^{p-1} \left(\sum_{k=0}^n \Delta_{sp} a_{m+1,k} S_k^{p-1}(y) + \sum_{t=0}^{p-1} \Delta_{st} a_{m+1,n+1} S_n^t(y) \right) S_m^s(x).
 \end{aligned}$$

Similarly we can have representation for $\sum_{01}^\lambda(m, n; x, y)$. □

4. PROOF OF THEOREMS

Proof of Theorem 1.1. For $m, n \geq 0$ and $p > 1$, we have from Lemma 3.1,

$$\begin{aligned}
 S_{mn}(x, y) & = \sum_{j=0}^m \sum_{k=0}^n \Delta_{pp} a_{jk} S_j^{p-1}(x) S_k^{p-1}(y) + \sum_{j=0}^m \sum_{t=0}^{p-1} \Delta_{pt} a_{j,n+1} S_j^{p-1}(x) S_n^t(y) \\
 & \quad + \sum_{k=0}^n \sum_{s=0}^{p-1} \Delta_{sp} a_{m+1,k} S_m^s(x) S_k^{p-1}(y) + \sum_{s=0}^{p-1} \sum_{t=0}^{p-1} \Delta_{st} a_{m+1,n+1} S_m^s(x) S_n^t(y)
 \end{aligned}$$

$$= \sum_1 + \sum_2 + \sum_3 + \sum_4.$$

Using the results as given in [6] that $S_j^p(x) = O\left(\frac{1}{x^p}\right)$, for all $p \geq 2$, $0 < x \leq \pi$, etc, we have for $0 < x, y \leq \pi$,

$$\sum_{j=0}^{\infty} \sum_{k=0}^{\infty} |\Delta_{pp} a_{jk} S_j^{p-1}(x) S_k^{p-1}(y)| < \infty \quad (\text{by (1.2)})$$

and also by (1.3)-(1.5), we have

$$\begin{aligned} \sum_{j=0}^m \sum_{t=0}^{p-1} \Delta_{pt} a_{j,n+1} &\leq \sum_{t=0}^{p-1} \sum_{v=0}^t \binom{t}{v} \left(\sum_{j=0}^m |\Delta_{p0} a_{j,n+v+1}| \right) \\ &\leq \sup_{n < k \leq n+p} \sum_{j=0}^m |\Delta_{p0} a_{jk}| \\ &\leq \sup_{n < k \leq n+p} \sum_{j=0}^m |\Delta_{p0} a_{jk}| \rightarrow 0 \text{ as } \min\{m, n\} \rightarrow \infty. \end{aligned}$$

Thus

$$\sum_{j=0}^m \sum_{t=0}^{p-1} \Delta_{pt} a_{j,n+1} S_j^{p-1}(x) S_n^t(y) \rightarrow 0 \text{ as } \min\{m, n\} \rightarrow \infty.$$

And similarly

$$\begin{aligned} \sum_{s=0}^{p-1} \sum_{k=0}^n \Delta_{sp} a_{m+1,k} &\leq \sum_{s=0}^{p-1} \sum_{u=0}^s \binom{s}{u} \left(\sum_{k=0}^n |\Delta_{0p} a_{m+u+1,k}| \right) \\ &\leq \sup_{m < j \leq m+p} \sum_{k=0}^n |\Delta_{0p} a_{jk}| \\ &\leq \sup_{m < j \leq m+p} \sum_{k=0}^n |\Delta_{0p} a_{jk}| \rightarrow 0 \text{ as } \min\{m, n\} \rightarrow \infty. \end{aligned}$$

Thus

$$\sum_{k=0}^n \sum_{s=0}^{p-1} \Delta_{sp} a_{m+1,k} S_m^s(x) S_k^{p-1}(y) \rightarrow 0,$$

as $\min\{m, n\} \rightarrow \infty$. Also

$$\begin{aligned} \sum_{s=0}^{p-1} \sum_{t=0}^{p-1} \Delta_{st} a_{m+1,n+1} &\leq \sum_{s=0}^{p-1} \sum_{t=0}^{p-1} \sum_{u=0}^s \sum_{v=0}^t \binom{s}{u} \binom{t}{v} |\Delta_{00} a_{m+u+1,n+v+1}| \\ &\leq \sup_{j > m, k > n} |a_{jk}| \rightarrow 0 \text{ as } \min\{m, n\} \rightarrow \infty. \end{aligned}$$

So,

$$\sum_{s=0}^{p-1} \sum_{t=0}^{p-1} \Delta_{st} a_{m+1,n+1} S_m^s(x) S_n^t(y) \rightarrow 0 \text{ as } \min\{m, n\} \rightarrow \infty.$$

Consequently, series (1.1) converges to the function $f(x, y)$ where

$$f(x, y) = \sum_{j=0}^{\infty} \sum_{k=0}^{\infty} \Delta_{pp} a_{jk} S_j^{p-1}(x) S_k^{p-1}(y) \quad \text{and} \quad \lim_{m,n \rightarrow \infty} S_{mn}(x, y) = f(x, y).$$

Now we will calculate $\|\Sigma_1\|$, $\|\Sigma_2\|$, $\|\Sigma_3\|$ and $\|\Sigma_4\|$ in the following way:

$$\begin{aligned} \|\Sigma_1\| &= \left\| \sum_{j=0}^m \sum_{k=0}^n \Delta_{pp} a_{jk} S_j^{p-1}(x) S_k^{p-1}(y) \right\| \\ &\leq \sum_{j=0}^m \sum_{k=0}^n |\Delta_{pp} a_{jk}| \int_0^\pi \int_0^\pi |S_j^{p-1}(x) S_k^{p-1}(y)| dx dy \\ &\leq \sum_{j=0}^m \sum_{k=0}^n |\Delta_{pp} a_{jk}| A_j^{p-1} A_k^{p-1} \int_0^\pi \int_0^\pi |T_j^{p-1}(x) T_k^{p-1}(y)| dx dy \\ &\leq C_p \sum_{j=0}^m \sum_{k=0}^n |\Delta_{pp} a_{jk}| j^{p-1} k^{p-1}, \\ \|\Sigma_2\| &= \left\| \sum_{j=0}^m \sum_{t=0}^{p-1} \Delta_{pt} a_{j,n+1} S_j^{p-1}(x) S_n^t(y) \right\| \\ &\leq \sum_{t=0}^{p-1} \sum_{v=0}^t \binom{t}{v} \left(\sum_{j=0}^m |\Delta_{p0} a_{j,n+v+1}| \right) A_j^{p-1} A_n^t \int_{-\pi}^\pi \int_{-\pi}^\pi |T_j^{p-1}(x) T_n^t(y)| dx dy \\ &\leq C_p \sup_{n < k \leq n+p} \sum_{j=0}^m |\Delta_{p0} a_{jk}| j^{p-1} \left(\sum_{t=0}^{p-1} n^t \right) \\ &\leq C_p \sup_{n < k \leq n+p} \sum_{j=0}^m |\Delta_{p0} a_{jk}| j^{p-1} k^{p-1}, \\ \|\Sigma_3\| &= \left\| \sum_{s=0}^{p-1} \sum_{k=0}^n \Delta_{sp} a_{m+1,k} S_m^s(x) S_k^{p-1}(y) \right\| \\ &\leq \sum_{s=0}^{p-1} \sum_{u=0}^s \binom{s}{u} \left(\sum_{k=0}^n |\Delta_{0p} a_{m+u+1,k}| \right) A_m^s A_k^{p-1} \int_0^\pi \int_0^\pi |T_m^s(x) T_k^{p-1}(y)| dx dy \\ &\leq C_p \sup_{m < j \leq m+p} \sum_{k=0}^n |\Delta_{0p} a_{jk}| k^{p-1} \left(\sum_{s=0}^{p-1} m^s \right) \\ &\leq C_p \sup_{m < j \leq m+p} \sum_{k=0}^n |\Delta_{0p} a_{jk}| j^{p-1} k^{p-1}, \\ \|\Sigma_4\| &= \left\| \sum_{s=0}^{p-1} \sum_{t=0}^{p-1} \Delta_{st} a_{m+1,n+1} S_m^s(x) S_n^t(y) \right\| \end{aligned}$$

$$\begin{aligned} &\leq \sum_{s=0}^{p-1} \sum_{t=0}^{p-1} \sum_{u=0}^s \sum_{v=0}^t \binom{s}{u} \binom{t}{v} |\Delta_{00} a_{m+u+1, n+v+1}| A_m^s A_n^t \int_0^\pi \int_0^\pi |T_m^s(x) T_n^t(y)| dx dy \\ &\leq C_p \sup_{j>m, k>n} |a_{jk}| j^{p-1} k^{p-1}. \end{aligned}$$

Now let R_{mn} consists of all (j, k) with $j > m$ or $k > n$, that is,

$$\sum \sum_{(j,k) \in R_{mn}} = \sum_{j=m+1}^\infty \sum_{k=0}^n + \sum_{j=0}^\infty \sum_{k=n+1}^\infty + \sum_{j=m+1}^\infty \sum_{k=n+1}^\infty .$$

Then

$$\begin{aligned} \|f - S_{mn}\|_r &= \left(\int_0^\pi \int_0^\pi |f(x, y) - S_{mn}(x, y)|^r dx dy \right)^{1/r}, \quad 1 \leq r < \infty, \\ &\leq \left\| \sum_{(j,k) \in R_{mn}} \Delta_{pp} a_{jk} S_j^{p-1}(x) S_k^{p-1}(y) \right\|_r \\ &\quad + \left\| \sum_{j=0}^m \sum_{t=0}^{p-1} \Delta_{pt} a_{j, n+1} S_j^{p-1}(x) S_n^t(y) \right\|_r \\ &\quad + \left\| \sum_{k=0}^n \sum_{s=0}^{p-1} \Delta_{sp} a_{m+1, k} S_m^s(x) S_k^{p-1}(y) \right\|_r \\ &\quad + \left\| \sum_{s=0}^{p-1} \sum_{t=0}^{p-1} \Delta_{st} a_{m+1, n+1} S_m^s(x) S_n^t(y) \right\|_r \\ &\leq C_{pr} \left\{ \left(\sum_{(j,k) \in R_{mn}} |\Delta_{pp} a_{jk}| j^{p-1} k^{p-1} \right) \right. \\ &\quad + \left(\sup_{n < k \leq n+p} \sum_{j=0}^m |\Delta_{p0} a_{jk}| j^{p-1} k^{p-1} \right) \\ &\quad + \left(\sup_{m < j \leq m+p} \sum_{k=0}^n |\Delta_{0p} a_{jk}| j^{p-1} k^{p-1} \right) \\ &\quad \left. + \left(\sup_{j>m, k>n} |a_{jk}| j^{p-1} k^{p-1} \right) \right\} \quad (\text{as discussed above}) \\ &\rightarrow 0 \quad \text{as } \min\{m, n\} \rightarrow \infty \quad (\text{by (1.2)-(1.5)}), \end{aligned}$$

which proves (ii) part.

Proof of Theorem 1.2. Using the relation (1.8), we find that (1.9) or (1.10) implies

$$(4.1) \quad \lim_{\lambda \downarrow 1} \overline{\lim}_{m, n \rightarrow \infty} \left(\sup_{(x,y) \in E} |\sum_{11}^\lambda(m, n; x, y)| \right) = 0.$$

Assume that $V_{mn}^\lambda(x, y)$ converges uniformly on E to $f(x, y)$. Then by Lemma 3.3, we get

$$\begin{aligned} \overline{\lim}_{m,n \rightarrow \infty} \left(\left\| \sup_{(x,y) \in E} (S_{mn}(x, y) - V_{mn}^\lambda(x, y)) \right\| \right) &\leq \overline{\lim}_{m,n \rightarrow \infty} \left(\sup_{(x,y) \in E} \left| \sum_{10}^\lambda(m, n; x, y) \right| \right) \\ &+ \overline{\lim}_{m,n \rightarrow \infty} \left(\sup_{(x,y) \in E} \left| \sum_{01}^\lambda(m, n; x, y) \right| \right) \\ &+ \overline{\lim}_{m,n \rightarrow \infty} \left(\sup_{(x,y) \in E} \left| \sum_{11}^\lambda(m, n; x, y) \right| \right). \end{aligned}$$

After taking $\lambda \downarrow 1$ the result follows from (1.9), (1.10) and (4.1).

For (ii) part of theorem, we have

$$\begin{aligned} \left\| \sum_{11}^\lambda(m, n; x, y) \right\|_r &= \frac{1}{\lambda_m - m} \sum_{u=m+1}^{\lambda_m} \left(\left\| \sum_{01}^\lambda(u, n; x, y) \right\|_r + \left\| \sum_{01}^\lambda(m, n; x, y) \right\|_r \right) \\ &\leq 2 \left(\sup_{m \leq u \leq \lambda_m} \left(\left\| \sum_{01}^\lambda(u, n; x, y) \right\|_r \right) \right). \end{aligned}$$

Thus (1.11) implies

$$\lim_{\lambda \downarrow 1} \overline{\lim}_{m,n \rightarrow \infty} \left\| \sum_{11}^\lambda(m, n; x, y) \right\|_r = 0.$$

Thus the result of Theorem 1.2 (ii) follows.

Proof of Theorem 1.3. Using the Lemma 3.4, we can write the expression for $\sum_{01}^\lambda(m, n; x, y)$ as

$$\begin{aligned} \sum_{01}^\lambda(m, n; x, y) &= \sum_{j=0}^m \sum_{k=n+1}^{\lambda_n} \frac{\lambda_n - k + 1}{\lambda_n - n} a_{jk} \cos jx \cos ky \\ &= \sum_{j=0}^m \sum_{k=n+1}^{\lambda_n} \frac{\lambda_n - k + 1}{\lambda_n - n} \Delta_{pp} a_{jk} S_j^{p-1}(x) S_k^{p-1}(y) \\ &\quad + \sum_{k=n+1}^{\lambda_n} \sum_{s=0}^{p-1} \frac{\lambda_n - k + 1}{\lambda_n - n} \Delta_{sp} a_{m+1,k} S_m^s(x) S_k^{p-1}(y) \\ &\quad + \frac{1}{\lambda_n - n} \sum_{j=0}^m \sum_{k=n+1}^{\lambda_n} \sum_{t=0}^{p-1} \Delta_{pt} a_{j,k+1} S_j^{p-1}(x) S_k^t(y) \\ &\quad + \frac{1}{\lambda_n - n} \sum_{k=n+1}^{\lambda_n} \sum_{s=0}^{p-1} \sum_{t=0}^{p-1} \Delta_{st} a_{m+1,k+1} S_m^s(x) S_k^t(y) \\ &\quad - \sum_{t=0}^{p-1} \sum_{j=0}^m \Delta_{pt} a_{j,n+1} S_j^{p-1}(x) S_n^t(y) \end{aligned}$$

$$\begin{aligned}
 & - \sum_{s=0}^{p-1} \sum_{t=0}^{p-1} \Delta_{st} a_{m+1, n+1} S_m^s(x) S_n^t(y) \\
 & = I_1 + I_2 + I_3 + I_4 + I_5 + I_6.
 \end{aligned}$$

Now by using (1.2)-(1.4) and (1.6) along with estimates of $S_j^{p-1}(x)$ etc., as mentioned in [6], we have the following estimates in brief:

$$\begin{aligned}
 |I_1| & = \left| \sum_{j=0}^m \sum_{k=n+1}^{\lambda_n} \frac{\lambda_n - k + 1}{\lambda_n - n} \Delta_{pp} a_{jk} S_j^{p-1}(x) S_k^{p-1}(y) \right| \\
 & \leq \sum_{j=0}^m \sum_{k=n+1}^{\lambda_n} \frac{\lambda_n - k + 1}{\lambda_n - n} |\Delta_{pp} a_{jk}| \\
 & \rightarrow 0 \quad \text{as } \min\{m, n\} \rightarrow \infty.
 \end{aligned}$$

Consequently, $\lim_{\lambda \downarrow 1} \overline{\lim}_{m, n \rightarrow \infty} \left(\sup_{(x, y) \in E} |I_1| \right) \rightarrow 0$ as $\min\{m, n\} \rightarrow \infty$. Also,

$$\begin{aligned}
 |I_2| & = \left| \sum_{k=n+1}^{\lambda_n} \sum_{s=0}^{p-1} \frac{\lambda_n - k + 1}{\lambda_n - n} \Delta_{sp} a_{m+1, k} S_m^s(x) S_k^{p-1}(y) \right| \\
 & \leq \sum_{s=0}^{p-1} \sum_{u=0}^s \binom{s}{u} \sum_{k=n+1}^{\lambda_n} |\Delta_{0p} a_{m+u+1, k}| \\
 & \leq \sup_{m < j \leq m+p} \sum_{k=n+1}^{\lambda_n} |\Delta_{0p} a_{jk}| \rightarrow 0 \quad \text{as } \min\{m, n\} \rightarrow \infty.
 \end{aligned}$$

So, $\lim_{\lambda \downarrow 1} \overline{\lim}_{m, n \rightarrow \infty} \left(\sup_{(x, y) \in E} |I_2| \right) \rightarrow 0$ as $\min\{m, n\} \rightarrow \infty$. Also,

$$\begin{aligned}
 |I_3| & \leq \sup_{n < k \leq \lambda_n} \sum_{t=0}^{p-1} \sum_{j=0}^m |\Delta_{pt} a_{j, k+1}| \\
 & \leq \sup_{n < k \leq \lambda_n} \sum_{t=0}^{p-1} \sum_{v=0}^t \binom{t}{v} \sum_{j=0}^m |\Delta_{pt} a_{j, k+v+1}| \\
 & \leq \sup_{n < k \leq \lambda_n + p} \sum_{j=0}^m |\Delta_{p0} a_{jk}| \rightarrow 0 \quad \text{as } \min\{m, n\} \rightarrow \infty,
 \end{aligned}$$

which implies $\lim_{\lambda \downarrow 1} \overline{\lim}_{m, n \rightarrow \infty} \left(\sup_{(x, y) \in E} |I_3| \right) \rightarrow 0$ as $\min\{m, n\} \rightarrow \infty$. Now,

$$\begin{aligned}
 |I_4| & \leq \sup_{n < k \leq \lambda_n} \sum_{s=0}^{p-1} \sum_{t=0}^{p-1} |\Delta_{st} a_{m+1, k+1}| \\
 & \leq \sup_{j > m, k > n} |a_{jk}| \rightarrow 0 \quad \text{as } \min\{m, n\} \rightarrow \infty.
 \end{aligned}$$

Thus $\lim_{\lambda \downarrow 1} \overline{\lim}_{m,n \rightarrow \infty} \left(\sup_{(x,y) \in E} |I_4| \right) \rightarrow 0$ as $\min\{m, n\} \rightarrow \infty$. Also,

$$|I_5| \leq \sum_{t=0}^{p-1} \sum_{v=0}^t \binom{t}{v} \sum_{j=0}^m |\Delta_{p0} a_{j,n+v+1}| \leq \sup_{n < k \leq n+p} \sum_{j=0}^m |\Delta_{p0} a_{jk}| \rightarrow 0 \text{ as } \min\{m, n\} \rightarrow \infty,$$

which implies $\lim_{\lambda \downarrow 1} \overline{\lim}_{m,n \rightarrow \infty} \left(\sup_{(x,y) \in E} |I_5| \right) \rightarrow 0$ as $\min\{m, n\} \rightarrow \infty$. Also,

$$\begin{aligned} |I_6| &\leq \sum_{s=0}^{p-1} \sum_{t=0}^{p-1} \sum_{u=0}^s \sum_{v=0}^t \binom{s}{u} \binom{t}{v} |\Delta_{00} a_{m+u+1,n+v+1}| \\ &\leq \sup_{j > m, k > n} |a_{jk}| \rightarrow 0 \text{ as } \min\{m, n\} \rightarrow \infty, \end{aligned}$$

and

$$\lim_{\lambda \downarrow 1} \overline{\lim}_{m,n \rightarrow \infty} \left(\sup_{(x,y) \in E} |I_6| \right) \rightarrow 0 \text{ as } \min\{m, n\} \rightarrow \infty.$$

Thus combining all these, we have

$$\lim_{\lambda \downarrow 1} \overline{\lim}_{m,n \rightarrow \infty} \left(\sup_{(x,y) \in E} \left| \sum_{01}^\lambda(m, n; x, y) \right| \right) = 0.$$

Similarly (1.2)–(1.4) and (1.7) results in

$$\lim_{\lambda \downarrow 1} \overline{\lim}_{m,n \rightarrow \infty} \left(\sup_{(x,y) \in E} \left| \sum_{10}^\lambda(m, n; x, y) \right| \right) = 0.$$

Thus, first part of theorem follows from Theorem 1.2.

Proof of (ii). We have

$$\|S_{mn} - f\|_r \leq \|S_{mn} - V_{mn}^\lambda\|_r + \|V_{mn}^\lambda - f\|_r.$$

By assumption $\|V_{mn}^\lambda - f\|_r \rightarrow 0$, so it is sufficient to show that

$$\|S_{mn} - V_{mn}^\lambda\|_r \rightarrow 0 \text{ as } \min\{m, n\} \rightarrow \infty.$$

By Lemma 3.3, we have

$$\begin{aligned} \|S_{mn} - V_{mn}^\lambda\|_r &\leq \left\| \sum_{10}^\lambda(m, n; x, y) \right\|_r + \left\| \sum_{01}^\lambda(m, n; x, y) \right\|_r \\ &\quad + \left\| \sum_{11}^\lambda(m, n; x, y) \right\|_r. \end{aligned}$$

Now in order to estimate $\left\| \sum_{01}^\lambda(m, n; x, y) \right\|_r$, we first find $\|I_1\|, \|I_2\|, \|I_3\|, \|I_4\|, \|I_5\|$ and $\|I_6\|$, so we have

$$\begin{aligned} \|I_1\| &= \left\| \sum_{j=0}^m \sum_{k=n+1}^{\lambda_n} \frac{\lambda_n - k + 1}{\lambda_n - n} \Delta_{pp} a_{jk} S_j^{p-1}(x) S_k^{p-1}(y) \right\| \\ &\leq \sum_{j=0}^m \sum_{k=n+1}^{\lambda_n} \frac{\lambda_n - k + 1}{\lambda_n - n} \Delta_{pp} a_{jk} A_j^{p-1} A_k^{p-1} \int_0^\pi \int_0^\pi |T_j^{p-1}(x) T_k^{p-1}(y)| dx dy \end{aligned}$$

$$\begin{aligned}
 &\leq C_p \sum_{j=0}^m \sum_{k=n+1}^{\lambda_n} \frac{\lambda_n - k + 1}{\lambda_n - n} |\Delta_{pp} a_{jk}| j^{p-1} k^{p-1}, \\
 \|I_2\| &= \left\| \sum_{k=n+1}^{\lambda_n} \sum_{s=0}^{p-1} \frac{\lambda_n - k + 1}{\lambda_n - n} \Delta_{sp} a_{m+1,k} S_m^s(x) S_k^{p-1}(y) \right\| \\
 &\leq C_p \sum_{s=0}^{p-1} \sum_{u=0}^s \binom{s}{u} \sum_{k=n+1}^{\lambda_n} |\Delta_{0p} a_{m+u+1,k}| k^{p-1} m^s \\
 &\leq C_p \sup_{m < j \leq m+p} \left(\sum_{k=n+1}^{\lambda_n} |\Delta_{0p} a_{jk}| k^{p-1} \right) \left(\sum_{s=0}^{p-1} m^s \right) \\
 &\leq C_p \sup_{m < j \leq m+p} \sum_{k=n+1}^{\lambda_n} |\Delta_{0p} a_{jk}| j^{p-1} k^{p-1}, \\
 \|I_3\| &\leq C_p \sup_{n < k \leq \lambda_n} \sum_{t=0}^{p-1} \sum_{j=0}^m |\Delta_{pt} a_{j,k+1}| j^{p-1} k^t \\
 &\leq C_p \sup_{n < k \leq \lambda_n} \sum_{t=0}^{p-1} \sum_{v=0}^t \binom{t}{v} \sum_{j=0}^m |\Delta_{pt} a_{j,k+v+1}| j^{p-1} k^t \\
 &\leq C_p \sup_{n < k \leq \lambda_n + p} \sum_{j=0}^m |\Delta_{p0} a_{jk}| j^{p-1} k^{p-1}, \\
 \|I_4\| &\leq C_p \sup_{n < k \leq \lambda_n} \sum_{s=0}^{p-1} \sum_{t=0}^{p-1} |\Delta_{st} a_{m+1,k+1}| m^s k^t \\
 &\leq C_p \sup_{j > m, k > n} |a_{jk}| j^{p-1} k^{p-1}, \\
 \|I_5\| &\leq C_p \sum_{t=0}^{p-1} \sum_{v=0}^t \binom{t}{v} \sum_{j=0}^m |\Delta_{p0} a_{j,n+v+1}| j^{p-1} n^t \\
 &\leq C_p \sup_{n < k \leq n+p} \sum_{j=0}^m |\Delta_{p0} a_{jk}| j^{p-1} k^{p-1}, \\
 \|I_6\| &\leq C_p \sum_{s=0}^{p-1} \sum_{t=0}^{p-1} \sum_{u=0}^s \sum_{v=0}^t \binom{s}{u} \binom{t}{v} |\Delta_{00} a_{m+u+1,n+v+1}| m^s n^t \\
 &\leq C_p \sup_{j > m, k > n} |a_{jk}| j^{p-1} k^{p-1}.
 \end{aligned}$$

Thus, we can estimate

$$\begin{aligned}
 \left\| \sum_{01}^{\lambda} (m, n; x, y) \right\|_r &\leq C_{pr} \sum_{k=n+1}^{\lambda_n} \sum_{j=0}^m \frac{\lambda_n - k + 1}{\lambda_n - n} |\Delta_{pp} a_{jk}| j^{p-1} k^{p-1} \\
 &\quad + C_{pr} \left(\sup_{m < j \leq m+p} \sum_{k=n+1}^{\lambda_n} |\Delta_{0p} a_{jk}| j^{p-1} k^{p-1} \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ C_{pr} \left(\sup_{n < k \leq \lambda_n + p} \sum_{j=0}^m |\Delta_{p0} a_{jk}| j^{p-1} k^{p-1} \right) \\
 &+ C_{pr} \left(\sup_{j > m, k > n} |a_{jk}| j^{p-1} k^{p-1} \right) \\
 &+ C_{pr} \left(\sup_{n < k \leq n+p} \sum_{j=0}^m |\Delta_{p0} a_{jk}| j^{p-1} k^{p-1} \right) \\
 &+ C_{pr} \left(\sup_{j > m, k > n} |a_{jk}| j^{p-1} k^{p-1} \right).
 \end{aligned}$$

By (1.2)–(1.4) and (1.6), we conclude that

$$\lim_{\lambda \downarrow 1} \overline{\lim}_{m, n \rightarrow \infty} \left(\left\| \sum_{01}^\lambda (m, n; x, y) \right\|_r \right) = 0.$$

Similarly, by conditions (1.2)–(1.4) and (1.7), we get

$$\lim_{\lambda \downarrow 1} \overline{\lim}_{m, n \rightarrow \infty} \left(\left\| \sum_{10}^\lambda (m, n; x, y) \right\|_r \right) = 0.$$

Also, by (1.8), we have

$$\lim_{\lambda \downarrow 1} \overline{\lim}_{m, n \rightarrow \infty} \left(\left\| \sum_{11}^\lambda (m, n; x, y) \right\|_r \right) = 0.$$

Thus, $\|S_{mn} - V_{mn}^\lambda\|_r \rightarrow 0$ as $\min\{m, n\} \rightarrow \infty$.

Acknowledgements. The authors are very thankful to the referee for his/her valuable suggestions.

REFERENCES

- [1] C. P. Chen and Y. W. Chauang, *L¹-convergence of double Fourier series*, Chin. J. Math. **19**(4) (1991), 391–410.
- [2] K. Kaur, S. S. Bhatia and B. Ram, *L¹-Convergence of complex double trigonometric series*, Proc. Indian Acad. Sci. **113**(3) (2003), 1–9.
- [3] A. N. Kolmogorov, *Sur l'ordre de grandeur des coefficients de la série de Fourier-Lebesgue*, Bulletin L'Academie Polonaise des Science (1923), 83–86.
- [4] F. Móricz, *Convergence and integrability of double trigonometric series with coefficients of bounded variation*, Proc. Amer. Math. Soc. **102** (1988), 633–640.
- [5] F. Móricz, *On the integrability and L¹-convergence of double trigonometric series*, Studia Math. (1991), 203–225.
- [6] T. M. Vukolova, *Certain properties of trigonometric series with monotone coefficients* (English, Russian original), Moscow Univ. Math. Bull. **39**(6) (1984), 24–30, translation from Vestnik Moskov. Univ. **1**(6) (1984), 18–23.
- [7] W. H. Young, *On the Fourier series of bounded functions*, Proc. London Math. Soc. **12**(2) (1913), 41–70.
- [8] A. Zygmund, *Trigonometric Series*, Vols. I, II, Cambridge University Press, Cambridge, 1959.

¹DEPARTMENT OF APPLIED MATHEMATICS,
GZS CAMPUS COLLEGE OF ENGINEERING AND TECHNOLOGY,
MAHARAJA RANJIT SINGH PUNJAB TECHNICAL UNIVERSITY BATHINDA, PUNJAB, INDIA
Email address: karanvir@rsptu.ac.in

²DEPARTMENT OF MATHEMATICS,
AMITY UNIVERSITY OF RAJASTHAN,
JAIPUR, INDIA
Email address: kmodi@jpr.amity.edu