

**ON DEGREE OF APPROXIMATION OF SIGNALS IN THE
GENERALIZED ZYGMUND CLASS BY USING $(E, r)(N, q_n)$ MEAN**

ANWESHA MISHRA¹, BIRUPAKHYA PRASAD PADHY¹, LAKSHMI NARAYAN MISHRA²,
AND UMAKANTA MISRA³

ABSTRACT. In the present article, we have established a result on degree of approximation of function (or signal) in the generalized Zygmund class $Z_l^{(m)}, (l \geq 1)$ by using $(E, r)(N, q_n)$ - mean of Trigonometric Fourier series.

1. INTRODUCTION

Signal Analysis describes the field of study whose objective is to collect, understand and deduce information and intelligence from various signals. Now-a-days the analysis of signals is a fundamental problem for many engineers and scientists. In the recent past, we have seen the applications of mathematical methods such as Probability theory, Mathematical statistics etc. in the analysis of signals. Very recently, approximation theory has got a large popularity as it has given a new dimension in approximating the signals (or functions). The estimation of error functions in Lipschitz class and Zygmund space using different summability techniques of Fourier series and conjugate Fourier series have been of great interest among the researchers in the last decades (for details see [2, 3, 9, 10, 13–18]). Later, the generalized Zygmund class $Z_l^{(m)}, l \geq 1$, is investigated by Leindler [8], Moricz [4], Moricz and Nemeth [5]. Very recently Nigam [7] and Singh et al. [11] proved approximation of functions in the generalized Zygmund class by using Hausdroff means. Lal and Shireen [12] proved a result on approximation of functions of generalized Zygmund class by Matrix Euler summability mean of Fourier series. In the present paper, we investigate on the degree

Key words and phrases. Degree of approximation, generalized Zygmund class, trigonometric Fourier series, (E, r) -summability mean, (N, q_n) -summability mean, $(E, r)(N, q_n)$ -summability mean.
2010 *Mathematics Subject Classification.* Primary: 41A24, 41A25. Secondary: 42B05, 42B08.
Received: January 11, 2020.
Accepted: July 23, 2020.

of approximation of a signal (or function) in the generalized Zygmund class $Z_l^{(m)}$, $l \geq 1$, by $(E, r)(N, q_n)$ product mean of the trigonometric Fourier series.

2. DEFINITIONS AND NOTATIONS

Let h be a function, which is periodic in $[0, 2\pi]$ such that $\int_0^{2\pi} |h(x)|^l dx < \infty$.

We denote

$$L_l[0, 2\pi] = \left\{ h : [0, 2\pi] \rightarrow \mathbb{R} : \int_0^{2\pi} |h(x)|^l dx < \infty \right\}, \quad l \geq 1.$$

The Fourier series of $h(x)$ is given by

$$(2.1) \quad \sum_{n=0}^{\infty} u_n(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx).$$

Let $S_p(h; x)$ denotes the p -th partial sum of $h(x)$ and is given by

$$S_p(h; x) = \frac{1}{\pi} \int_{-\pi}^{\pi} h(x+v) \frac{\sin\left(p + \frac{1}{2}\right)v}{2 \sin \frac{v}{2}} dv.$$

We define

$$\|h\|_l = \left(\frac{1}{2\pi} \int_0^{2\pi} |h(x)|^l dx \right)^{\frac{1}{l}}, \quad 1 \leq l < \infty,$$

and

$$\|h\|_l = \text{esssup}_{0 \leq x \leq 2\pi} |h(x)|, \quad l = \infty.$$

Let the Zygmund modulus of continuity of $h(x)$ be

$$m(h; r) = \sup_{0 \leq r, x \in \mathbb{R}} |h(x+v) + h(x-v) - 2h(x)| \quad (\text{see [1]}).$$

Suppose \mathbf{B} represents the Banach space of all 2π periodic functions which are continuous and defined over $[0, 2\pi]$ under the supremum norm. Clearly,

$$Z_{(\alpha)} = \left\{ h \in \mathbf{B} : |h(x+v) + h(x-v) - 2h(x)| = O(|v|^\alpha), 0 < \alpha \leq 1 \right\}$$

is a Banach space under the norm $\|\cdot\|_{(\alpha)}$ defined by

$$\|h\|_{(\alpha)} = \sup_{0 \leq x \leq 2\pi} |h(x)| + \sup_{x, v \neq 0} \frac{|h(x+v) + h(x-v) - 2h(x)|}{|v|^\alpha}.$$

For $h \in L_l[0, 2\pi]$, $l \geq 1$, the integral Zygmund modulus of continuity is defined by

$$m_l(h; r) = \sup_{0 < v \leq r} \left\{ \frac{1}{2\pi} \int_0^{2\pi} |h(x+v) + h(x-v) - 2h(x)|^l dx \right\}^{\frac{1}{l}},$$

and for $h \in \mathbf{B}$, $l = \infty$,

$$m_\infty(h; r) = \sup_{0 < v \leq r} \max_x |h(x+v) + h(x-v) - 2h(x)|.$$

Clearly, $m_l(h; r) \rightarrow 0$ as $l \rightarrow 0$. Again,

$$Z_{(\alpha), l} = \left\{ h \in L_l[0, 2\pi] : \left(\int_0^{2\pi} |h(x+v) + h(x-v) - 2h(x)|^l dx \right)^{\frac{1}{l}} = O(|v|^\alpha) \right\}$$

is a Banach space under the norm $\|\cdot\|_{(\alpha), l}$ for $0 < \alpha \leq 1$ and $l \geq 1$. Clearly,

$$\|h\|_{(\alpha), l} = \|h\|_l + \sup_{v \neq 0} \frac{\|h(\cdot + v) + h(\cdot - v) - 2h(\cdot)\|_l}{|v|^\alpha}.$$

Let

$$Z^{(m)} = \left\{ h \in \mathbf{B} : |h(x+v) + h(x-v) - 2h(x)| = O(m(v)) \right\},$$

where m is a Zygmund modulus of continuity satisfying

- (a) $m(0) = 0$;
- (b) $m(v_1 + v_2) \leq m(v_1) + m(v_2)$.

Define

$$Z_l^{(m)} = \left\{ h \in L_l : 1 \leq l < \infty, \sup_{v \neq 0} \frac{\|h(\cdot + v) + h(\cdot - v) - 2h(\cdot)\|_l}{m(v)} < \infty \right\},$$

where

$$\|h\|_l^{(m)} = \|h\|_l + \sup_{v \neq 0} \frac{\|h(\cdot + v) + h(\cdot - v) - 2h(\cdot)\|_l}{m(v)}, \quad l \geq 1.$$

Clearly, $\|\cdot\|_l^{(m)}$ is a norm $Z_l^{(m)}$. Also, $Z_l^{(m)}$ is complete since $L_l, l \geq 1$, is complete. So, $Z_l^{(m)}$ is a Banach space under $\|\cdot\|_l^{(m)}$. Again, suppose $m(v)$ and $\mu(v)$ represents the Zygmund moduli of continuity such that $\frac{m(v)}{\mu(v)}$ is positive and non-decreasing then

$$(2.2) \quad \|h\|_l^{(\mu)} \leq \max \left\{ 1, \frac{m(2\pi)}{\mu(2\pi)} \right\} \|h\|_l^{(m)} \leq \infty.$$

Clearly,

$$Z_l^{(m)} \subseteq Z_l^{(\mu)} \subseteq L_l, \quad l \geq 1.$$

Let $\sum u_n$ be an infinite series with sequence of partial sums $\{s_n\}$. Suppose $\{q_k\}$ represents the sequence of non-negative integers such that

$$(2.3) \quad Q_n = \sum_{k=0}^n q_k \rightarrow \infty \quad \text{as } n \rightarrow \infty.$$

If

$$(2.4) \quad \tau_n^N = \frac{1}{Q_n} \sum_{k=0}^n q_{n-k} s_k, \quad n = 0, 1, 2, \dots$$

represents the (N, q_n) mean of $\{s_n\}$ generated by the sequence $\{q_n\}$, then the series $\sum u_n$ is said to be summable to 's' whenever

$$\lim_{n \rightarrow \infty} \tau_n^N \rightarrow s.$$

We know, (N, q_n) method is regular [6]. The (E, r) transform of $\{s_n\}$ is given by

$$(2.5) \quad E_n^r = \frac{1}{(1+r)^n} \sum_{k=0}^n C(n, k) r^{n-k} s_k.$$

If $E_n^r \rightarrow s$ as $n \rightarrow \infty$, then $\sum u_n$ is summable to 's' by (E, r) summability. Also, (E, r) method is regular [6].

The $(E, r)(N, q_n)$ transform of $\{s_n\}$ is given by

$$(2.6) \quad \tau_n^{E_r, N} = \frac{1}{(1+r)^n} \sum_{k=0}^n C(n, k) \left\{ \frac{1}{Q_k} \sum_{\nu=0}^k q_{k-\nu} s_\nu \right\}.$$

The series $\sum u_n$ is summable to s by the $(E, r)(N, q_n)$ transform if $\tau_n^{E_r, N} \rightarrow s$ as $n \rightarrow \infty$.

The following notations are used in the rest part of our paper:

$$\varphi(x, v) = h(x+v) + h(x-v) - 2h(x),$$

$$\kappa_n^{E_r, N}(v) = \frac{1}{2\pi(1+r)^n} \sum_{k=0}^n C(n, k) \left\{ \frac{1}{Q_k} \sum_{\nu=0}^k q_{k-\nu} \frac{\sin\left(\nu + \frac{1}{2}\right)v}{\sin\left(\frac{v}{2}\right)} \right\}.$$

3. KNOWN RESULTS

Using Matrix Euler summability means, Lal and Shireen [12] proved the following theorems.

Theorem 3.1. *Let the lower triangular matrix $A = (a_{n,k})$ satisfy the following conditions:*

$$(3.1) \quad \sum_{k=0}^n a_{n,k} = 1, \quad a_{n,k} \geq 0, \quad n = 0, 1, 2, \dots, \quad k = 0, 1, 2, \dots,$$

$$(3.2) \quad \sum_{k=0}^n |\Delta a_{n,k}| = O\left(\frac{1}{n+1}\right) \quad \text{and} \quad (n+1)a_{n,n} = O(1).$$

The best approximation of the Fourier series (2.1) by Matrix-Euler mean is given by

$$(3.3) \quad E_n(h) = \inf_{t_n^{\Delta, E}} \|t_n^{\Delta, E} - h\|_l^\mu = O\left(\frac{1}{n+1} \int_{\frac{1}{(n+1)}}^\pi \frac{m(v)}{v^2 \mu(v)} dv\right),$$

where

$$t_n^{\Delta, E} = \sum_{k=0}^n a_{n,k} \frac{1}{2^k} \sum_{\nu=0}^k C(k, \nu) s_\nu,$$

represents the Matrix-Euler mean of a 2π periodic and Lebesgue integrable function $h : [0, 2\pi] \rightarrow \mathbb{R}$, that belongs to $Z_l^{(m)}$, $l \geq 1$. Here, m and μ are the Zygmund moduli of continuity and $\frac{m(v)}{\mu(v)}$ is positive and non-decreasing.

Theorem 3.2. *Let $A = (a_{n,k})$ be a lower triangular matrix satisfying (3.1) and (3.2) in Theorem 3.1 along with the condition that $\frac{m(v)}{\mu(v)}$ is non-increasing. Then, for $h \in Z_l^{(m)}$, $l \geq 1$, the best approximation by Matrix-Euler mean $(t_n^{\Delta, E})$ is given by*

$$(3.4) \quad E_n(h) = O \left(\frac{m\left(\frac{1}{n+1}\right)}{\mu\left(\frac{1}{n+1}\right)} \log(n+1)\pi \right).$$

4. MAIN THEOREMS

Theorem 4.1. *Let $h : [0, 2\pi] \rightarrow \mathbb{R}$ be a periodic function (with period 2π) belonging to $Z_l^{(m)}$, $l \geq 1$, which is integrable in the sense of Lebesgue. Then the degree of approximation of h by using $(E, r)(N, q_n)$ mean of (2.1) is given by*

$$(4.1) \quad E_n(h) = \inf_{\tau_n^{E_r, N}} \|\tau_n^{E_r, N} - h\|_l^\mu = O \left(\int_{\frac{1}{n+1}}^\pi \frac{m(v)}{v\mu(v)} dv \right),$$

where $m(v)$ and $\mu(v)$ are the Zygmund moduli of continuity and $\frac{m(v)}{v\mu(v)}$ is positive and non-decreasing.

Theorem 4.2. *The degree of approximation of a 2π periodic and Lebesgue integrable function $h, h : [0, 2\pi] \rightarrow \mathbb{R}$, using $(E, r)(N, q_n)$ mean of (2.1) is given by*

$$(4.2) \quad E_n(h) = \inf_{\tau_n^{E_r, N}} \|\tau_n^{E_r, N} - h\|_l^\mu = O \left(\frac{m\left(\frac{1}{n+1}\right)}{\mu\left(\frac{1}{n+1}\right)} \left(\frac{\pi}{n+1} - \frac{1}{(n+1)^2} \right) \right),$$

where $h \in Z_l^{(m)}$, $l \geq 1$, $m(v)$ and $\mu(v)$ are the Zygmund moduli of continuity and $\frac{m(v)}{v\mu(v)}$ is positive and non-increasing.

We require the below mentioned lemmas to prove our main theorems.

5. LEMMAS

Lemma 5.1.

$$|\kappa_n^{E_r, N}| = O(n), \quad \text{for } 0 \leq v \leq \frac{1}{n+1}.$$

Lemma 5.2.

$$|\kappa_n^{E_r, N}| = O\left(\frac{1}{v}\right), \quad \text{for } \frac{1}{n+1} \leq v \leq \pi.$$

Lemma 5.3. *Let $h \in Z_l^{(m)}$. Then, for $0 < v \leq \pi$,*

- (i) $\|\varphi(\cdot, v)\|_l = O(m(v))$;
- (ii) $\|\varphi(\cdot + y, v) + \varphi(\cdot - y, v) - 2\varphi(\cdot, v)\|_l = O(m(v))$ or $O(m(y))$;

(iii) If $m(v)$ and $\mu(v)$ are as defined in Theorem 4.1, then

$$\|\varphi(\cdot + y, v) + \varphi(\cdot - y, v) - 2\varphi(\cdot, v)\|_l = O\left(\mu(y) \frac{m(v)}{\mu(v)}\right),$$

where $\varphi(x, v) = h(x + v) + h(x - v) - 2h(x)$.

6. PROOF OF THE LEMMAS

Proof of Lemma 5.1. For $0 \leq v \leq \frac{1}{n+1}$ and $\sin nv \leq n \sin v$, we have

$$\begin{aligned} |\kappa_n^{E_r, N}(v)| &= \frac{1}{2\pi(1+r)^n} \left| \sum_{k=0}^n C(n, k) r^{n-k} \left\{ \frac{1}{Q_k} \sum_{\nu=0}^k q_{k-\nu} \frac{\sin\left(\nu + \frac{1}{2}\right)v}{\sin \frac{v}{2}} \right\} \right| \\ &\leq \frac{1}{2\pi(1+r)^n} \left| \sum_{k=0}^n C(n, k) r^{n-k} \left\{ \frac{1}{Q_k} \sum_{\nu=0}^k q_{k-\nu} \frac{(2\nu+1) \sin \frac{v}{2}}{\sin \frac{v}{2}} \right\} \right| \\ &\leq \frac{1}{2\pi(1+r)^n} \left| \sum_{k=0}^n C(n, k) r^{n-k} (2k+1) \left\{ \frac{1}{Q_k} \sum_{\nu=0}^k q_{k-\nu} \right\} \right| \\ &\leq \frac{(2n+1)}{2\pi(1+r)^n} \left| \sum_{k=0}^n C(n, k) r^{n-k} \right| \\ &= O(n). \end{aligned} \quad \square$$

Proof of Lemma 5.2. By Jordan's lemma

$$\sin\left(\frac{v}{2}\right) \geq \frac{v}{\pi}, \quad \sin nv \leq 1, \quad \frac{1}{n+1} \leq v \leq \pi.$$

Now,

$$\begin{aligned} |\kappa_n^{E_r, N}(v)| &= \frac{1}{2\pi(1+r)^n} \left| \sum_{k=0}^n C(n, k) r^{n-k} \left\{ \frac{1}{Q_k} \sum_{\nu=0}^k q_{k-\nu} \frac{\sin\left(\nu + \frac{1}{2}\right)v}{\sin \frac{v}{2}} \right\} \right| \\ &\leq \frac{1}{2\pi(1+r)^n} \left| \sum_{k=0}^n C(n, k) r^{n-k} \left\{ \frac{1}{Q_k} \sum_{\nu=0}^k \frac{\pi}{v} q_{k-\nu} \right\} \right| \\ &= \frac{1}{2v(1+r)^n} \left| \sum_{k=0}^n C(n, k) r^{n-k} \left\{ \frac{1}{Q_k} \sum_{\nu=0}^k q_{k-\nu} \right\} \right| \\ &= \frac{1}{2v(1+r)^n} \left| \sum_{k=0}^n C(n, k) r^{n-k} \right| \\ &= O\left(\frac{1}{v}\right). \end{aligned} \quad \square$$

Proof of Lemma 5.3. See [12]. □

7. PROOF OF MAIN THEOREMS

Proof of Theorem 4.1. Let $S_k(h; x)$ denotes the k -th partial sum of the series (2.1). We have

$$S_k(h; x) - h(x) = \frac{1}{2\pi} \int_0^\pi \varphi(x, v) \frac{\sin\left(k + \frac{1}{2}\right)v}{\sin \frac{v}{2}} dv$$

and the (N, q_n) transform of it is given by

$$\frac{1}{Q_n} \sum_{k=0}^n q_{n-k} \{S_k(h; x) - h(x)\} = \frac{1}{2\pi} \int_0^\pi \varphi(x, v) \frac{1}{Q_n} \sum_{k=0}^n q_{n-k} \frac{\sin\left(k + \frac{1}{2}\right)v}{\sin \frac{v}{2}} dv.$$

Let the $(E, r)(N, q_n)$ transform of $S_k(h; x)$ by $\tau_n^{E_r, N}$. Then

$$\begin{aligned} \tau_n^{E_r, N} - h(x) &= \frac{1}{2\pi(1+r)^n} \int_0^\pi \varphi(x, v) \sum_{k=0}^n C(n, k) r^{n-k} \left\{ \frac{1}{Q_n} \sum_{k=0}^n q_{n-k} \frac{\sin\left(k + \frac{1}{2}\right)v}{\sin \frac{v}{2}} \right\} dv \\ &= \int_0^\pi \varphi(x; v) \kappa_n^{E_r, N}(v) dv \\ &= \chi_n(x). \end{aligned}$$

Then

$$\chi_n(x+y) + \chi_n(x-y) - 2\chi_n(x) = \int_0^\pi \left\{ \varphi(x+y, v) + \varphi(x-y, v) - 2\varphi(x, v) \right\} \kappa_n^{E_r, N}(v) dv.$$

Using Minkowski's inequality, we have

$$\begin{aligned} &\|\chi_n(\cdot + y) + \chi_n(\cdot - y) - 2\chi_n(\cdot)\|_l \\ &= \left\{ \frac{1}{2\pi} \int_0^{2\pi} |\chi_n(x+y) + \chi_n(x-y) - 2\chi_n(x)|^l dx \right\}^{\frac{1}{l}} \\ &= \frac{1}{2\pi} \left[\int_0^{2\pi} \left| \int_0^\pi \left\{ \varphi(x+y, v) + \varphi(x-y, v) - 2\varphi(x, v) \right\} \kappa_n^{E_r, N}(v) dv \right|^l dx \right]^{\frac{1}{l}} \\ &\leq \int_0^\pi |\kappa_n^{E_r, N}(v)| \left\{ \frac{1}{2\pi} \int_0^{2\pi} |\varphi(x+y, v) + \varphi(x-y, v) - 2\varphi(x, v)|^l dx \right\}^{\frac{1}{l}} dv \\ &= \int_0^\pi \|\varphi(\cdot + y, v) + \varphi(\cdot - y, v) - 2\varphi(\cdot, v)\|_l |\kappa_n^{E_r, N}(v)| dv \\ &= \int_0^{\frac{1}{n+1}} \|\varphi(\cdot + y, v) + \varphi(\cdot - y, v) - 2\varphi(\cdot, v)\|_l |\kappa_n^{E_r, N}(v)| dv \\ &\quad + \int_{\frac{1}{n+1}}^\pi \|\varphi(\cdot + y, v) + \varphi(\cdot - y, v) - 2\varphi(\cdot, v)\|_l |\kappa_n^{E_r, N}(v)| dv \\ (7.1) \quad &= \Gamma_1 + \Gamma_2. \end{aligned}$$

Using Lemma 5.1, Lemma 5.3 and monotonicity of $\frac{m(v)}{\mu(v)}$, with respect to 'v', we have

$$\begin{aligned}\Gamma_1 &= \int_0^{\frac{1}{n+1}} \|\varphi(\cdot + y, v) + \varphi(\cdot - y, v) - 2\varphi(\cdot, v)\|_l |\kappa_n^{E_r, N}(v)| dv \\ &= \int_0^{\frac{1}{n+1}} O\left(\mu(y) \frac{m(v)}{\mu(v)}\right) O(n) dv.\end{aligned}$$

By using the second mean value theorem of integral, we have

$$\begin{aligned}(7.2) \quad \Gamma_1 &\leq O\left(\frac{n\mu(y) \frac{m\left(\frac{1}{n+1}\right)}{\mu\left(\frac{1}{n+1}\right)} \int_0^{\frac{1}{n+1}} dv}{\mu\left(\frac{1}{n+1}\right)}\right) \\ &= O\left(\frac{n}{n+1} \mu(y) \frac{m\left(\frac{1}{n+1}\right)}{\mu\left(\frac{1}{n+1}\right)}\right) = O\left(\mu(y) \frac{m\left(\frac{1}{n+1}\right)}{\mu\left(\frac{1}{n+1}\right)}\right).\end{aligned}$$

Again, by using Lemma 5.2 and Lemma 5.3, we get

$$\begin{aligned}(7.3) \quad \Gamma_2 &= \int_{\frac{1}{n+1}}^{\pi} \|\varphi(\cdot + y, v) + \varphi(\cdot - y, v) - 2\varphi(\cdot, v)\|_l |\kappa_n^{E_r, N}(v)| dv \\ &\leq \int_{\frac{1}{n+1}}^{\pi} O\left(\mu(y) \frac{m(v)}{\mu(v)}\right) \frac{1}{v} dv \\ &= O\left(\mu(y) \int_{\frac{1}{n+1}}^{\pi} \frac{m(v)}{v\mu(v)} dv\right).\end{aligned}$$

By (7.1), (7.2) and (7.3), we have

$$\|\chi_n(\cdot + y) + \chi_n(\cdot - y) - 2\chi_n(\cdot)\|_l = O\left(\mu(y) \frac{m\left(\frac{1}{n+1}\right)}{\mu\left(\frac{1}{n+1}\right)}\right) + O\left(\mu(y) \int_{\frac{1}{n+1}}^{\pi} \frac{m(v)}{v\mu(v)} dv\right).$$

Therefore, we have

$$(7.4) \quad \sup_{y \neq 0} \frac{\|\chi_n(\cdot + y) + \chi_n(\cdot - y) - 2\chi_n(\cdot)\|_l}{\mu(y)} = O\left(\frac{m\left(\frac{1}{n+1}\right)}{\mu\left(\frac{1}{n+1}\right)}\right) + O\left(\int_{\frac{1}{n+1}}^{\pi} \frac{m(v)}{v\mu(v)} dv\right).$$

As

$$\varphi(x, v) = |h(x + v) + h(x - v) - 2h(x)|,$$

by applying Minkowski's inequality, we get

$$(7.5) \quad \|\varphi(x, v)\|_l = \|h(x + v) + h(x - v) - 2h(x)\|_l = O(m(v)).$$

Now, using Lemma 5.1, Lemma 5.2 and (7.5),

$$\|\chi_n(\cdot)\|_l \leq \left(\int_0^{\frac{1}{n+1}} + \int_{\frac{1}{n+1}}^{\pi}\right) \|\varphi(\cdot, v)\|_l |\kappa_n^{E_r, N}(v)| dv$$

$$\begin{aligned}
&= O\left(n \int_0^{\frac{1}{n+1}} m(v) dv\right) + O\left(\int_{\frac{1}{n+1}}^{\pi} \frac{m(v)}{v} dv\right) \\
(7.6) \quad &= O\left(m\left(\frac{1}{n+1}\right)\right) + O\left(\int_{\frac{1}{n+1}}^{\pi} \frac{m(v)}{v} dv\right).
\end{aligned}$$

From (7.4) and (7.6), we have

$$\begin{aligned}
&\|\chi_n(\cdot)\|_l^\mu \\
&= \|\chi_n(\cdot)\|_l + \sup_{y \neq 0} \frac{\|\chi_n(\cdot + y) + \chi_n(\cdot - y) - 2\chi_n(\cdot)\|_l}{\mu(y)} \\
&= O\left(m\left(\frac{1}{n+1}\right)\right) + O\left(\int_{\frac{1}{n+1}}^{\pi} \frac{m(v)}{v} dv\right) + O\left(\frac{m\left(\frac{1}{n+1}\right)}{\mu\left(\frac{1}{n+1}\right)}\right) + O\left(\int_{\frac{1}{n+1}}^{\pi} \frac{m(v)}{v \mu(v)} dv\right) \\
&= \sum_{j=1}^4 G_j.
\end{aligned}$$

In view of monotonicity of $\mu(v)$ for $0 < v \leq \pi$, we have

$$m(v) = \frac{m(v)}{\mu(v)} \mu(v) \leq \mu(\pi) \frac{m(v)}{\mu(v)} = O\left(\frac{m(v)}{\mu(v)}\right).$$

Therefore,

$$G_1 = O(G_3).$$

Again, by using monotonicity of $\mu(v)$,

$$G_2 = \int_{\frac{1}{n+1}}^{\pi} \frac{m(v)}{v} dv = \int_{\frac{1}{n+1}}^{\pi} \frac{m(v)}{v \mu(v)} \mu(v) dv \leq \mu(\pi) \int_{\frac{1}{n+1}}^{\pi} \frac{m(v)}{v \mu(v)} dv = O(G_4).$$

Since $\frac{m(v)}{\mu(v)}$ is positive and increasing

$$G_4 = \int_{\frac{1}{n+1}}^{\pi} \frac{m(v)}{v \mu(v)} dv = \frac{m\left(\frac{1}{n+1}\right)}{\mu\left(\frac{1}{n+1}\right)} \int_{\frac{1}{n+1}}^{\pi} \frac{dv}{v} \geq \frac{m\left(\frac{1}{n+1}\right)}{\mu\left(\frac{1}{n+1}\right)}.$$

Therefore,

$$G_3 = O(G_4).$$

Thus,

$$\|\chi_n(\cdot)\|_l^\mu = O(G_4) = O\left(\int_{\frac{1}{n+1}}^{\pi} \frac{m(v)}{v \mu(v)} dv\right).$$

Hence,

$$E_n(h) = \inf_n \|\chi_n(\cdot)\|_l^\mu = O\left(\int_{\frac{1}{n+1}}^{\pi} \frac{m(v)}{v \mu(v)} dv\right).$$

This completes the proof of the Theorem 4.1. \square

Proof of Theorem 4.2. In this theorem, since we have assumed $\frac{m(v)}{v\mu(v)}$ is positive and decreasing, proceeding as in Theorem 4.1. We have

$$E_n(h) = \inf_n \|\chi_n(\cdot)\|_I^\mu = O \left(\frac{m\left(\frac{1}{n+1}\right)}{(n+1)\mu\left(\frac{1}{n+1}\right)} \int_{\frac{1}{n+1}}^\pi dv \right),$$

i.e.,

$$E_n(h) = O \left(\frac{m\left(\frac{1}{n+1}\right)}{\mu\left(\frac{1}{n+1}\right)} \left(\frac{\pi}{(n+1)} - \frac{1}{(n+1)^2} \right) \right).$$

This is what we need to prove in Theorem 4.2. \square

REFERENCES

- [1] A. Zygmund, *Trigonometric Series*, 2nd revised Edition, Cambridge university Press, Cambridge, 1959.
- [2] B. P. Padhy, P. K. Das, M. Misra, P. Samanta and U. K. Misra, *Trigonometric Fourier approximation of the conjugate series of a function of generalized Lipschitz class by product summability*, Computational Intelligence in Data Mining **1** (2016), 181–190.
- [3] Deepmala, L. N. Mishra and V. N. Mishra, *Trigonometric Approximation of signals (functions) belonging to the $W(L_r, \xi(t))$, ($r \geq 1$)-class by (E, q) ($q > 0$)-means of the conjugate series of its Fourier series*, GJMS Special Issue for Recent Advances in Mathematical Sciences and Applications, Global Journal of Mathematical Sciences **2**(2) (2014), 61–69.
- [4] F. Moricz, *Enlarged Lipschitz and Zygmund classes of functions and Fourier transforms*, East Journal on Approximations **16**(3) (2010), 259–271.
- [5] F. Moricz and J. Nemeth, *Generalized Zygmund classes of functions and strong approximation by Fourier series*, Acta Sci. Math. **73**(3–4) (2007), 637–648.
- [6] G. H. Hardy, *Divergent Series*, Oxford University Press, Oxford, 1949.
- [7] H. K. Nigam, *On approximation in generalized Zygmund class*, Demonstr. Math. **52**(1) (2019), 370–387.
- [8] L. Leindler, *Strong approximation and generalized Zygmund class*, Acta Sci. Math. **43**(3–4) (1981), 301–309.
- [9] L. N. Mishra, V. N. Mishra, K. Khatri and Deepmala, *On the trigonometric approximation of signals belonging to generalized weighted Lipschitz class $W(L^r, \xi(t))$, ($r \geq 1$)-class by matrix (C^1, N_p) operator of conjugate series of its Fourier series*, Appl. Math. Comput. **237** (2014), 252–263.
- [10] L. N. Mishra, V. N. Mishra and V. Sonavane, *Trigonometric approximation of functions belonging to Lipschitz class by matrix (C^1, N_p) operator of conjugate series of Fourier series*, Adv. Difference Equ. **2013**(1) (2013), Article ID 127.
- [11] M. V. Singh, M. L. Mittal and B. E. Rhoades, *Approximation of functions in the generalized Zygmund class using Hausdroff means*, J. Inequal. Appl. **2017**(1) (2017), Article ID 101.
- [12] S. Lal and Shireen, *Best approximation of functions of generalized Zygmund class by matrix-Euler summability mean of Fourier series*, Bull. Math. Anal. Appl. **5**(4) (2013), 1–13.
- [13] V. N. Mishra, *Some Problems on Approximations of Functions in Banach Spaces*, Ph. D. Thesis, 2007.

- [14] V. N. Mishra, K. Khatri and L. N. Mishra, *Approximation of functions belonging to $Lip(\xi(t), r)$ class by $(N, p_n)(E, q)$ -summability of conjugate series of Fourier series*, J. Inequal. Appl. **2012** (2012), Article ID 296.
- [15] V. N. Mishra, K. Khatri and L. N. Mishra, *Product $(N, p_n)(C, 1)$ -summability of a sequence of a sequence of Fourier coefficients*, Math. Sci. **6** (2012), Article ID 38.
- [16] V. N. Mishra, K. Khatri, L. N. Mishra and Deepmala, *Trigonometric approximation of periodic signals belonging to generalized weighted Lipschitz $W(L_r, \xi(t))$, $(r \geq 1)$ -class by Norlund Euler $(N, p_n)(E, q)$ operator of conjugate series of its Fourier series*, J. Class. Anal. **5**(2) (2014), 91–105.
- [17] V. N. Mishra and L. N. Mishra, *Trigonometric approximation of signals (functions) in L^p , $(p \geq 1)$ -norm*, International Journal of Contemporary Mathematical Sciences **7**(19) (2012), 909–918.
- [18] V. N. Mishra, V. Sonavane and L. N. Mishra, *L_r -Approximation of signals (functions) belonging to weighted $W(L_r, \xi(t))$ -class by (C^1, N_p) summability method of conjugate series of its Fourier series*, J. Inequal. Appl. **2013** (2013), Article ID 440.

¹DEPARTMENT OF MATHEMATICS,
KIIT DEEMED TO BE UNIVERSITY,
BHUBANESWAR-24, ODISHA, INDIA
Email address: m.anvesha17@gmail.com
Email address: birupakhya.padhyfma@kiit.ac.in

²DEPARTMENT OF MATHEMATICS,
SCHOOL OF ADVANCED SCIENCES,
VELLORE INSTITUTE OF TECHNOLOGY (VIT) UNIVERSITY,
VELLORE 632 014, TAMIL NADU, INDIA
Email address: lakshminarayanmishra04@gmail.com

³DEPARTMENT OF MATHEMATICS,
NATIONAL INSTITUTE OF SCIENCE AND TECHNOLOGY,
BERHAMPUR, ODISHA, INDIA
Email address: umakanta_misra@yahoo.com