

## COEFFICIENT ESTIMATES FOR SUBCLASS OF $m$ -FOLD SYMMETRIC BI-UNIVALENT FUNCTIONS

A. MOTAMEDNEZHAD<sup>1</sup>, S. SALEHIAN<sup>2</sup>, AND N. MAGESH<sup>3</sup>

ABSTRACT. In the present paper, a general subclass  $\mathcal{M}_{\Sigma_m}^{h,p}(\lambda, \gamma)$  of the  $m$ -Fold symmetric bi-univalent functions is defined. Also, the estimates of the Taylor-Maclaurin coefficients  $|a_{m+1}|$ ,  $|a_{2m+1}|$  and Fekete-Szegő problems are obtained for functions in this new subclass. The results presented in this paper would generalize and improve some recent works of several earlier authors.

### 1. INTRODUCTION

Let  $\mathcal{A}$  be a class of analytic functions in the open unit disk  $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$  of the form

$$(1.1) \quad f(z) = z + \sum_{n=2}^{\infty} a_n z^n.$$

Denote by  $\mathcal{S}$  the class of all functions in the normalized analytic function class  $\mathcal{A}$  which are univalent in  $\mathbb{U}$  (see details in [2, 3]).

Since univalent functions are one-to-one, they are invertible and the inverse functions need not be defined on the entire unit disk  $\mathbb{U}$ . In fact, the Koebe one-quarter theorem [3] ensures that the image of  $\mathbb{U}$  under every univalent function  $f \in \mathcal{S}$  contains a disk of radius  $1/4$ . Therefore, every function  $f \in \mathcal{S}$  has an inverse  $f^{-1}$  satisfying  $f^{-1}(f(z)) = z$  ( $z \in \mathbb{U}$ ) and

$$f(f^{-1}(w)) = w \quad \left( |w| < r_0(f), r_0(f) \geq \frac{1}{4} \right).$$

---

*Key words and phrases.* Bi-univalent functions,  $m$ -fold symmetric univalent functions,  $m$ -fold symmetric bi-univalent functions, coefficient estimates, Fekete-Szegő problem.

2010 *Mathematics Subject Classification.* Primary: 30C45. Secondary: 30C50.

*Received:* March 30, 2019.

*Accepted:* January 03, 2020.

In fact, the inverse function  $f^{-1}$  is given by

$$(1.2) \quad f^{-1}(w) = w - a_2w^2 + (2a_2^2 - a_3)w^3 - (5a_2^3 - 5a_2a_3 + a_4)w^4 + \dots$$

A function  $f \in \mathcal{A}$  is said to be bi-univalent in  $\mathbb{U}$ , if both  $f$  and  $f^{-1}$  are univalent in  $\mathbb{U}$  (see [10]). We denote  $\sigma_{\mathcal{B}}$  the class of bi-univalent functions in  $\mathbb{U}$  given by (1.1). For examples the functions  $\frac{z}{1-z}$  and  $-\log(1-z)$  belong to the class  $\sigma_{\mathcal{B}}$ .

The first time in 1967, Lewin [4] introduced the class  $\sigma_{\mathcal{B}}$  and proved that the bound for the second coefficients of every  $f \in \sigma_{\mathcal{B}}$  satisfies the inequality  $|a_2| < 1.51$ . Also, Smith [5] showed that  $|a_2| < 2/\sqrt{27}$  and  $|a_3| < 4/27$  for bi-univalent polynomial  $f(z) = z + a_2z^2 + a_3z^3$  with real coefficients.

Recently many researchers introduced subclasses of bi-univalent functions and obtain non-sharp estimates on the first two Taylor-Maclaurin coefficients  $|a_2|$  and  $|a_3|$ . For example, we refer the reader to Srivastava et al. [6, 8, 10] and others [13, 14]. The coefficient estimate problem, i.e., bound of  $|a_n|$  ( $n \in \mathbb{N} - \{2, 3\}$ ) for each  $f \in \sigma_{\mathcal{B}}$ , is still an open problem.

Let  $m$  be a positive integer. A domain  $E$  is known as  $m$ -Fold symmetric if a rotation of  $E$  around origin with an angle  $2\pi/m$  maps  $E$  on itself. A function  $f(z)$  analytic in  $\mathbb{U}$  is said to be  $m$ -Fold symmetric if

$$f\left(e^{i\frac{2\pi}{m}}z\right) = e^{i\frac{2\pi}{m}}f(z).$$

For each function  $f \in \mathcal{S}$ , function

$$(1.3) \quad h(z) = \sqrt[m]{f(z^m)}$$

is univalent and maps unit disk  $\mathbb{U}$  into a region with  $m$ -Fold symmetry.

We denote by  $\mathcal{S}_m$  the class of  $m$ -Fold symmetric univalent functions in  $\mathbb{U}$  and clearly  $\mathcal{S}_1 = \mathcal{S}$ . Every  $f \in \mathcal{S}_m$  has a series expansion of the form

$$(1.4) \quad f(z) = z + \sum_{k=1}^{\infty} a_{mk+1}z^{mk+1} \quad (z \in \mathbb{U}, m \in \mathbb{N}).$$

Srivastava et al. [11], introduced a natural extensions of  $m$ -Fold symmetric univalent functions and defined the class  $\Sigma_m$  of symmetric bi-univalent functions. They obtained the series expansion for  $g = f^{-1}$  as:

$$(1.5) \quad \begin{aligned} f^{-1}(w) = & w - a_{m+1}w^{m+1} + [(m+1)a_{m+1}^2 - a_{2m+1}]w^{2m+1} \\ & - \left[ \frac{1}{2}(m+1)(3m+2)a_{m+1}^3 - (3m+2)a_{m+1}a_{2m+1} + a_{3m+1} \right] w^{3m+1} + \dots \end{aligned}$$

For  $m = 1$  formula (1.5) coincides with formula (1.2) of the class  $\sigma_{\mathcal{B}}$ .

In fact, this widely-cited work by Srivastava et al. [7] actually revived the study of  $m$ -Fold bi-univalent functions in recent years and that it has led to a flood of papers on the subject by (for example) Srivastava et al. [7, 9, 11, 12].

The aim of the this paper is to introduce new subclass  $\mathcal{M}_{\Sigma_m}^{h,p}(\lambda, \gamma)$  of the  $m$ -Fold symmetric bi-univalent functions class  $\Sigma_m$ . Moreover, we obtain estimates on initial coefficients  $|a_{m+1}|$ ,  $|a_{2m+1}|$  and Fekete-Szegő problems for functions in this subclass.

The results presented in this paper would generalize and improve some recent works of Altinkaya et al. [1] and Li et al. [13].

### 2. SUBCLASS $\mathcal{M}_{\Sigma_m}^{h,p}(\lambda, \gamma)$

In this section, we introduce and consider the subclass  $\mathcal{M}_{\Sigma_m}^{h,p}(\lambda, \gamma)$ .

**Definition 2.1.** Assume that  $h : \mathbb{U} \rightarrow \mathbb{C}$  and  $p : \mathbb{U} \rightarrow \mathbb{C}$ , are analytic functions of the form

$$\begin{aligned} h(z) &= 1 + h_m z^m + h_{2m} z^{2m} + h_{3m} z^{3m} + \dots, \\ p(w) &= 1 + p_m w^m + p_{2m} w^{2m} + p_{3m} w^{3m} + \dots, \end{aligned}$$

such that

$$\min\{\operatorname{Re}(h(z)), \operatorname{Re}(p(z))\} > 0 \quad (z \in \mathbb{U}).$$

Let  $\lambda \geq 0$  and  $\gamma \in \mathbb{C} - \{0\}$ . We say that a function  $f$  given by (1.4) is in the subclass  $\mathcal{M}_{\Sigma_m}^{h,p}(\lambda, \gamma)$ , if the following conditions are satisfied:

$$(2.1) \quad 1 + \frac{1}{\gamma} \left[ (1 - \lambda) \frac{z f'(z)}{f(z)} + \lambda \left( 1 + \frac{z f''(z)}{f'(z)} \right) - 1 \right] \in h(\mathbb{U}) \quad (z \in \mathbb{U})$$

and

$$(2.2) \quad 1 + \frac{1}{\gamma} \left[ (1 - \lambda) \frac{w g'(w)}{g(w)} + \lambda \left( 1 + \frac{w g''(w)}{g'(w)} \right) - 1 \right] \in p(\mathbb{U}) \quad (w \in \mathbb{U}),$$

where  $g$  is the extension of  $f^{-1}$  to  $\mathbb{U}$ .

**Definition 2.2.** A function  $f \in \Sigma_m$  given by (1.4) is said to be in the subclass  $\mathcal{C}_{\Sigma_m}(\beta)$  ( $0 \leq \beta < 1$ ), if two following conditions are satisfied:

$$\operatorname{Re} \left( 1 + \frac{z f''(z)}{f'(z)} \right) > \beta \quad \text{and} \quad \operatorname{Re} \left( 1 + \frac{w g''(w)}{g'(w)} \right) > \beta \quad (z, w \in \mathbb{U}),$$

where  $g$  is the extension of  $f^{-1}$  to  $\mathbb{U}$ .

*Remark 2.1.* There are many selections of the functions  $h(z)$  and  $p(z)$  which would provide interesting classes of  $m$ -Fold symmetric bi-univalent functions  $\Sigma_m$ . For example, if we let

$$h(z) = p(z) = \left( \frac{1 + z^m}{1 - z^m} \right)^\alpha = 1 + 2\alpha z^m + 2\alpha^2 z^{2m} + \dots \quad (0 < \alpha \leq 1),$$

it is easy to verify that the functions  $h(z)$  and  $p(z)$  satisfy the hypotheses of Definition 2.1. If  $f \in \mathcal{M}_{\Sigma_m}^{h,p}(\lambda, \gamma)$ , then

$$\left| \arg \left( 1 + \frac{1}{\gamma} \left[ (1 - \lambda) \frac{z f'(z)}{f(z)} + \lambda \left( 1 + \frac{z f''(z)}{f'(z)} \right) - 1 \right] \right) \right| < \frac{\alpha\pi}{2}$$

and

$$\left| \arg \left( 1 + \frac{1}{\gamma} \left[ (1 - \lambda) \frac{w g'(w)}{g(w)} + \lambda \left( 1 + \frac{w g''(w)}{g'(w)} \right) - 1 \right] \right) \right| < \frac{\alpha\pi}{2}.$$

In this case we say that  $f$  belongs to the subclass  $\mathcal{M}_{\Sigma_m}(\alpha, \lambda, \gamma)$ .

Also, for  $h(z) = p(z) = \left(\frac{1+z^m}{1-z^m}\right)^\alpha$ ,  $\gamma = 1$  and  $\lambda = 0$ , the subclass  $\mathcal{M}_{\Sigma_m}^{h,p}(\lambda, \gamma)$  reduces to the subclass  $\mathcal{S}_{\Sigma_m}^\alpha$  which was considered by Altinkaya and Yalcin [1].

If we let

$$h(z) = p(z) = \frac{1 + (1 - 2\beta)z^m}{1 - z^m} = 1 + 2(1 - \beta)z^m + 2(1 - \beta)z^{2m} + \dots \quad (0 \leq \beta < 1),$$

it is easy to verify that the functions  $h(z)$  and  $p(z)$  satisfy the hypotheses of Definition 2.1. If  $f \in \mathcal{M}_{\Sigma_m}^{h,p}(\lambda, \gamma)$ , then

$$\operatorname{Re} \left( 1 + \frac{1}{\gamma} \left[ (1 - \lambda) \frac{zf'(z)}{f(z)} + \lambda \left( 1 + \frac{zf''(z)}{f'(z)} \right) - 1 \right] \right) > \beta$$

and

$$\operatorname{Re} \left( 1 + \frac{1}{\gamma} \left[ (1 - \lambda) \frac{wg'(w)}{g(w)} + \lambda \left( 1 + \frac{wg''(w)}{g'(w)} \right) - 1 \right] \right) > \beta.$$

In this case we say that  $f$  belongs to the subclass  $\mathcal{M}_{\Sigma_m}(\beta, \lambda, \gamma)$ .

Also, for  $h(z) = p(z) = \frac{1+(1-2\beta)z^m}{1-z^m}$ ,  $\gamma = 1$  and  $\lambda = 0$ , the subclass  $\mathcal{M}_{\Sigma_m}^{h,p}(\lambda, \gamma)$  reduces to the subclass  $\mathcal{S}_{\Sigma_m}^\beta$  considered by Altinkaya and Yalcin [1].

Furthermore, for  $h(z) = p(z) = \frac{1+(1-2\beta)z^m}{1-z^m}$ ,  $\gamma = 1$  and  $\lambda = 1$ , the subclass  $\mathcal{M}_{\Sigma_m}^{h,p}(\lambda, \gamma)$  reduces to Definition 2.2.

*Remark 2.2.* For one-fold symmetric bi-univalent functions, we denote the subclass  $\mathcal{M}_{\Sigma_1}^{h,p}(\lambda, \gamma) = \mathcal{M}_{\Sigma}^{h,p}(\lambda, \gamma)$ . Special cases of this subclass are illustrated below.

- (i) By putting  $h(z) = p(z) = \left(\frac{1+z}{1-z}\right)^\alpha$  and  $\gamma = 1$ , the subclass  $\mathcal{M}_{\Sigma}^{h,p}(\lambda, \gamma)$  reduces to the subclass  $M_{\Sigma}(\alpha, \lambda)$  studied by Li and Wang [13].
- (ii) By putting  $h(z) = p(z) = \left(\frac{1+z}{1-z}\right)^\alpha$ ,  $\gamma = 1$  and  $\lambda = 0$ , the subclass  $\mathcal{M}_{\Sigma}^{h,p}(\lambda, \gamma)$  reduces to the subclass  $\mathcal{S}_{\sigma_B}^\alpha$  of strongly bi-starlike functions of order  $\alpha$  ( $0 < \alpha \leq 1$ ).
- (iii) By putting  $h(z) = p(z) = \frac{1+(1-2\beta)z}{1-z}$  and  $\gamma = 1$ , the subclass  $\mathcal{M}_{\Sigma}^{h,p}(\lambda, \gamma)$  reduces to the subclass  $B_{\Sigma}(\beta, \lambda)$  studied by Li and Wang [13].
- (iv) By putting  $h(z) = p(z) = \frac{1+(1-2\beta)z}{1-z}$ ,  $\gamma = 1$  and  $\lambda = 0$ , the subclass  $\mathcal{M}_{\Sigma}^{h,p}(\lambda, \gamma)$  reduces to the subclass  $\mathcal{S}_{\sigma_B}(\beta)$  of bi-starlike functions of order  $\beta$  ( $0 \leq \beta < 1$ ).
- (v) By putting  $h(z) = p(z) = \frac{1+(1-2\beta)z}{1-z}$  and  $\lambda = \gamma = 1$ , the subclass  $\mathcal{M}_{\Sigma}^{h,p}(\lambda, \gamma)$  reduces to the subclass  $\mathcal{C}_{\sigma_B}(\beta)$  of bi-convex functions of order  $\beta$  ( $0 \leq \beta < 1$ ).

**Theorem 2.1.** *Let  $f$  given by (1.4) be in the subclass  $\mathcal{M}_{\Sigma_m}^{h,p}(\lambda, \gamma)$  ( $\lambda \geq 0, \gamma \in \mathbb{C} - \{0\}$ ). Then*

$$|a_{m+1}| \leq \min \left\{ \frac{|\gamma||h_m|}{m(1 + \lambda m)}, \sqrt{\frac{|\gamma|(|h_{2m}| + |p_{2m}|)}{2m^2(1 + \lambda m)}} \right\}$$

and

$$|a_{2m+1}| \leq \min \left\{ \frac{|\gamma|(|h_{2m}| + |p_{2m}|)}{4m(1 + 2\lambda m)} + \frac{(m + 1)|\gamma|^2(|h_m|^2 + |p_m|^2)}{4m^2(1 + \lambda m)^2}, \right. \\ \left. \frac{(3\lambda m^2 + 2\lambda m + 2m + 1)|\gamma||h_{2m}| + (\lambda m^2 + 2\lambda m + 1)|\gamma||p_{2m}|}{4m^2(1 + 2\lambda m)(1 + \lambda m)} \right\}.$$

*Proof.* The main idea in the proof of Theorem 2.1 is to get the desired bounds for the coefficient  $|a_{m+1}|$  and  $|a_{2m+1}|$ . Indeed, by considering the relations (2.1) and (2.2), we have

$$(2.3) \quad 1 + \frac{1}{\gamma} \left[ (1 - \lambda) \frac{zf'(z)}{f(z)} + \lambda \left( 1 + \frac{zf''(z)}{f'(z)} \right) - 1 \right] = h(z) \quad (z \in \mathbb{U})$$

and

$$(2.4) \quad 1 + \frac{1}{\gamma} \left[ (1 - \lambda) \frac{wg'(w)}{g(w)} + \lambda \left( 1 + \frac{wg''(w)}{g'(w)} \right) - 1 \right] = p(w) \quad (w \in \mathbb{U}),$$

where each of the functions  $h$  and  $p$  satisfies the conditions of Definition 2.1. For precise comparison of the coefficients of the above equations, in the following we obtain Taylor-Maclaurin series expansions each side of the equations

$$(2.5) \quad 1 + \frac{1}{\gamma} \left[ (1 - \lambda) \frac{zf'(z)}{f(z)} + \lambda \left( 1 + \frac{zf''(z)}{f'(z)} \right) - 1 \right] \\ = 1 + \frac{m(1 + \lambda m)}{\gamma} a_{m+1} z^m + \left\{ \frac{2m(1 + 2\lambda m)}{\gamma} a_{2m+1} - \frac{m(1 + 2\lambda m + \lambda m^2)}{\gamma} a_{m+1}^2 \right\} z^{2m} \\ + \dots,$$

and

$$(2.6) \quad 1 + \frac{1}{\gamma} \left[ (1 - \lambda) \frac{wg'(w)}{g(w)} + \lambda \left( 1 + \frac{wg''(w)}{g'(w)} \right) - 1 \right] \\ = 1 - \frac{m(1 + \lambda m)}{\gamma} a_{m+1} w^m + \left\{ -\frac{2m(1 + 2\lambda m)}{\gamma} a_{2m+1} \right. \\ \left. + \frac{m(1 + 2m + 2\lambda m + 3\lambda m^2)}{\gamma} a_{m+1}^2 \right\} w^{2m} + \dots.$$

Also from the Definition 2.1, the analytic functions  $h$  and  $p$  have the following Taylor-Maclaurin series expansions

$$(2.7) \quad h(z) = 1 + h_m z^m + h_{2m} z^{2m} + h_{3m} z^{3m} + \dots$$

and

$$(2.8) \quad p(w) = 1 + p_m w^m + p_{2m} w^{2m} + p_{3m} w^{3m} + \dots.$$

By comparing the coefficients of the equations (2.5), (2.7), (2.6) and (2.8), respectively, we get

$$(2.9) \quad \frac{m(1 + \lambda m)}{\gamma} a_{m+1} = h_m,$$

$$(2.10) \quad \frac{2m(1 + 2\lambda m)}{\gamma} a_{2m+1} - \frac{m(1 + 2\lambda m + \lambda m^2)}{\gamma} a_{m+1}^2 = h_{2m},$$

$$(2.11) \quad -\frac{m(1 + \lambda m)}{\gamma} a_{m+1} = p_m$$

and

$$(2.12) \quad -\frac{2m(1 + 2\lambda m)}{\gamma} a_{2m+1} + \frac{m(1 + 2m + 2\lambda m + 3\lambda m^2)}{\gamma} a_{m+1}^2 = p_{2m}.$$

From (2.9) and (2.11), we get

$$(2.13) \quad h_m = -p_m$$

and

$$(2.14) \quad a_{m+1}^2 = \frac{\gamma^2(h_m^2 + p_m^2)}{2m^2(1 + \lambda m)^2}.$$

Adding (2.10) and (2.12), we get

$$(2.15) \quad a_{m+1}^2 = \frac{\gamma(h_{2m} + p_{2m})}{2m^2(1 + \lambda m)}.$$

Therefore, we find from the equations (2.13), (2.14) and (2.15) that

$$|a_{m+1}| \leq \frac{|\gamma||h_m|}{m(1 + \lambda m)} \quad \text{and} \quad |a_{m+1}| \leq \sqrt{\frac{|\gamma|(|h_{2m}| + |p_{2m}|)}{2m^2(1 + \lambda m)}},$$

respectively. So, we get the desired estimate on the coefficient  $|a_{m+1}|$ .

The proof is completed by finding the bound on the coefficient  $|a_{2m+1}|$ . Upon subtracting (2.12) from (2.10), we get

$$(2.16) \quad a_{2m+1} = \frac{\gamma(h_{2m} - p_{2m})}{4m(1 + 2\lambda m)} + \frac{(m + 1)}{2} a_{m+1}^2.$$

Putting the value of  $a_{m+1}^2$  from (2.14) into (2.16), it follows that

$$(2.17) \quad a_{2m+1} = \frac{\gamma(h_{2m} - p_{2m})}{4m(1 + 2\lambda m)} + \frac{(m + 1)\gamma^2(h_m^2 + p_m^2)}{4m^2(1 + \lambda m)^2}.$$

By substituting the value of  $a_{m+1}^2$  from (2.15) into (2.16), we obtain

$$(2.18) \quad a_{2m+1} = \frac{\gamma(h_{2m} - p_{2m})}{4m(1 + 2\lambda m)} + \frac{(m + 1)\gamma(h_{2m} + p_{2m})}{4m^2(1 + \lambda m)}.$$

Therefore, from the equations (2.17) and (2.18), we get

$$|a_{2m+1}| \leq \frac{|\gamma|(|h_{2m}| + |p_{2m}|)}{4m(1 + 2\lambda m)} + \frac{(m + 1)|\gamma|^2(|h_m|^2 + |p_m|^2)}{4m^2(1 + \lambda m)^2}$$

and

$$|a_{2m+1}| \leq \frac{(3\lambda m^2 + 2\lambda m + 2m + 1)|\gamma||h_{2m}| + (\lambda m^2 + 2\lambda m + 1)|\gamma||p_{2m}|}{4m^2(1 + 2\lambda m)(1 + \lambda m)}. \quad \square$$

**Theorem 2.2.** *Let  $f$  given by (1.4) be in the subclass  $\mathcal{M}_{\Sigma_m}^{h,p}(\lambda, \gamma)$  ( $\lambda \geq 0, \gamma \in \mathbb{C} - \{0\}$ ). Also let  $\rho$  be real number. Then*

$$|a_{2m+1} - \rho a_{m+1}^2| \leq \begin{cases} \frac{|\gamma|}{4m(1 + 2\lambda m)} \{(1 + T(\rho))|h_{2m}| + (1 - T(\rho))|p_{2m}|\}, & |T(\rho)| \leq 1, \\ \frac{|\gamma|}{4m(1 + 2\lambda m)} \{|1 + T(\rho)||h_{2m}| + |T(\rho) - 1||p_{2m}|\}, & |T(\rho)| \geq 1, \end{cases}$$

where

$$T(\rho) = \frac{(m - 2\rho + 1)(1 + 2\lambda m)}{m(1 + \lambda m)}.$$

*Proof.* From the equation (2.16), we get

$$(2.19) \quad a_{2m+1} - \rho a_{m+1}^2 = \frac{\gamma(h_{2m} - p_{2m})}{4m(1 + 2\lambda m)} + \frac{m - 2\rho + 1}{2} a_{m+1}^2.$$

From the equation (2.15) and (2.19), we have

$$a_{2m+1} - \rho a_{m+1}^2 = \frac{|\gamma|}{4m(1 + 2\lambda m)} \left\{ \left[ 1 + \frac{(m - 2\rho + 1)(1 + 2\lambda m)}{m(1 + \lambda m)} \right] h_{2m} + \left[ \frac{(m - 2\rho + 1)(1 + 2\lambda m)}{m(1 + \lambda m)} - 1 \right] p_{2m} \right\}.$$

Next, taking the absolute values we obtain

$$|a_{2m+1} - \rho a_{m+1}^2| \leq \frac{|\gamma|}{4m(1 + 2\lambda m)} \left\{ \left| 1 + \frac{(m - 2\rho + 1)(1 + 2\lambda m)}{m(1 + \lambda m)} \right| |h_{2m}| + \left| \frac{(m - 2\rho + 1)(1 + 2\lambda m)}{m(1 + \lambda m)} - 1 \right| |p_{2m}| \right\}.$$

Then, we conclude that

$$|a_{2m+1} - \rho a_{m+1}^2| \leq \begin{cases} \frac{|\gamma|}{4m(1 + 2\lambda m)} \{(1 + T(\rho))|h_{2m}| + (1 - T(\rho))|p_{2m}|\}, & |T(\rho)| \leq 1, \\ \frac{|\gamma|}{4m(1 + 2\lambda m)} \{|1 + T(\rho)||h_{2m}| + |T(\rho) - 1||p_{2m}|\}, & |T(\rho)| \geq 1. \end{cases}$$

□

## 3. COROLLARIES AND CONSEQUENCES

By setting

$$h(z) = p(z) = \left( \frac{1+z^m}{1-z^m} \right)^\alpha = 1 + 2\alpha z^m + 2\alpha^2 z^{2m} + \dots \quad (0 < \alpha \leq 1, z \in \mathbb{U}),$$

in Theorem 2.1, we conclude the following result.

**Corollary 3.1.** *Let  $f$  given by (1.4) be in the subclass  $\mathcal{M}_{\Sigma_m}(\alpha, \lambda, \gamma)$  ( $0 < \alpha \leq 1$ ,  $\lambda \geq 0$ ,  $\gamma \in \mathbb{C} - \{0\}$ ). Then*

$$|a_{m+1}| \leq \min \left\{ \frac{2\alpha|\gamma|}{m(1+\lambda m)}, \frac{\alpha}{m} \sqrt{\frac{2|\gamma|}{1+\lambda m}} \right\}$$

and

$$|a_{2m+1}| \leq \min \left\{ \frac{\alpha^2|\gamma|}{m(1+2\lambda m)} + \frac{2\alpha^2(m+1)|\gamma|^2}{m^2(1+\lambda m)^2}, \frac{\alpha^2|\gamma|(m+1)}{m^2(1+\lambda m)} \right\}.$$

By setting  $h(z) = p(z) = \left( \frac{1+z^m}{1-z^m} \right)^\alpha$  ( $0 < \alpha \leq 1$ ) in Theorem 2.2, we conclude the following result.

**Corollary 3.2.** *Let  $f$  given by (1.4) be in the subclass  $\mathcal{M}_{\Sigma_m}(\alpha, \lambda, \gamma)$  ( $0 < \alpha \leq 1$ ,  $\lambda \geq 0$ ,  $\gamma \in \mathbb{C} - \{0\}$ ). Also let  $\rho$  be real number. Then*

$$|a_{2m+1} - \rho a_{m+1}^2| \leq \begin{cases} \frac{\alpha^2|\gamma|}{m(1+2\lambda m)}, & |T(\rho)| \leq 1, \\ \frac{\alpha^2|T(\rho)||\gamma|}{m(1+2\lambda m)}, & |T(\rho)| \geq 1, \end{cases}$$

where

$$T(\rho) = \frac{(m-2\rho+1)(1+2\lambda m)}{m(1+\lambda m)}.$$

By setting  $\gamma = 1$  and  $\lambda = 0$  in Corollary 3.1, we conclude the following result.

**Corollary 3.3.** *Let  $f$  given by (1.4) be in the subclass  $\mathcal{S}_{\Sigma_m}^\alpha$  ( $0 < \alpha \leq 1$ ). Then*

$$|a_{m+1}| \leq \frac{\sqrt{2}\alpha}{m}$$

and

$$|a_{2m+1}| \leq \min \left\{ \frac{\alpha^2}{m} + \frac{2(m+1)\alpha^2}{m^2}, \frac{(m+1)\alpha^2}{m^2} \right\} = \frac{(m+1)\alpha^2}{m^2}.$$

*Remark 3.1.* The bounds on  $|a_{m+1}|$  and  $|a_{2m+1}|$  given in Corollary 3.3 are better than those given in [1, Corollary 6], because of

$$\frac{\sqrt{2}\alpha}{m} \leq \frac{2\alpha}{m\sqrt{\alpha+1}}$$

and

$$\frac{(m+1)\alpha^2}{m^2} \leq \frac{\alpha^2}{m} + \frac{2(m+1)\alpha^2}{m^2} \leq \frac{\alpha}{m} + \frac{2(m+1)\alpha^2}{m^2}.$$



By setting  $m = 1$  and  $\gamma = 1$  in Corollary 3.1, we conclude the following result.

**Corollary 3.4.** *Let  $f$  given by (1.1) be in the subclass  $M_{\Sigma}(\alpha, \lambda)$  ( $0 < \alpha \leq 1$ ,  $\lambda \geq 0$ ). Then*

$$|a_2| \leq \begin{cases} \alpha \sqrt{\frac{2}{1+\lambda}}, & 0 \leq \lambda \leq 1, \\ \frac{2\alpha}{1+\lambda}, & \lambda \geq 1, \end{cases}$$

and

$$|a_3| \leq \begin{cases} \frac{2\alpha^2}{1+\lambda}, & 0 \leq \lambda \leq \frac{2+\sqrt{13}}{3}, \\ \frac{\alpha^2}{1+2\lambda} + \frac{4\alpha^2}{(1+\lambda)^2}, & \lambda \geq \frac{2+\sqrt{13}}{3}. \end{cases}$$

*Remark 3.2.* The bounds on  $|a_2|$  and  $|a_3|$  given in Corollary 3.4 are better than those given in [13, Theorem 2.2].

By setting  $m = 1$  in Corollary 3.3, we conclude the following result.

**Corollary 3.5.** *Let  $f$  given by (1.1) be in the subclass  $\mathcal{S}_{\sigma_B}^{\alpha}$  of strongly bi-starlike functions of order  $\alpha$  ( $0 < \alpha \leq 1$ ). Then*

$$|a_2| \leq \sqrt{2}\alpha \quad \text{and} \quad |a_3| \leq 2\alpha^2.$$

By setting

$$\begin{aligned} h(z) = p(z) &= \frac{1 + (1 - 2\beta)z^m}{1 - z^m} \\ &= 1 + 2(1 - \beta)z^m + 2(1 - \beta)z^{2m} + \cdots \quad (0 \leq \beta < 1, z \in \mathbb{U}), \end{aligned}$$

in Theorem 2.1, we conclude the following result.

**Corollary 3.6.** *Let  $f$  given by (1.4) be in the subclass  $\mathcal{M}_{\Sigma_m}(\beta, \lambda, \gamma)$  ( $0 \leq \beta < 1$ ,  $\lambda \geq 0$ ,  $\gamma \in \mathbb{C} - \{0\}$ ). Then*

$$|a_{m+1}| \leq \min \left\{ \frac{2(1-\beta)|\gamma|}{m(1+\lambda m)}, \sqrt{\frac{2(1-\beta)|\gamma|}{m^2(1+\lambda m)}} \right\}$$

and

$$|a_{2m+1}| \leq \min \left\{ \frac{(1-\beta)|\gamma|}{m(1+2\lambda m)} + \frac{2(1-\beta)^2(m+1)|\gamma|^2}{m^2(1+\lambda m)^2}, \frac{(1-\beta)(m+1)|\gamma|}{m^2(1+\lambda m)} \right\}.$$

By setting  $h(z) = p(z) = \frac{1+(1-2\beta)z^m}{1-z^m}$  ( $0 \leq \beta < 1$ ) in Theorem 2.2, we conclude the following result.

**Corollary 3.7.** *Let  $f$  given by (1.4) be in the subclass  $\mathcal{M}_{\Sigma_m}(\beta, \lambda, \gamma)$  ( $0 \leq \beta < 1$ ,  $\lambda \geq 0$ ,  $\gamma \in \mathbb{C} - \{0\}$ ). Also let  $\rho$  be real number. Then*

$$|a_{2m+1} - \rho a_{m+1}^2| \leq \begin{cases} \frac{(1 - \beta)|\gamma|}{m(1 + 2\lambda m)}, & |T(\rho)| \leq 1, \\ \frac{(1 - \beta)|\gamma||T(\rho)|}{m(1 + 2\lambda m)}, & |T(\rho)| \geq 1, \end{cases}$$

where

$$T(\rho) = \frac{(m - 2\rho + 1)(1 + 2\lambda m)}{m(1 + \lambda m)}.$$

By setting  $\gamma = 1$  and  $\lambda = 0$  in Corollary 3.6, we conclude the following result.

**Corollary 3.8.** *Let  $f$  given by (1.4) be in the subclass  $\mathcal{S}_{\Sigma_m}^\beta$  ( $0 \leq \beta < 1$ ). Then*

$$|a_{m+1}| \leq \begin{cases} \frac{\sqrt{2(1 - \beta)}}{m}, & 0 \leq \beta \leq \frac{1}{2}, \\ \frac{2(1 - \beta)}{m}, & \frac{1}{2} \leq \beta < 1, \end{cases}$$

and

$$|a_{2m+1}| \leq \begin{cases} \frac{(m + 1)(1 - \beta)}{m^2}, & 0 \leq \beta \leq \frac{1 + 2m}{2(1 + m)}, \\ \frac{2(m + 1)(1 - \beta)^2}{m^2} + \frac{1 - \beta}{m}, & \frac{1 + 2m}{2(1 + m)} \leq \beta < 1. \end{cases}$$

*Remark 3.3.* The bounds on  $|a_{m+1}|$  and  $|a_{2m+1}|$  given in Corollary 3.8 are better than those given in [1, Corolary 7].

By setting  $\gamma = 1$  and  $\lambda = 1$  in Corollary 3.6, we conclude the following result.

**Corollary 3.9.** *Let  $f$  given by (1.4) be in the subclass  $\mathcal{C}_{\Sigma_m}(\beta)$  ( $0 \leq \beta < 1$ ). Then*

$$|a_{m+1}| \leq \begin{cases} \frac{1}{m} \sqrt{\frac{2(1-\beta)}{(1+m)}}, & 2\beta + m \leq 1, \\ \frac{2(1 - \beta)}{m(1 + m)}, & 2\beta + m \geq 1, \end{cases}$$

and

$$|a_{2m+1}| \leq \begin{cases} \frac{1 - \beta}{m^2}, & 0 \leq \beta \leq \frac{1 + 2m - m^2}{2(1 + 2m)}, \\ \frac{1 - \beta}{m(1 + 2m)} + \frac{2(1 - \beta)^2}{m^2(1 + m)}, & \frac{1 + 2m - m^2}{2(1 + 2m)} \leq \beta < 1. \end{cases}$$

By setting  $m = 1$  and  $\gamma = 1$  in Corollary 3.6, we conclude the following result.

**Corollary 3.10.** *Let  $f$  given by (1.1) be in the subclass  $B_{\Sigma}(\beta, \lambda)$  ( $0 \leq \beta < 1$ ,  $\lambda \geq 0$ ). Then*

$$|a_2| \leq \begin{cases} \sqrt{\frac{2(1-\beta)}{1+\lambda}}, & \lambda + 2\beta \leq 1, \\ \frac{2(1-\beta)}{1+\lambda}, & \lambda + 2\beta \geq 1, \end{cases}$$

and

$$|a_3| \leq \begin{cases} \frac{2(1-\beta)}{1+\lambda}, & 0 \leq \beta \leq \frac{3+4\lambda-3\lambda^2}{4(1+2\lambda)}, \\ \frac{1-\beta}{1+2\lambda} + \frac{4(1-\beta)^2}{(1+\lambda)^2}, & \frac{3+4\lambda-3\lambda^2}{4(1+2\lambda)} \leq \beta < 1. \end{cases}$$

*Remark 3.4.* The bounds on  $|a_2|$  and  $|a_3|$  given in Corollary 3.10 are better than those given in [13, Theorem 3.2].

By setting  $m = 1$  in Corollary 3.8, we conclude the following result.

**Corollary 3.11.** *Let  $f$  given by (1.1) be in the subclass  $\mathcal{S}_{\sigma_B}(\beta)$  of bi-starlike functions of order  $\beta$  ( $0 \leq \beta < 1$ ). Then*

$$|a_2| \leq \begin{cases} \sqrt{2(1-\beta)}, & 0 \leq \beta \leq \frac{1}{2}, \\ 2(1-\beta), & \frac{1}{2} \leq \beta < 1, \end{cases}$$

and

$$|a_3| \leq \begin{cases} 2(1-\beta), & 0 \leq \beta \leq \frac{3}{4}, \\ 4(1-\beta)^2 + (1-\beta), & \frac{3}{4} \leq \beta < 1. \end{cases}$$

By setting  $m = 1$  in Corollary 3.9, we conclude the following result.

**Corollary 3.12.** *Let  $f$  given by (1.1) be in the subclass  $\mathcal{C}_{\sigma_B}(\beta)$  of bi-convex functions of order  $\beta$  ( $0 \leq \beta < 1$ ). Then*

$$|a_2| \leq 1 - \beta \quad \text{and} \quad |a_3| \leq \begin{cases} 1 - \beta, & 0 \leq \beta \leq \frac{1}{3}, \\ \frac{1-\beta}{3} + (1-\beta)^2, & \frac{1}{3} \leq \beta < 1. \end{cases}$$

#### REFERENCES

- [1] Ş. Altinkaya and S. Yalçın, *Coefficient bounds for certain subclasses of  $m$ -fold symmetric bi-univalent functions*, J. Math. **2015** (2015), Article ID 241683, 5 pages.
- [2] D. Breaz, N. Breaz and H. M. Sirvastava, *An extention of the univalent conditions for a family of integral operators*, Appl. Math. Lett. **22** (2009), 41–44.
- [3] P. L. Duren, *Univalent Functions*, Grundlehren der Mathematischen Wissenschaften, Band 259, Springer-Verlag, New York, Berlin, Heidelberg. Tokyo, 1983.

- [4] M. Lewin, *On a coefficient problem for bi-univalent functions*, Proc. Amer. Math. Soc. **18** (1967), 63–68.
- [5] H. V. Smith, *Bi-univalent polynomials*, Simon Stevin **50**(2) (1976/77), 115–122.
- [6] H. M. Srivastava, S. Bulut, M. Çağlar and N. Yağmur, *Coefficient estimates for a general subclass of analytic and bi-univalent functions*, Filomat **27**(5) (2013), 831–842.
- [7] H. M. Srivastava, S. Gaboury and F. Ghanim, *Initial coefficient estimates for some subclasses of  $m$ -fold symmetric bi-univalent functions*, Acta Math. Sci. Ser. B **36**(3) (2016), 863–871.
- [8] H. M. Srivastava, S. Gaboury and F. Ghanim, *Coefficient estimates for some general subclasses of analytic and bi-univalent functions*, Afr. Mat. **28** (2017), 693–706.
- [9] H. M. Srivastava, S. Gaboury and F. Ghanim, *Coefficient estimates for some subclasses of  $m$ -fold symmetric bi-univalent functions*, Acta Univ. Apulensis Math. Inform. **41** (2015), 153–164.
- [10] H. M. Srivastava, A. K. Mishra and P. Gochhayat, *Certain subclasses of analytic and bi-univalent functions*, Appl. Math. Lett. **23** (2010), 1188–1192.
- [11] H. M. Srivastava, S. Sivasubramanian and R. Sivakumar, *Initial coefficient bounds for a subclass of  $m$ -fold symmetric bi-univalent functions*, Tbilisi Math. J. **7**(2) (2014), 1–10.
- [12] H. M. Srivastava, A. Zireh and S. Hajiparvaneh, *Coefficient estimates for some subclasses of  $m$ -fold symmetric bi-univalent functions*, Filomat **32**(9) (2018), 3143–3153.
- [13] X. -F. Li and A. -P. Wang, *Two new subclasses of bi-univalent functions*, International Mathematical Forum **7**(30) (2012), 1495–1504.
- [14] A. Zireh and S. Salehian, *On the certain subclass of analytic and bi-univalent functions defined by convolution*, Acta Univ. Apulensis Math. Inform. **44** (2015), 9–19.

<sup>1</sup>FACULTY OF MATHEMATICAL SCIENCES,  
SHAHROOD UNIVERSITY OF TECHNOLOGY,  
P. O. BOX 316-36155, SHAHROOD, IRAN  
Email address: a.motamedne@gmail.com

<sup>2</sup>DEPARTMENT OF MATHEMATICS,  
KORDKUY CENTER, GORGAN BRANCH,  
ISLAMIC AZAD UNIVERSITY, KORDKUY, IRAN  
Email address: s.salehian84@gmail.com

<sup>3</sup>POST-GRADUATE AND RESEARCH DEPARTMENT OF MATHEMATICS  
GOVERNMENT ARTS COLLEGE FOR MEN,  
KRISHNAGIRI 635001, TAMILNADU, INDIA  
Email address: nmagi\_2000@yahoo.co.in