

## BEURLING'S THEOREM FOR THE $Q$ -FOURIER-DUNKL TRANSFORM

EL MEHDI LOUALID<sup>1\*</sup>, AZZEDINE ACHAK<sup>1</sup>, AND RADOUAN DAHER<sup>1</sup>

ABSTRACT. The  $Q$ -Fourier-Dunkl transform satisfies some uncertainty principles in a similar way to the Euclidean Fourier transform. By using the heat kernel associated to the  $Q$ -Fourier-Dunkl operator, we establish an analogue of Beurling's theorem for the  $Q$ -Fourier-Dunkl transform  $\mathcal{F}_Q$  on  $\mathbb{R}$ .

### 1. INTRODUCTION AND PRELIMINARIES

There are many known theorems which state that a function and its classical Fourier transform on  $\mathbb{R}$  cannot both be sharply localized. That is, it is impossible for a nonzero function and its Fourier transform to be simultaneously small. This principle has several versions which were proved by A. Beurling [3]. The Beurling theorem for the classical Fourier transform on  $\mathbb{R}$  which was proved by L. Hörmander [5], says that for any non-trivial function  $f$  in  $L^2(\mathbb{R})$ , the function  $f(x)\mathcal{F}(y)$  is never integrable on  $\mathbb{R}^2$  with respect to the measure  $e^{|xy|}dxdy$ . A far-reaching generalization of this result has been recently proved in [4]. In this paper the author proved that a square-integrable function  $f$  on  $\mathbb{R}$  satisfying for an integer  $N$

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \frac{|f(x)||\mathcal{F}(y)|}{(1+|x|+|y|)^N} e^{|xy|} dxdy < \infty,$$

has the form  $f(x) = P(x)e^{-rx^2}$ , where  $P$  is a polynomial of degree strictly lower than  $\frac{N-1}{2}$  and  $r > 0$ . Many authors have established the analogue of Beurling's theorem in other various settings of harmonic analysis (see for instance [1, 6]). In this paper we study an analogue of Beurling's theorem, in the next we deduce an analogue of Gelfand-Shilov, for the  $Q$ -Fourier-Dunkl transform.

---

*Key words and phrases.*  $Q$ -Fourier-Dunkl transform, Beurling's theorem, uncertainty principles.  
2010 *Mathematics Subject Classification.* Primary: 42A38. Secondary: 44A35, 34B30.

*Received:* August 29, 2017.

*Accepted:* September 03, 2018.

The outline of the content of this paper is as follows. Section 2 is dedicated to some properties and results concerning the Q-Fourier-Dunkl transform. In Section 3 we give an analogue of Beurling's theorem and Gelfand-Shilov theorems for the Q-Fourier-Dunkl transform. Let us now be more precise and describe our results. To do so, we need to introduce some notations. Throughout this paper  $\alpha > -\frac{1}{2}$ ,

- $Q(x) = \exp(-\int_0^x q(t)dt)$ ,  $x \in \mathbb{R}$ , where  $q$  is a  $\mathcal{C}^\infty$  real-valued odd function on  $\mathbb{R}$ ;
- $L_{\alpha}^p(\mathbb{R})$  the class of measurable functions  $f$  on  $\mathbb{R}$  for which  $\|f\|_{p,\alpha} < \infty$ , where

$$\|f\|_{p,\alpha} = \left( \int_{\mathbb{R}} |f(x)|^p |x|^{2\alpha+1} dx \right)^{\frac{1}{p}}, \quad \text{if } p < \infty,$$

and  $\|f\|_{\infty,\alpha} = \|f\|_{\infty} = \text{esssup}_{x \in \mathbb{R}} |f(x)|$ .

- $L_Q^p(\mathbb{R})$  the class of measurable functions  $f$  on  $\mathbb{R}$  for which  $\|f\|_{p,Q} = \|Qf\|_{p,\alpha} < \infty$ , where  $Q$  is given by  $Q(x) = \exp(-\int_0^x q(t)dt)$ ,  $x \in \mathbb{R}$ .

We consider the first singular differential-difference operator  $\Lambda$  defined on  $\mathbb{R}$

$$\Lambda f(x) = f'(x) + \left( \alpha + \frac{1}{2} \right) \frac{f(x) - f(-x)}{x} + q(x)f(x),$$

where  $q$  is a  $\mathcal{C}^\infty$  real-valued odd function on  $\mathbb{R}$ . For  $q = 0$  we regain the Dunkl operator  $\Lambda_\alpha$  associated with reflection group  $\mathbb{Z}_2$  on  $\mathbb{R}$  given by

$$\Lambda_\alpha f(x) = f'(x) + \left( \alpha + \frac{1}{2} \right) \frac{f(x) - f(-x)}{x}.$$

**1.1. Q-Fourier-Dunkl Transform.** The following statements are proved in [2].

**Lemma 1.1.** (a) For each  $\lambda \in \mathbb{C}$ , the differential-difference equation

$$\Lambda u = i\lambda u, \quad u(0) = 1,$$

admits a unique  $\mathcal{C}^\infty$  solution on  $\mathbb{R}$ , denoted by  $\Psi_\lambda$ , given by

$$\Psi_\lambda(x) = Q(x)e_\alpha(i\lambda x),$$

where  $e_\alpha$  denotes the one-dimensional Dunkl kernel defined by

$$e_\alpha(z) = j_\alpha(iz) + \frac{z}{2(\alpha+1)} j_{\alpha+1}(z), \quad z \in \mathbb{C},$$

and  $j_\alpha$  being the normalized spherical Bessel function of index  $\alpha$  given by

$$j_\alpha(z) = \Gamma(\alpha+1) \sum_{n=0}^{\infty} \frac{(-1)^n \left(\frac{z}{2}\right)^{2n}}{n! \Gamma(n+\alpha+1)}, \quad z \in \mathbb{C}.$$

(b) For all  $x \in \mathbb{R}$ ,  $\lambda \in \mathbb{C}$  and  $n = 0, 1, \dots$ , we have

$$\left| \frac{\partial^n}{\partial \lambda^n} \Psi_\lambda(x) \right| \leq Q(x) |x|^n e^{|\text{Im}(\lambda)| |x|}.$$

In particular,

$$|\Psi_\lambda(x)| \leq Q(x) e^{|\text{Im}(\lambda)| |x|}.$$

(c) For all  $x \in \mathbb{R}$ ,  $\lambda \in \mathbb{C}$ , we have the Laplace type integral representation

$$\Psi_\lambda(x) = a_\alpha Q(x) \int_{-1}^1 (1-t^2)^{\alpha-\frac{1}{2}} (1+t) e^{i\lambda xt} dt,$$

$$\text{where } a_\alpha = \frac{\Gamma(\alpha+1)}{\sqrt{\pi}\Gamma(\alpha+\frac{1}{2})}.$$

**Definition 1.1.** The Q-Fourier-Dunkl transform associated with  $\Lambda$  for a function in  $L^1_Q(\mathbb{R})$  is defined by

$$\mathcal{F}_Q(f)(\lambda) = \int_{\mathbb{R}} f(x) \Psi_{-\lambda}(x) |x|^{2\alpha+1} dx.$$

**Theorem 1.1.** (a) Let  $f \in L^1_Q(\mathbb{R})$  such that  $\mathcal{F}_Q(f) \in L^1_\alpha(\mathbb{R})$ . Then for almost  $x \in \mathbb{R}$  we have the inversion formula

$$f(x) (Q(x))^2 = m_\alpha \int_{\mathbb{R}} \mathcal{F}_Q(f)(\lambda) \Psi_\lambda(x) |\lambda|^{2\alpha+1} d\lambda,$$

where

$$m_\alpha = \frac{1}{2^{2(\alpha+1)} (\Gamma(\alpha+1))^2}.$$

(b) For every  $f \in L^2_Q(\mathbb{R})$ , we have the Plancherel formula

$$\int_{\mathbb{R}} |f(x)|^2 (Q(x))^2 |x|^{2\alpha+1} dx = m_\alpha \int_{\mathbb{R}} |\mathcal{F}_Q(f)(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda.$$

(c) The Q-Fourier-Dunkl transform  $\mathcal{F}_Q$  extends uniquely to an isomorphism from  $L^2_Q(\mathbb{R})$  onto  $L^2_\alpha(\mathbb{R})$ .

The heat kernel  $N(x, s)$ ,  $x \in \mathbb{R}$ ,  $s > 0$ , associated with the Q-Fourier-Dunkl transform is given by

$$N(x, s) = m_\alpha \frac{e^{-\frac{x^2}{4s}}}{(2s)^{\alpha+\frac{1}{2}} Q(x)}.$$

Some basic properties of  $N(x, s)$  are the following:

- $N(x, s) Q^2(x) = m_\alpha \int_{\mathbb{R}} e^{-sy^2} \Psi_y(x) |y|^{2\alpha+1} dy;$
- $\mathcal{F}_Q(N(\cdot, s))(x) = e^{-sx^2}.$

We define the heat functions  $W_l$ ,  $l \in \mathbb{N}$ , as

$$(1.1) \quad Q^2(x) W_l(x, s) = \int_{\mathbb{R}} y^l e^{-\frac{y^2}{4s}} \Psi_y(x) |y|^{2\alpha+1} dy,$$

$$(1.2) \quad \mathcal{F}_Q(W_l(\cdot, s)) = i^l y^l e^{-sy^2}.$$

The intertwining operators associated with a Q-Fourier-Dunkl transform on the real line is given by

$$X_Q(f)(x) = a_\alpha Q(x) \int_{-1}^1 f(tx) (1-t^2)^{\alpha-\frac{1}{2}} dt,$$

its dual is given by

$$(1.3) \quad {}^t X_Q(f)(y) = a_\alpha \int_{|x| \geq |y|} f(x) Q(x) \operatorname{sgn}(x) (x^2 - y^2)^{\alpha-\frac{1}{2}} (x+y) dx.$$

**Proposition 1.1.** *If  $f \in L^1_Q(\mathbb{R})$ , then  ${}^tX_Q(f) \in L^1(\mathbb{R})$  and  $\|{}^tX_Q(f)\|_1 \leq \|f\|_{1,Q}$ .*

For every  $f \in L^1_Q(\mathbb{R})$  we have

$$(1.4) \quad \mathcal{F}_Q = \mathcal{F} \circ {}^tX_Q(f),$$

where  $\mathcal{F}$  is the usual Fourier transform defined by

$$\mathcal{F}(f)(\lambda) = \int_{\mathbb{R}} f(x) e^{-i\lambda x} dx.$$

## 2. BEURLING'S THEOREM FOR THE Q-FOURIER-DUNKL TRANSFORM

**Theorem 2.1.** *Let  $N \in \mathbb{N}$  and  $f \in L^2_Q(\mathbb{R})$  satisfy*

$$(2.1) \quad \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{|f(x)| |\mathcal{F}_Q(f)(y)| Q(x)}{(1 + |x| + |y|)^N} e^{|x||y|} |x|^{2\alpha+1} dx dy < \infty.$$

*If  $N > 1$ , then  $f(y) = \sum_{|s| < \frac{N-1}{2}} b_s W_s(r, y)$  a.e. where  $r > 0$ ,  $b_s \in \mathbb{C}$  and  $W_s(r, \cdot)$  is given by (1.1). Otherwise,  $f(y) = 0$  a.e.*

*Proof.* We start with the following lemma.

**Lemma 2.1.** *We suppose that  $f \in L^2_Q(\mathbb{R})$  satisfies (2.1). Then  $f \in L^1_Q(\mathbb{R})$ .*

*Proof.* We may suppose that  $f \neq 0$  in  $L^2_Q(\mathbb{R})$ . (2.1) and Fubini theorem imply that for almost every  $y \in \mathbb{R}$ ,

$$\frac{|\mathcal{F}_Q(f)(y)|}{(1 + |y|)^N} \int_{\mathbb{R}} \frac{Q(x) |f(x)|}{(1 + |x|)^N} e^{|x||y|} |x|^{2\alpha+1} dx < \infty.$$

Since  $\mathcal{F}_Q(f) \neq 0$ , there exist  $y_0 \in \mathbb{R}$ ,  $y_0 \neq 0$ , such that  $\mathcal{F}_Q(f)(y_0) \neq 0$ .

Therefore,

$$(2.2) \quad \int_{\mathbb{R}} \frac{Q(x) |f(x)|}{(1 + |x|)^N} e^{|x||y_0|} |x|^{2\alpha+1} dx < \infty.$$

Since  $\frac{e^{|x||y_0|}}{(1 + |x|)^N} \geq 1$  for large  $|x|$ , it follows that  $\int_{\mathbb{R}} Q(x) |f(x)| |x|^{2\alpha+1} dx < \infty$ .  $\square$

This Lemma and Proposition 1.1 imply that  ${}^tX_Q(f)$  is well-defined almost everywhere on  $\mathbb{R}$ . We shall prove that we have

$$(2.3) \quad \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{|{}^tX_Q(f)(x)| |\mathcal{F}({}^tX_Q(f))(y)|}{(1 + |x| + |y|)^N} e^{|x||y|} dx dy < \infty.$$

Take  $y_0$  as in Lemma 2.1, we write the above integral as a sum of the following integrals

$$I = \int_{\mathbb{R}} \int_{|y| \leq |y_0|} \frac{e^{|x||y|}}{(1 + |x| + |y|)^N} |{}^tX_Q f(x)| |\mathcal{F}({}^tX_Q(f))(y)| dy dx$$

and

$$J = \int_{\mathbb{R}} \int_{|y| \geq |y_0|} \frac{e^{|x||y|}}{(1 + |x| + |y|)^N} |{}^tX_Q f(x)| |\mathcal{F}({}^tX_Q(f))(y)| dy dx.$$

We will prove that I and J are finite, which implies (2.3).

- As the functions  $|\mathcal{F}_Q(f)(y)|$  is continuous in the compact  $\{y \in \mathbb{R} \mid |y| \leq |y_0|\}$ , so we get

$$I \leq C \int_{\mathbb{R}} \frac{e^{|x||y_0|^t} |X_Q f(x)|}{(1+|x|)^N} dx.$$

Writing the integral of the second member as  $I_1 + I_2$  with

$$I_1 = \int_{|x| \leq \frac{N}{|y_0|}} \frac{e^{|x||y_0|^t} |X_Q f(x)|}{(1+|x|)^N} dx$$

and

$$I_2 = \int_{|x| \geq \frac{N}{|y_0|}} \frac{e^{|x||y_0|^t} |X_Q f(x)|}{(1+|x|)^N} dx.$$

Therefore, we have the following results.

- As the function  $x \rightarrow \frac{e^{|x||y_0|^t}}{(1+|x|)^N}$  is continuous in the compact  $\{x \in \mathbb{R} \mid |x| \leq \frac{N}{|y_0|}\}$ , and  $f \in L^1_Q(\mathbb{R})$ , we deduce by using proposition (1.1) that  $|{}^t X_Q(f)|$  belongs to  $L^1(\mathbb{R})$ . Hence,  $I_1$  is finite.
- On the other hand, for  $t > \frac{N}{|y_0|}$ , the function  $t \mapsto \frac{e^{t|y_0|}}{(1+t)^N}$  is increasing, so we obtain by using Proposition 1.1 that

$$I_2 \leq \int_{\mathbb{R}} \frac{Q(\xi) e^{|\xi||y_0|}}{(1+|\xi|)^N} |f(\xi)| |\xi|^{2\alpha+1} d\xi.$$

The inequality (2.2) assert that  $I_2$  is finite. This proves that  $I$  is finite.

- We suppose  $|y_0| \leq N$ . Then  $J = J_1 + J_2 + J_3$ , with

$$J_1 = \int_{|x| \leq \frac{N}{|y_0|}} \int_{|y_0| \leq |y| \leq N} \frac{e^{|x||y|}}{(1+|x|+|y|)^N} |{}^t X_Q(f)(x)| |\mathcal{F}_Q(f)(y)| dy dx,$$

$$J_2 = \int_{|x| \geq \frac{N}{|y_0|}} \int_{|y_0| \leq |y| \leq N} \frac{e^{|x||y|}}{(1+|x|+|y|)^N} |{}^t X_Q(f)(x)| |\mathcal{F}_Q(f)(y)| dy dx,$$

$$J_3 = \int_{\mathbb{R}} \int_{|y| \geq N} \frac{e^{|x||y|}}{(1+|x|+|y|)^N} |{}^t X_Q(f)(x)| |\mathcal{F}_Q(f)(y)| dy dx.$$

- As the function  $(x, y) \mapsto \frac{e^{|x||y|}}{(1+|x|+|y|)^N} |\mathcal{F}_Q(f)(y)|$  is bounded in the compact  $\{x \in \mathbb{R} \mid |x| \leq \frac{N}{|y_0|}\} \times \{y \in \mathbb{R} \mid |y_0| \leq |y| \leq N\}$  and  ${}^t X_Q(|f|)(x)$  is Lebesgue-integrable on  $\mathbb{R}$ , then  $J_1$  is finite.
- Let  $\lambda > 0$ . As the function  $t \mapsto \frac{e^{\lambda t}}{(1+t+\lambda)^N}$  is increasing for  $t > \frac{N}{\lambda}$ . Thus, for all  $(x, y) \in C(\xi, y_0, N)$  we have the inequality

$$\frac{e^{|x||y|}}{(1+|x|+|y|)^N} \leq \frac{e^{|\xi||y|}}{(1+|\xi|+|y|)^N},$$

with

$$C(\xi, y_0, N) = \left\{ (x, y) \in \mathbb{R} \times \mathbb{R} \mid \frac{N}{|y_0|} \leq |x| \leq |\xi| \text{ et } |y_0| \leq |y| \leq N \right\}.$$

Therefore, from Fubini-Tonelli's theorem and Proposition 1.1 we get

$$J_2 \leq \int_{\mathbb{R}} \int_{\mathbb{R}} |Q(\xi)f(\xi)||\mathcal{F}_Q(f)(y)| \frac{e^{|\xi||y|}}{(1+|\xi|+|y|)^N} |\xi|^{2\alpha+1} d\xi dy.$$

Taking account of the condition (2.1), we deduce that  $J_2$  is finite.

- For  $|y| > N$ , the function  $t \mapsto \frac{e^{t|y|}}{(1+t+|y|)^N}$  is increasing. We deduce, by using Fubini-Tonelli's theorem and Proposition 1.1, that

$$J_3 \leq \int_{\mathbb{R}} \int_{|y| \geq N} |(f)(\xi)||F_Q(f)(y)| \frac{e^{|\xi||y|}}{(1+|\xi|+|y|)^N} dy |\xi|^{2\alpha+1} d\xi < +\infty.$$

This implies that  $J_3$  is finite. Finally for  $|y_0| > N$ , we have  $J \leq J_3 < \infty$ . This completes the proof of the relation (2.3).

According to Corollary 3.1, ii) of [4], we conclude that

$${}^t X_Q(f)(x) = R(x)e^{-\delta x^2}, \quad \text{for all } x \in \mathbb{R},$$

with  $\delta > 0$  and  $R$  a polynomial of degree strictly lower than  $\frac{N-1}{2}$ .

Using this relation and (1.4), we deduce that

$$\mathcal{F}_Q(f)(y) = \mathcal{F} \circ {}^t X_Q(f)(y) = \mathcal{F}(R(x)e^{-\delta x^2})(y), \quad \text{for all } x \in \mathbb{R},$$

but

$$\mathcal{F}(P(x)e^{-\delta x^2})(y) = S(y)e^{-\frac{y^2}{4\delta}}, \quad \text{for all } x \in \mathbb{R},$$

with  $S$  a polynomial of degree strictly lower than  $\frac{N-1}{2}$ .

Thus from (1.2) we obtain

$$\mathcal{F}_Q(f)(y) = \mathcal{F}_Q \left( \sum_{|s| < \frac{N-1}{2}} b_s W_s \left( \frac{1}{4\delta}, \cdot \right) \right) (y), \quad \text{for all } x \in \mathbb{R}.$$

The injectivity of the transform  $\mathcal{F}_Q$  implies

$$f(x) = \sum_{|s| < \frac{N-1}{2}} b_s W_s \left( \frac{1}{4\delta}, \cdot \right) (x) \text{ a.e.}, \quad \text{for all } x \in \mathbb{R},$$

and the theorem is proved. □

As an application of Beurling's Theorem, we can deduce a Gelfand-Shilov type theorem for the Q-Fourier-Dunkl transform.

**Theorem 2.2.** *Let  $N \in \mathbb{N}$ ,  $a, b > 0$  and  $1 < p, q < \infty$ , with  $\frac{1}{p} + \frac{1}{q} = 1$  and let  $f \in L_Q^2(\mathbb{R})$  satisfy*

$$(2.4) \quad \int_{\mathbb{R}} \frac{Q(x)|f(x)|e^{\frac{(2a)^p}{p}|x|^p}}{(1+|x|)^N} |x|^{2\alpha+1} dx < \infty$$

and

$$(2.5) \quad \int_{\mathbb{R}} \frac{|\mathcal{F}_Q(f)(y)| e^{\frac{(2b)^q}{q}|y|^q}}{(1+|y|)^N} dy < \infty.$$

If  $ab > \frac{1}{4}$  or  $(p, q) \neq (2, 2)$ , then  $f(x) = 0$  a.e. If  $ab = \frac{1}{4}$  and  $(p, q) = (2, 2)$ , then  $f(x) = \sum_{|s| < \frac{N-1}{2}} b_s W_s(r, \cdot)(x)$ , whenever  $N > 1$  and  $r = 2b^2$ . Otherwise,  $f(x) = 0$  a.e.

*Proof.* Since

$$4ab|x||y| \leq \frac{(2a)^p}{p}|x|^p + \frac{(2b)^q}{q}|y|^q,$$

it follows from (2.4) and (2.5) that

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \frac{Q(x)|f(x)||\mathcal{F}_Q(f)(y)|}{(1+|x|+|y|)^{2N}} e^{4ab|x||y|} |x|^{2\alpha+1} dx dy < \infty.$$

Then (2.1) is satisfied, because  $4ab \leq 1$ . Especially, according to the proof of Theorem 2.1, we can deduce that

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \frac{|{}^t X_Q(f)(x)||\mathcal{F}_Q(f)(y)|}{(1+|x|+|y|)^{2N}} e^{4ab|x||y|} dx dy < \infty,$$

and  ${}^t X_Q(f)$  and  $f$  are of the forms  ${}^t X_Q(f) = R(x)e^{-\frac{x^2}{4r}}$  and  $\mathcal{F}_Q(f)(y) = S(y)e^{-ry^2}$ , where  $r > 0$  and  $S, R$  are polynomials of the same degree strictly lower than  $\frac{2N-1}{2}$ . Therefore, substituting these, we can deduce that

$$(2.6) \quad \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{e^{-(\sqrt{r}|y| - \frac{1}{2\sqrt{r}}|x|)^2} e^{(4ab-1)|x||y|} R(x)S(y)}{(1+|x|+|y|)^{2N}} e^{4ab|x||y|} dx dy < \infty.$$

When  $4ab > 1$ , this integral is not finite unless  $f = 0$  almost everywhere. Indeed, as  $ab > \frac{1}{4}$ , there exists  $\varepsilon > 0$  such that  $4ab - 1 - \varepsilon > 0$ . If  $R$  is non null,  $S$  is also non null and we have

$$\begin{aligned} & \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{|R(x)||S(y)|}{(1+|x|+|y|)^{2N}} e^{-(\sqrt{r}|y| - \frac{1}{2\sqrt{r}}|x|)^2} e^{(4ab-1)|x||y|} dx dy \\ & \geq C \int_{\mathbb{R}} \int_{\mathbb{R}} e^{-(\sqrt{r}|y| - \frac{1}{2\sqrt{r}}|x|)^2} e^{(4ab-1-\varepsilon)|x||y|} dx dy, \end{aligned}$$

where  $C$  is a positive constant. But the function

$$e^{-(\sqrt{r}|y| - \frac{1}{2\sqrt{r}}|x|)^2} e^{(4ab-1-\varepsilon)|x||y|}$$

is not integrable, (2.6) does not hold. Hence,  $f(x) = 0$  a.e.

Moreover, it follows from (2.4) and (2.5) that

$$(2.7) \quad \int_{\mathbb{R}} \frac{|f(x)|Q(x)e^{\frac{(2a)^p}{p}|x|^p}}{(1+|x|)^N} |x|^{2\alpha+1} dx = \int_{\mathbb{R}} \frac{e^{-\frac{1}{4}x^2} e^{\frac{(2a)^p}{p}|x|^p} R(x)Q(x)}{(1+|x|)^N} |x|^{2\alpha+1} dx < \infty$$

and

$$(2.8) \quad \int_{\mathbb{R}} \frac{|\mathcal{F}_Q(f)(y)| e^{\frac{(2b)^q}{q}|y|^q}}{(1+|y|)^N} dy = \int_{\mathbb{R}} \frac{e^{-ry^2} e^{\frac{(2b)^q}{q}|y|^q} S(y)}{(1+|y|)^N} dy < \infty.$$

Hence, one of these integrals is not finite unless  $(p, q) = (2, 2)$ . When  $4ab = 1$  and  $(p, q) = (2, 2)$ , the finiteness of above integrals implies that  $r = 2b^2$  and the rest follows from Theorem 2.1.  $\square$

**Acknowledgements.** The authors thank the reviewers for their valuable comments and suggestions.

#### REFERENCES

- [1] A. Achak, R. Daher and H. Lahlali, *Beurling's theorem for Bessel-Struve transform*, C. R. Math. **354**(1) (2016), 81–85.
- [3] A. Beurling, *The Collected Works of Arne Beurling*, Birkhauser, Boston, 1989.
- [4] A. Bonami, B. Demange and P. Jaming, *Hermite functions and uncertainty principles for the Fourier and the windowed Fourier transforms*, Rev. Mat. Iberoamericana **19** (2002), 22–35.
- [5] L. Hörmander, *A uniqueness theorem of Beurling for Fourier transform pairs*, Ark. Mat. **2**(2) (1991), 237–240.
- [6] S. Negzaoui, *Beurling-Hörmander's theorem related to Bessel-Struve transform*, Integral Transforms Spec. Funct. **27**(9) (2016), 685–697.
- [2] E. A. Al Zahrani, H. El Mir and M. A. Mourou, *Intertwining operators associated with a Dunkl-type operator on the real line and applications*, Far East Journal of Applied Mathematics and Applications **64**(2) (2012), 129–144.

LABORATORY: TOPOLOGY, ALGEBRA, GEOMETRY AND DISCRETE STRUCTURES,  
 DEPARTMENT OF MATHEMATICS AND INFORMATICS,  
 FACULTY OF SCIENCES AÏN CHOCK,  
 UNIVERSITY OF HASSAN II,  
 B.P 5366 MAARIF, CASABLANCA, MOROCCO  
*Email address:* mehdi.loualid@gmail.com  
*Email address:* achakachak@hotmail.fr  
*Email address:* r.daher@fsac.ac.ma

\*CORRESPONDING AUTHOR