

A NEW COMPANION OF OSTROWSKI'S INEQUALITY AND ITS APPLICATIONS

HUANG HONG¹

ABSTRACT. In this paper, we establish a new companion of Ostrowski type inequality for differentiable functions whose first derivatives are bounded, and give its applications to probability density function.

1. INTRODUCTION

In 1938, Ostrowski established the following inequality which can be used to estimate the absolute deviation of a function from its integral mean (see [2]).

Theorem 1.1. *Let $f : I \rightarrow \mathbb{R}$ be a differentiable mapping in the interior I of I , where $I \subset \mathbb{R}$ is an interval, and let $a, b \in I$, $a < b$. If $|f'(t)| \leq M, t \in [a, b]$, then we have*

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[\frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right] (b-a)M, \quad x \in [a, b].$$

The following result is well known in literature as Grüss's inequality ([4], p. 296).

Theorem 1.2. [4] *Let $f, g : [a, b] \rightarrow \mathbb{R}$ be two integrable functions such that $\phi \leq f(x) \leq \Phi$ and $\gamma \leq g(x) \leq \Gamma$, for all $x \in (a, b)$. Then we have*

$$\left| \frac{1}{b-a} \int_a^b f(x)g(x)dx - \frac{1}{(b-a)^2} \int_a^b f(x)dx \int_a^b g(x)dx \right| \leq \frac{1}{4}(\Phi - \phi)(\Gamma - \gamma)$$

and the inequality is sharp in the sense that the constant $\frac{1}{4}$ can not be replaced by a smaller one.

Key words and phrases. Ostrowski's inequality, bounded functions, probability density functions.
2010 *Mathematics Subject Classification.* Primary: 26D10, 26A15, 26A16. Secondary: 26A51.
Received: June 06, 2017.
Accepted: January 11, 2018.

For differentiable mappings whose first derivatives are bounded, Alomari proved two companion inequalities.

Theorem 1.3. [1] *Let $f : I \rightarrow R$ be a differentiable mapping in the interior I of I , where $I \subset R$ is an interval, and let $a, b \in I$, $a < b$. If $f' \in L^1[a, b]$ and $\gamma \leq f'(t) \leq \Gamma$, for all $x \in [a, b]$, then we have*

$$(1.1) \quad \left| \frac{f(x) + f(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{\Gamma - \gamma}{2(b-a)} \left[(x-a)^2 + \left(\frac{a+b}{2} - x \right)^2 \right],$$

for all $x \in [a, \frac{a+b}{2}]$.

Theorem 1.4. [5] *Let the assumptions of Theorem 1.3 hold. Then we have*

$$(1.2) \quad \left| \frac{f(x) + f(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{8}(b-a)(\Gamma - \gamma),$$

for all $x \in [a, \frac{a+b}{2}]$.

We remark that the bound of (1.1) is better than that of (1.2). In fact, for $x \in [a, \frac{a+b}{2}]$, we have

$$\begin{aligned} \frac{1}{8}(b-a) &= \frac{1}{2} \frac{(x-a + \frac{a+b}{2} - x)^2}{b-a} \\ &\geq \frac{1}{2} \frac{(x-a)^2 + \left(\frac{a+b}{2} - x \right)^2}{b-a}. \end{aligned}$$

In this paper, using a different mean, we at first give (1.1) a new proof, and point out that (1.1) is sharp in the sense that the constant $\frac{1}{2}$ cannot be replaced by a smaller one. Secondly, we establish a new companion inequality of Ostrowski type for differentiable mappings whose first derivatives are bounded and consider its applications to probability density function.

2. MAIN RESULTS

A new proof of Theorem 1.3.

Proof. We define a mapping as

$$k(x, t) = \begin{cases} t - a, & t \in [a, x], \\ t - \frac{a+b}{2}, & (x, a+b-x], \\ t - b, & (a+b-x, b], \end{cases}$$

for all $x \in [a, \frac{a+b}{2}]$.

Integrating by parts, we have

$$(2.1) \quad \frac{1}{b-a} \int_a^b k(x, t) f'(t) dt = \frac{f(x) + f(a+b-x)}{2} - \frac{1}{b-a} \int_a^b f(t) dt.$$

If $t \in [a, b]$ and $x \in [a, \frac{a+b}{2}]$, then from the definition of $k(x, t)$, we have

$$p(x, t) \geq 0, \quad t \in [a, x] \cup \left(\frac{a+b}{2}, a+b-x \right),$$

$$p(x, t) \leq 0, \quad t \in \left[x, \frac{a+b}{2} \right] \cup (a+b-x, b].$$

Hence,

$$(2.2) \quad \begin{aligned} & \frac{1}{b-a} \int_a^b p(x, t) f'(t) dt \\ &= \frac{1}{b-a} \left[\int_a^x (t-a) f'(t) dt + \int_x^{\frac{a+b}{2}} \left(t - \frac{a+b}{2} \right) f'(t) dt \right. \\ & \quad \left. + \int_{\frac{a+b}{2}}^{a+b-x} \left(t - \frac{a+b}{2} \right) f'(t) dt + \int_{a+b-x}^b (t-b) f'(t) dt \right] \\ &\leq \frac{\Gamma}{b-a} \left[\int_a^x (t-a) dt + \int_{\frac{a+b}{2}}^{a+b-x} \left(t - \frac{a+b}{2} \right) dt \right] \\ & \quad + \frac{\gamma}{b-a} \left[\int_x^{\frac{a+b}{2}} \left(t - \frac{a+b}{2} \right) dt + \int_{a+b-x}^b (t-b) dt \right] \\ &= \frac{\Gamma}{2(b-a)} \left[(x-a)^2 + \left(\frac{a+b}{2} - x \right)^2 \right] \\ & \quad - \frac{\gamma}{2(b-a)} \left[(x-a)^2 + \left(\frac{a+b}{2} - x \right)^2 \right] \\ &= \frac{\Gamma - \gamma}{2(b-a)} \left[(x-a)^2 + \left(\frac{a+b}{2} - x \right)^2 \right]. \end{aligned}$$

A similar argument can prove that

$$(2.3) \quad -\frac{1}{b-a} \int_a^b p(x, t) f'(x) \leq \frac{\Gamma - \gamma}{2(b-a)} \left[(x-a)^2 + \left(\frac{a+b}{2} - x \right)^2 \right].$$

Form (2.1)–(2.3), we get the inequality (1.1).

Form the proof, let

$$f(x) = \begin{cases} \Gamma(t-a), & t \in [a, x], \\ \Gamma(x-a) + \gamma(t-x), & \left(x, \frac{a+b}{2} \right], \\ \Gamma \left(t+x - \frac{3a+b}{2} \right) + \gamma \left(\frac{a+b}{2} - x \right), & \left(\frac{a+b}{2}, a+b-x \right], \\ \Gamma \frac{b-a}{2} + \gamma \left(t - \frac{a+b}{2} \right), & (a+b-x, b], \end{cases}$$

then the inequality (1.1) holds equality. Thus the constant $1/2$ is sharp in the sense that it cannot be replaced by a smaller one. \square

Theorem 2.1. *Let $f : I \rightarrow R$ be a differentiable mapping in the interior I of I , where $I \subset R$ is an interval, and let $a, b \in I$, $a < b$. If $f' \in L^1[a, b]$ and $\gamma \leq f'(t) \leq \Gamma$, for all $t \in [a, b]$, then we have*

$$(2.4) \quad \left| \frac{1}{4} \left(f(x) + f(a+b-x) + f\left(\frac{a+x}{2}\right) + f\left(\frac{a+2b-x}{2}\right) \right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ \leq \begin{cases} \frac{1}{2(b-a)} \left(\left(\frac{x-a}{2}\right)^2 + \frac{5}{4} \left(\frac{a+b}{2} - x\right)^2 - \left(\frac{3a+b}{4} - x\right)^2 \right) (\Gamma - \gamma), & a \leq x \leq \frac{3a+b}{4}, \\ \frac{1}{2(b-a)} \left(\left(\frac{x-a}{2}\right)^2 + \frac{5}{4} \left(\frac{a+b}{2} - x\right)^2 + \left(\frac{3a+b}{4} - x\right)^2 \right) (\Gamma - \gamma), & \frac{3a+b}{4} \leq x \leq \frac{a+b}{2}, \end{cases}$$

for all $x \in [a, \frac{a+b}{2}]$.

Proof. We define the function given by

$$k(x, t) = \begin{cases} t - a, & t \in \left[a, \frac{a+x}{2} \right], \\ t - \frac{3a+b}{4}, & \left(\frac{a+x}{2}, x \right], \\ t - \frac{a+b}{2}, & (x, a+b-x], \\ t - \frac{a+3b}{4}, & \left(a+b-x, \frac{a+2b-x}{2} \right], \\ t - b, & \left(\frac{a+2b-x}{2}, b \right] \end{cases}$$

for all $x \in [a, \frac{a+b}{2}]$.

Integrating by parts, we can state that

$$(2.5) \quad \frac{1}{b-a} \int_a^b k(x, t) f'(t) dt \\ = \frac{1}{4} \left(f(x) + f(a+b-x) + f\left(\frac{a+x}{2}\right) + f\left(\frac{a+2b-x}{2}\right) \right) - \frac{1}{b-a} \int_a^b f(t) dt.$$

We also have

$$(2.6) \quad \frac{1}{b-a} \int_a^b k(x, t) dt = 0.$$

Let $C = \frac{\Gamma+\gamma}{2}$. From (2.5) and (2.6), we have

$$(2.7) \quad \frac{1}{b-a} \int_a^b k(x, t) (f'(t) - C) dt \\ = \frac{1}{4} \left(f(x) + f(a+b-x) + f\left(\frac{a+x}{2}\right) + f\left(\frac{a+2b-x}{2}\right) \right) - \frac{1}{b-a} \int_a^b f(t) dt.$$

Hence, we have

$$(2.8) \quad \left| \frac{1}{b-a} \int_a^b k(x,t)(f'(t) - C)dt \right| \leq \max_{t \in [a,b]} |f'(t) - C| \frac{1}{b-a} \int_a^b |k(x,t)|dt.$$

Noting

$$(2.9) \quad \max_{t \in [a,b]} |f'(t) - C| \leq \frac{\Gamma - \gamma}{2}.$$

If $x \in [a, \frac{3a+b}{4}]$, we have

$$(2.10) \quad \int_a^b |k(x,t)|dt = \int_a^{\frac{a+x}{2}} (t-a)dt + \int_{\frac{a+b}{2}}^{a+b-x} \left(t - \frac{a+b}{2}\right) dt + \int_{a+b-x}^{\frac{a+2b-x}{2}} \left(t - \frac{a+3b}{4}\right) dt + \int_{\frac{a+x}{2}}^x \left(\frac{3a+b}{4} - t\right) dt + \int_x^{\frac{a+b}{2}} \left(\frac{a+b}{2} - t\right) dt + \int_{\frac{a+2b-x}{2}}^b (b-t)dt = \left(\left(\frac{x-a}{2}\right)^2 + \frac{5}{4} \left(\frac{a+b}{2} - x\right)^2 - \left(\frac{3a+b}{4} - x\right)^2 \right).$$

If $x \in [\frac{3a+b}{4}, \frac{a+b}{2}]$, we have

$$(2.11) \quad \int_a^b |k(x,t)| dt = \int_a^{\frac{a+x}{2}} (t-a)dt + \int_{\frac{a+b}{2}}^{a+b-x} \left(t - \frac{a+b}{2}\right) dt + \int_{\frac{3a+b}{4}}^x \left(t - \frac{3a+b}{4}\right) dt + \int_{a+b-x}^{\frac{a+3b}{4}} \left(\frac{a+3b}{4} - t\right) dt + \int_{\frac{3a+b}{4}}^{\frac{a+2b-x}{2}} \left(t - \frac{a+3b}{4}\right) dt + \int_x^{\frac{a+b}{2}} \left(\frac{a+b}{2} - t\right) dt + \int_{\frac{a+2b-x}{2}}^b (b-t)dt + \int_{\frac{a+x}{2}}^{\frac{a+3b}{4}} \left(\frac{3a+b}{4} - t\right) dt = \left(\left(\frac{x-a}{2}\right)^2 + \frac{5}{4} \left(\frac{a+b}{2} - x\right)^2 + \left(\frac{3a+b}{4} - x\right)^2 \right).$$

We have

$$(2.12) \quad \left| \frac{1}{b-a} \int_a^b k(x,t)(f'(t) - C)dt \right| \leq \begin{cases} \frac{1}{2(b-a)} \left(\left(\frac{x-a}{2}\right)^2 + \frac{5}{4} \left(\frac{a+b}{2} - x\right)^2 - \left(\frac{3a+b}{4} - x\right)^2 \right) (\Gamma - \gamma), & a \leq x \leq \frac{3a+b}{4}, \\ \frac{1}{2(b-a)} \left(\left(\frac{x-a}{2}\right)^2 + \frac{5}{4} \left(\frac{a+b}{2} - x\right)^2 + \left(\frac{3a+b}{4} - x\right)^2 \right) (\Gamma - \gamma), & \frac{3a+b}{4} \leq x \leq \frac{a+b}{2}. \end{cases}$$

Form (2.7) and (2.12), we get (2.4). This completes the proof. □

Corollary 2.1. *In inequality (2.4), choosing*

(1) $x = a$, we have the same results of [5]:

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{8} (b-a)(\Gamma - \gamma);$$

(2) $x = \frac{3a+b}{4}$, we have

$$\left| \frac{1}{4} \left[f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right) + f\left(\frac{7a+b}{8}\right) + f\left(\frac{a+7b}{8}\right) \right] - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{3}{64} (b-a)(\Gamma - \gamma);$$

(3) $x = \frac{a+b}{2}$, we have

$$\left| \frac{1}{4} \left[2f\left(\frac{a+b}{2}\right) + f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right) \right] - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{16} (b-a)(\Gamma - \gamma).$$

3. APPLICATIONS TO PROBABILITY DENSITY FUNCTION

Let X be a random variable taking values in the finite interval $[a, b]$, with the probability density function $f : [a, b] \rightarrow [0, 1]$ with the cumulative distribution function $F(x) = Pr(X \leq x) = \int_a^x f(t) dt$.

Theorem 3.1. *With the above assumptions and f being as in Theorem 2.1, we have*

$$(3.1) \quad \left| \frac{1}{4} \left(F(x) + F(a+b-x) + F\left(\frac{a+x}{2}\right) + F\left(\frac{a+2b-x}{2}\right) \right) - \frac{b-E(x)}{b-a} \right| \leq \begin{cases} \frac{1}{2(b-a)} \left(\left(\frac{x-a}{2}\right)^2 + \frac{5}{4} \left(\frac{a+b}{2} - x\right)^2 - \left(\frac{3a+b}{4} - x\right)^2 \right) (\Gamma - \gamma), & a \leq x \leq \frac{3a+b}{4}, \\ \frac{1}{2(b-a)} \left(\left(\frac{x-a}{2}\right)^2 + \frac{5}{4} \left(\frac{a+b}{2} - x\right)^2 + \left(\frac{3a+b}{4} - x\right)^2 \right) (\Gamma - \gamma), & \frac{3a+b}{4} \leq x \leq \frac{a+b}{2}, \end{cases}$$

for all $x \in [a, \frac{a+b}{2}]$, where $E(X)$ is the expectation of X .

Proof. Let $f = F$ in (2.4), and taking into account

$$E(X) = \int_a^b t dF(t) = b - \int_a^b F(t) dt,$$

we obtain (3.1). □

Corollary 3.1. *In Theorem 3.1, putting $x = \frac{3a+b}{4}$ we have*

$$\left| \frac{1}{4} \left[F\left(\frac{3a+b}{4}\right) + F\left(\frac{a+3b}{4}\right) + F\left(\frac{7a+b}{8}\right) + F\left(\frac{a+7b}{8}\right) \right] - \frac{b-E(X)}{b-a} \right| \leq \frac{3(b-a)}{64} (\Gamma - \gamma).$$

Corollary 3.2. *In Theorem 3.1, if $x = \frac{a+b}{2}$, we have*

$$\left| \frac{1}{4} \left[2F \left(\frac{a+b}{2} \right) + F \left(\frac{3a+b}{4} \right) + F \left(\frac{a+3b}{4} \right) \right] - \frac{b-E(X)}{b-a} \right| \leq \frac{(b-a)}{16} (\Gamma - \gamma).$$

Acknowledgements. The author wish to express his heartfelt thanks to the referees for their detailed and helpful suggestions for revising the manuscript.

REFERENCES

- [1] M. W. Alomari, *A companion of Ostrowski's inequality with applications*, Transylvanian Journal of Mathematics and Mechanics **3** (2011), 9–14.
- [2] S. S. Dragomir and Th. M. Rassias (ed.), *Ostrowski Type Inequalities and Applications in Numerical Integration*, Kluwer academic Publishers, Dordrecht, 2002.
- [3] N. S. Barnett, S. S. Dragomir and I. Gomma, *A companion for the Ostrowski and the generalized trapezoid inequalities*, Mathematical and Computer Modeling **50** (2009), 179–187.
- [4] D. S. Mitrinovic, J. E. Pečarić and A. M. Fink, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht, 1993.
- [5] M. W. Alomari, *A companion of Ostrowski's inequality for mappings whose first derivatives are bounded and applications in numerical integration*, Kragujevac J. Math. **36**(1) (2012), 77–82.

¹SCHOOL OF MATHEMATICS AND STATISTICS,
HUBEI ENGINEERING UNIVERSITY,
XIAOGAN, HUBEI, P.R. CHINA
Email address: 2369844949@qq.com