

INTEGRAL INVOLVING THE PRODUCT OF MULTIVARIABLE  
 ALEPH-FUNCTION, GENERAL CLASS OF SRIVASTAVA  
 POLYNOMIALS AND ALEPH-FUNCTION OF ONE VARIABLE

DINESH KUMAR<sup>1</sup>, FRÉDÉRIC AYANT<sup>2,3</sup>, AND NARESH<sup>4</sup>

ABSTRACT. In this paper, we derive an integral involving the multivariable Aleph-function, the general class of Srivastava polynomials, and the Aleph-function of one variable, all of which are sufficiently general in nature and are capable of yielding a large number of simpler and more useful results simply by specialization of their parameters. Moreover, we establish certain specific instances.

1. INTRODUCTION AND PRELIMINARIES

The Aleph ( $\aleph$ )-function was established by Südland et al. [30], but its notation and complete definition are offered below in terms of the Mellin-Barnes type integral (see also, [2, 3, 7, 13, 23, 25]):

$$\begin{aligned} \aleph(z) &= \aleph_{P_i, Q_i, c_i; r'}^{M, N} \left( z \left| \begin{array}{l} (a_j, A_j)_{1, n}, [c_i (a_{ji}, A_{ji})]_{n+1, p_i; r'} \\ (b_j, B_j)_{1, m}, [c_i (b_{ji}, B_{ji})]_{m+1, q_i; r'} \end{array} \right. \right) \\ (1.1) \quad &= \frac{1}{2\pi\omega} \int_L \Omega_{P_i, Q_i, c_i; r'}^{M, N}(s) z^{-s} ds, \end{aligned}$$

for all  $z$  different to 0 and

$$(1.2) \quad \Omega_{P_i, Q_i, c_i; r'}^{M, N}(s) = \frac{\prod_{j=1}^M \Gamma(b_j + B_j s) \prod_{j=1}^N \Gamma(1 - a_j - A_j s)}{\sum_{i=1}^{r'} c_i \left\{ \prod_{j=N+1}^{P_i} \Gamma(a_{ji} + A_{ji} s) \prod_{j=M+1}^{Q_i} \Gamma(1 - b_{ji} - B_{ji} s) \right\}},$$

---

*Key words and phrases.* Aleph-function of several variables, general class of Srivastava polynomials, Aleph-function of one and two variables,  $I$ -function of two and several variables.

2020 *Mathematics Subject Classification.* Primary: 33C99, 33C60. Secondary: 44A20.

DOI

*Received:* August 05, 2022.

*Accepted:* March 17, 2023.

where  $|\arg z| < \frac{1}{2}\pi$  and  $\Omega = \sum_{j=1}^M B_j + \sum_{j=1}^N A_j - c_i \left( \sum_{j=M+1}^{Q_i} B_{ji} + \sum_{j=N+1}^{P_i} A_{ji} \right) > 0$  for  $i = 1, \dots, r'$ . For convergence conditions and other details of Aleph-function (one variable), see Südland et al. [30] (see also, [23, 24]). The series representation of Aleph-function is given by Chaurasia and Singh [6], defined as

$$(1.3) \quad \aleph_{P_i, Q_i, c_i; r'}^{M, N}(z) = \sum_{G=1}^M \sum_{g=0}^{+\infty} \frac{(-1)^g \Omega_{P_i, Q_i, c_i; r'}^{M, N}(s)}{B_G g!} z^{-s},$$

with  $s = \eta_{G, g} = \frac{b_G + g}{B_G}$ ,  $P_i < Q_i$ ,  $|z| < 1$  and  $\Omega_{P_i, Q_i, c_i; r'}^{M, N}(s)$  is given in (1.2).

The generalized polynomials defined by Srivastava [29], and studied by many authors, e.g., [5, 7, 8, 10–12, 14, 18, 20], is given in the following manner:

$$(1.4) \quad \begin{aligned} S_{N_1, \dots, N_s}^{M_1, \dots, M_s}(y_1, \dots, y_s) &= \sum_{K_1=0}^{[N_1/M_1]} \cdots \sum_{K_s=0}^{[N_s/M_s]} \frac{(-N_1)_{M_1 K_1}}{K_1!} \times \cdots \times \frac{(-N_s)_{M_s K_s}}{K_s!} \\ &\times A[N_1, K_1; \dots; N_s, K_s] y_1^{K_1} \cdots y_s^{K_s}, \end{aligned}$$

where  $M_1, \dots, M_s$  are arbitrary positive integers and the coefficients  $A[N_1, K_1; \dots; N_s, K_s]$  are arbitrary constants, real or complex. In the present paper, we use the following notation:

$$(1.5) \quad a_1 = \frac{(-N_1)_{M_1 K_1}}{K_1!} \times \cdots \times \frac{(-N_s)_{M_s K_s}}{K_s!} A[N_1, K_1; \dots; N_s, K_s].$$

The Aleph-function of several variables generalizes the multivariable  $I$ -function defined by Sharma and Ahmad [26], which is a generalization of  $G$  and  $H$ -functions [8, 21] of multiple variables. The multiple Mellin-Barnes integral occurring in this paper will be referred to as the multivariable Aleph-function throughout our present study and will be defined and represented as follows (see also, [4, 15–17, 19]).

$$(1.6) \quad \begin{aligned} &\aleph(z_1, \dots, z_r) \\ &= \aleph_{P_i, Q_i; \tau_i; R; p_i(1), q_i(1), \tau_i(1); R^{(1)}; \dots; p_i(r), q_i(r); \tau_i(r); R^{(r)}}^{0, n; m_1, n_1, \dots, m_r, n_r} \left( \begin{array}{c} z_1 \\ \vdots \\ z_r \end{array} \middle| \begin{array}{c} (a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)})_{1, n} \\ \dots \end{array} \right) \\ &\quad \left[ \tau_i \left( a_{ji}; \alpha_{ji}^{(1)}, \dots, \alpha_{ji}^{(r)} \right) \right]_{n+1, p_i} : (c_j^{(1)}, \gamma_j^{(1)})_{1, n_1}, \\ &\quad \left[ \tau_i \left( b_{ji}; \beta_{ji}^{(1)}, \dots, \beta_{ji}^{(r)} \right) \right]_{1, q_i} : (d_j^{(1)}, \delta_j^{(1)})_{1, m_1}, \\ &\quad \left[ \tau_i(1) \left( c_{ji(1)}^{(1)}, \gamma_{ji(1)}^{(1)} \right) \right]_{n_1+1, p_i(1)} ; \dots ; (c_j^{(r)}, \gamma_j^{(r)})_{1, n_r}, \left[ \tau_i(r) \left( c_{ji(r)}^{(r)}, \gamma_{ji(r)}^{(r)} \right) \right]_{n_r+1, p_i(r)} \\ &\quad \left[ \tau_i(1) \left( d_{ji(1)}^{(1)}, \delta_{ji(1)}^{(1)} \right) \right]_{m_1+1, q_i(1)} ; \dots ; (d_j^{(r)}, \delta_j^{(r)})_{1, m_r}, \left[ \tau_i(r) \left( d_{ji(r)}^{(r)}, \delta_{ji(r)}^{(r)} \right) \right]_{m_r+1, q_i(r)} \end{aligned}$$

$$= \frac{1}{(2\pi\omega)^r} \int_{L_1} \cdots \int_{L_r} \psi(s_1, \dots, s_r) \prod_{k=1}^r \theta_k(s_k) z_k^{s_k} ds_1 \cdots ds_r,$$

with  $\omega = \sqrt{-1}$ . For more details, see Ayant [1]. The real numbers  $\tau_i$  are positives for  $i = 1, \dots, R$ ,  $\tau_{i^{(k)}}$  are positives for  $i^{(k)} = 1, \dots, R^{(k)}$ . The condition for absolute convergence of multiple Mellin-Barnes type contour (1.6) can be obtained by extension of the corresponding conditions for multivariable  $H$ -function given by  $|\arg z_k| < \frac{1}{2}A_i^{(k)}\pi$ , where

$$(1.7) \quad \begin{aligned} A_i^{(k)} &= \sum_{j=1}^n \alpha_j^{(k)} - \tau_i \sum_{j=n+1}^{p_i} \alpha_{ji}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{ji}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} - \tau_{i^{(k)}} \sum_{j=n_k+1}^{p_{i^{(k)}}} \gamma_{ji^{(k)}}^{(k)} + \sum_{j=1}^{m_k} \delta_j^{(k)} \\ &- \tau_{i^{(k)}} \sum_{j=m_k+1}^{q_{i^{(k)}}} \delta_{ji^{(k)}}^{(k)} > 0, \quad \text{with } k = 1, \dots, r, \quad i = 1, \dots, R, \quad i^{(k)} = 1, \dots, R^{(k)}. \end{aligned}$$

The complex numbers  $z_i$  are not zero. Throughout this paper, we assume the existence and absolute convergence conditions of the multivariable Aleph-function. We may establish the asymptotic expansion in the following convenient form:

$$\aleph(z_1, \dots, z_r) = 0 (|z_1|^{\alpha_1}, \dots, |z_r|^{\alpha_r}), \quad \max\{|z_1|, \dots, |z_r|\} \rightarrow 0,$$

$$\aleph(z_1, \dots, z_r) = 0 (|z_1|^{\beta_1}, \dots, |z_r|^{\beta_r}), \quad \min\{|z_1|, \dots, |z_r|\} \rightarrow +\infty,$$

where  $k = 1, \dots, r$ ,  $\alpha_k = \min\{\text{Re}(d_j^{(k)}/\delta_j^{(k)}) : j = 1, \dots, m_k\}$  and  $\beta_k = \max\{\text{Re}((c_j^{(k)} - 1)/\gamma_j^{(k)}) : j = 1, \dots, n_k\}$ . We will use these following notations:

$$(1.8) \quad U = p_i, q_i, \tau_i; R, \quad V = m_1, n_1; \dots; m_r, n_r,$$

$$(1.9) \quad W = p_{i(1)}, q_{i(1)}, \tau_{i(1)}; R^{(1)}, \dots, p_{i(r)}, q_{i(r)}, \tau_{i(r)}; R^{(r)},$$

$$(1.10) \quad A = (a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)})_{1,n}, \quad [\tau_i (a_{ji}; \alpha_{ji}^{(1)}, \dots, \alpha_{ji}^{(r)})]_{n+1, p_i},$$

$$(1.11) \quad B = [\tau_i (b_{ji}; \beta_{ji}^{(1)}, \dots, \beta_{ji}^{(r)})]_{1, q_i},$$

$$(1.12) \quad C = (c_j^{(1)}, \gamma_j^{(1)})_{1, n_1}, \quad [\tau_{i(1)} (c_{ji(1)}^{(1)}, \gamma_{ji(1)}^{(1)})]_{n_1+1, p_{i(1)}}, \dots,$$

$$(1.12) \quad (c_j^{(r)}, \gamma_j^{(r)})_{1, n_r}, \quad [\tau_{i(r)} (c_{ji(r)}^{(r)}, \gamma_{ji(r)}^{(r)})]_{n_r+1, p_{i(r)}},$$

$$(1.13) \quad D = (d_j^{(1)}, \delta_j^{(1)})_{1, m_1}, \quad [\tau_{i(1)} (d_{ji(1)}^{(1)}, \delta_{ji(1)}^{(1)})]_{m_1+1, q_{i(1)}}, \dots,$$

$$(1.13) \quad (d_j^{(r)}, \delta_j^{(r)})_{1, m_r}, \quad [\tau_{i(r)} (d_{ji(r)}^{(r)}, \delta_{ji(r)}^{(r)})]_{m_r+1, q_{i(r)}}.$$

We denote the multivariable Aleph-function as

$$(1.14) \quad \aleph(z_1, \dots, z_r) = \aleph_{U;W}^{0,n;V} \left( \begin{array}{c|c} z_1 & A : C \\ \vdots & B : D \\ z_r & \end{array} \right).$$

We have the following required integral [9]:

$$(1.15) \quad \int_0^{\pi/2} \frac{\sin^{2\alpha-1} \theta \cos^{2\beta-1} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{\alpha+\beta}} d\theta = \frac{1}{2a^{2\beta}b^{2\alpha}} B(\alpha, \beta), \quad \text{Re}(\alpha) > 0, \text{Re}(\beta) > 0,$$

where  $a, b \in \mathbb{C} \setminus \{0\}$  and  $B(\cdot, \cdot)$  is the Beta function.

### 2. MAIN INTEGRAL

In this section we evaluate the integral involving multivariable Aleph-function, a class of polynomials of several variables and a Aleph-function of one variable.

#### Theorem 2.1.

$$\begin{aligned} & \int_0^{\pi/2} \frac{\sin^{2\alpha-1} \theta \cos^{2\beta-1} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{\alpha+\beta}} \aleph_{P_i, Q_i, c_i; r'}^{M, N} \left( t \frac{\sin^{2c} \theta \cos^{2d} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{c+d}} \right) \\ & \times S_{N_1, \dots, N_s}^{M_1, \dots, M_s} \left( \begin{matrix} t_1 \frac{\sin^{2c_1} \theta \cos^{2d_1} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{c_1+d_1}} \\ \vdots \\ t_s \frac{\sin^{2c_s} \theta \cos^{2d_s} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{c_s+d_s}} \end{matrix} \right) \aleph_{U:W}^{0, n; V} \left( \begin{matrix} z_1 \frac{\sin^{2h_1} \theta \cos^{2l_1} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{h_1+l_1}} \\ \vdots \\ z_r \frac{\sin^{2h_r} \theta \cos^{2l_r} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{h_r+l_r}} \end{matrix} \right) d\theta \\ & = \frac{1}{2} \sum_{G=1}^M \sum_{g=0}^{+\infty} \sum_{K_1=0}^{[N_1/M_1]} \cdots \sum_{K_s=0}^{[N_s/M_s]} a_1 \frac{(-1)^g \Omega_{P_i, Q_i, c_i; r'}^{M, N}(\eta_{G, g})}{B_G g!} t^{\eta_{G, g}} t_1^{K_1} \dots t_s^{K_s} \\ & \times a^{-2(\beta+d\eta_{G, g} + \sum_{i=1}^s K_i d_i)} b^{-2(\alpha+c\eta_{G, g} + \sum_{i=1}^s K_i c_i)} \\ & \times \aleph_{U_{21}:W}^{0, n+2; V} \left( \begin{matrix} \frac{z_1}{a^{2l_1} b^{2h_1}} \\ \vdots \\ \frac{z_r}{a^{2l_r} b^{2h_r}} \end{matrix} \middle| \begin{matrix} (1 - \alpha - c\eta_{G, g} - \sum_{i=1}^s K_i c_i; h_1, \dots, h_r), \\ - \\ (1 - \beta - d\eta_{G, g} - \sum_{i=1}^s K_i d_i; l_1, \dots, l_r), A : C \\ (1 - \alpha - \beta - (c + d)\eta_{G, g} - \sum_{i=1}^s K(c_i + d_i); h_1 + l_1, \dots, h_r + l_r), B : D \end{matrix} \right), \end{aligned} \tag{2.1}$$

where  $U_{21} = p_i + 2, q_i + 1, \tau_i, R$ , also satisfy the following conditions:

- (a)  $\min \{c, c_i, h_j\} \leq 0, i = 1, \dots, s; j = 1, \dots, r$  ( $h_j$  are not simultaneously zero);
- (b)  $\min \{d, d_i, l_j\} \leq 0, i = 1, \dots, s; j = 1, \dots, r$  ( $l_j$  are not simultaneously zero);
- (c)  $\text{Re}(\alpha) + c + \sum_{i=1}^r c_i \text{Re}(\alpha) + c \min_{1 \leq l \leq M} \text{Re} \left( \frac{b_j^{(i)}}{\beta_j} \right) + \sum_{i=1}^r h_i \min_{1 \leq j \leq m_i} \text{Re} \left( \frac{d_j^{(i)}}{\delta_j^{(i)}} \right) > 0;$
- (d)  $\text{Re}(\alpha) + d \min_{1 \leq l \leq M} \text{Re} \left( \frac{b_j^{(i)}}{\beta_j} \right) + \sum_{i=1}^r l_i \min_{1 \leq j \leq m_i} \text{Re} \left( \frac{d_j^{(i)}}{\delta_j^{(i)}} \right) > 0;$
- (e)  $|\arg z_k| < \frac{1}{2} A_i^{(k)} \pi$ , where  $A_i^{(k)}$  is given in (1.7);
- (f)  $|\arg t| < \frac{1}{2} \pi \Omega$ , where

$$\Omega = \sum_{j=1}^M \beta_j + \sum_{j=1}^N \alpha_j - c_i \left( \sum_{j=M+1}^{Q_i} \beta_{ji} + \sum_{j=N+1}^{P_i} \alpha_{ji} \right) > 0.$$

*Proof.* Expressing the Aleph-function of one variable in series form with the help of (1.3), the general class of polynomials of several variables in series with the help of (1.4), and the Aleph-function of  $r$  variables in Mellin-Barnes contour integral with the help of (1.6). The conditions (e) and (f) are satisfied, then the integral representing multivariable Aleph function converges uniformly, and we can invert the sums and multiple Mellin-Barnes integrals. Next, by changing the order of integration and summation (which is easily seen to be justified due to the absolute convergence of the integral and summations involved in the process) and then evaluating the resulting integral with the help of equation (1.15). Finally interpreting the result thus obtained with the Mellin-barnes contour integral, we arrive at the desired result (2.1).  $\square$

### 3. MULTIVARIABLE $I$ -FUNCTION

**Corollary 3.1.** *If  $\tau_i, \tau_{i(1)}, \dots, \tau_{i(r)} \rightarrow 1$ , the Aleph-function of several variables renounces to the  $I$ -function of several variables. The simple integral has been derived in this section for multivariable  $I$ -functions defined by Sharma and Ahmad [26]*

$$\begin{aligned}
 & \int_0^{\pi/2} \frac{\sin^{2\alpha-1} \theta \cos^{2\beta-1} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{\alpha+\beta}} \aleph_{P_i, Q_i, c_i, r'}^{M, N} \left( t \frac{\sin^{2c} \theta \cos^{2d} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{c+d}} \right) \\
 & \times S_{N_1, \dots, N_s}^{M_1, \dots, M_s} \left( \begin{matrix} t_1 \frac{\sin^{2c_1} \theta \cos^{2d_1} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{c_1+d_1}} \\ \vdots \\ t_s \frac{\sin^{2c_s} \theta \cos^{2d_s} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{c_s+d_s}} \end{matrix} \right) I_{U:W}^{0, n:V} \left( \begin{matrix} z_1 \frac{\sin^{2h_1} \theta \cos^{2l_1} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{h_1+l_1}} \\ \vdots \\ z_r \frac{\sin^{2h_r} \theta \cos^{2l_r} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{h_r+l_r}} \end{matrix} \right) d\theta \\
 & = \frac{1}{2} \sum_{G=1}^M \sum_{g=0}^{\infty} \sum_{K_1=0}^{[N_1/M_1]} \cdots \sum_{K_s=0}^{[N_s/M_s]} a_1 \frac{(-1)^g \Omega_{P_i, Q_i, c_i, r'}^{M, N}(\eta_{G, g})}{B_G g!} t^{\eta_{G, g}} t_1^{K_1} \dots t_s^{K_s} \\
 & \times a^{-2(\beta + d\eta_{G, g} + \sum_{i=1}^s K_i d_i)} b^{-2(\alpha + c\eta_{G, g} + \sum_{i=1}^s K_i c_i)} \\
 & \times I_{U:21:W}^{0, n+2:V} \left( \begin{matrix} \frac{z_1}{a^{2l_1} b^{2h_1}} \\ \vdots \\ \frac{z_r}{a^{2l_r} b^{2h_r}} \end{matrix} \middle| \begin{matrix} (1 - \alpha - c\eta_{G, g} - \sum_{i=1}^s K_i c_i; h_1, \dots, h_r), \\ - \\ (1 - \beta - d\eta_{G, g} - \sum_{i=1}^s K_i d_i; l_1, \dots, l_r), A' : C' \\ (1 - \alpha - \beta - (c + d)\eta_{G, g} - \sum_{i=1}^s K(c_i + d_i); h_1 + l_1, \dots, h_r + l_r), B' : D' \end{matrix} \right),
 \end{aligned}
 \tag{3.1}$$

where  $A' = (a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)})_{1, n}$ ,  $(a_{ji}; \alpha_{ji}^{(1)}, \dots, \alpha_{ji}^{(r)})_{n+1, p_i}$ ,  $B' = (b_{ji}; \beta_{ji}^{(1)}, \dots, \beta_{ji}^{(r)})_{1, q_i}$ ,  
 $C' = (c_j^{(1)}, \gamma_j^{(1)})_{1, n_1}$ ,  $(c_{ji(1)}^{(1)}, \gamma_{ji(1)}^{(1)})_{n_1+1, p_{i(1)}}$ ;  $\dots$ ;  $(c_j^{(r)}, \gamma_j^{(r)})_{1, n_r}$ ,  $(c_{ji(r)}^{(r)}, \gamma_{ji(r)}^{(r)})_{n_r+1, p_{i(r)}}$ ,  
 $D' = (d_j^{(1)}, \delta_j^{(1)})_{1, m_1}$ ,  $(d_{ji(1)}^{(1)}, \delta_{ji(1)}^{(1)})_{m_1+1, q_{i(1)}}$ ;  $\dots$ ;  $(d_j^{(r)}, \delta_j^{(r)})_{1, m_r}$ ,  $(d_{ji(r)}^{(r)}, \delta_{ji(r)}^{(r)})_{m_r+1, q_{i(r)}}$ ,  
 also under the same conditions that (2.1).

4. ALEPH-FUNCTION OF TWO VARIABLES

**Corollary 4.1.** *If we set  $r = 2$  in (1.6), then we obtain the Aleph-function of two variables defined by Sharma [28] and further generalized by Kumar [13]. We have the following simple integral*

$$\begin{aligned}
 & \int_0^{\pi/2} \frac{\sin^{2\alpha-1} \theta \cos^{2\beta-1} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{\alpha+\beta}} \aleph_{P_i, Q_i, c_i, r'}^{M, N} \left( t \frac{\sin^{2c} \theta \cos^{2d} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{c+d}} \right) \\
 & \times S_{N_1, \dots, N_s}^{M_1, \dots, M_s} \left( \begin{matrix} t_1 \frac{\sin^{2c_1} \theta \cos^{2d_1} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{c_1+d_1}} \\ \vdots \\ t_s \frac{\sin^{2c_s} \theta \cos^{2d_s} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{c_s+d_s}} \end{matrix} \right) \aleph_{U:W}^{0, n:V} \left( \begin{matrix} z_1 \frac{\sin^{2h_1} \theta \cos^{2l_1} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{h_1+l_1}} \\ z_2 \frac{\sin^{2h_2} \theta \cos^{2l_2} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{h_2+l_2}} \end{matrix} \right) d\theta \\
 & = \frac{1}{2} \sum_{G=1}^M \sum_{g=0}^{+\infty} \sum_{K_1=0}^{[N_1/M_1]} \cdots \sum_{K_s=0}^{[N_s/M_s]} a_1 \frac{(-1)^g \Omega_{P_i, Q_i, c_i, r'}^{M, N}(\eta_{G, g})}{B_G g!} t^{\eta_{G, g}} t_1^{K_1} \dots t_s^{K_s} \\
 & \times a^{-2(\beta+d\eta_{G, g} + \sum_{i=1}^s K_s d_i)} b^{-2(\alpha+c\eta_{G, g} + \sum_{i=1}^s K_s c_i)} \\
 & \times \aleph_{U_{21}:W}^{0, n+2:V} \left( \begin{matrix} \frac{z_1}{a^{2l_1} b^{2h_1}} \\ \frac{z_2}{a^{2l_2} b^{2h_2}} \end{matrix} \middle| \begin{matrix} (1 - \alpha - c\eta_{G, g} - \sum_{i=1}^s K_i c_i; h_1, h_2), \\ - \\ (1 - \beta - d\eta_{G, g} - \sum_{i=1}^s K_i d_i; l_1, l_2), A'' : C''; E'' \\ (1 - \alpha - \beta - (c + d)\eta_{G, g} - \sum_{i=1}^s K(c_i + d_i); h_1 + l_1, h_2 + l_2), B'' : D''; F'' \end{matrix} \right),
 \end{aligned}
 \tag{4.1}$$

where  $A'' = (a_j; \alpha'_j, \alpha''_j)_{1, n}$ ,  $[\tau_i(a_{ji}; \alpha'_{ji}, \alpha''_{ji})]_{n+1, p_i}$ ;  $B'' = [\tau_i(b_{ji}; \beta'_{ji}, \beta''_{ji})]_{1, q_i}$ ,  $C'' = (c_j, \gamma_j)_{1, n_1}$ ,  $[\tau_{i'}(c_{ji'}; \gamma_{ji'})]_{n_1+1, p_{i'}}$ ;  $D'' = (d_j, \delta_j)_{1, m_1}$ ,  $[\tau_{i'}(d_{ji'}; \delta_{ji'})]_{m_1+1, q_{i'}}$ ,  $E'' = (e_j, \eta_j)_{1, n_2}$ ,  $[\tau_{i''}(e_{ji''}; \eta_{ji''})]_{n_2+1, p_{i''}}$ ;  $F'' = (f_j, \zeta_j)_{1, m_2}$ ,  $[\tau_{i''}(f_{ji''}; \zeta_{ji''})]_{m_2+1, q_{i''}}$ , also satisfy the existence conditions provided in (2.1).

5. I-FUNCTION OF TWO VARIABLES

**Corollary 5.1.** *If we set  $\tau_i, \tau_{i'}, \tau_{i''} \rightarrow 1$  in (4.1), the Aleph-function of two variables reduces to the I-function of two variables defined by Sharma and Mishra [27], and we obtain the same formula with the I-function of two variables.*

$$\begin{aligned}
 & \int_0^{\pi/2} \frac{\sin^{2\alpha-1} \theta \cos^{2\beta-1} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{\alpha+\beta}} \aleph_{P_i, Q_i, c_i, r'}^{M, N} \left( t \frac{\sin^{2c} \theta \cos^{2d} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{c+d}} \right) \\
 & \times S_{N_1, \dots, N_s}^{M_1, \dots, M_s} \left( \begin{matrix} t_1 \frac{\sin^{2c_1} \theta \cos^{2d_1} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{c_1+d_1}} \\ \vdots \\ t_s \frac{\sin^{2c_s} \theta \cos^{2d_s} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{c_s+d_s}} \end{matrix} \right) I_{U:W}^{0, n:V} \left( \begin{matrix} z_1 \frac{\sin^{2h_1} \theta \cos^{2l_1} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{h_1+l_1}} \\ z_2 \frac{\sin^{2h_2} \theta \cos^{2l_2} \theta}{(a^2 \cos^2 \theta + b^2 \sin^2 \theta)^{h_2+l_2}} \end{matrix} \right) d\theta
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{2} \sum_{G=1}^M \sum_{g=0}^{\infty} \sum_{K_1=0}^{[N_1/M_1]} \cdots \sum_{K_s=0}^{[N_s/M_s]} a_1 \frac{(-1)^g \Omega_{P_i, Q_i, c_i, r'}^{M, N}(\eta_{G, g})}{B_G g!} t^{\eta_{G, g}} t_1^{K_1} \dots t_s^{K_s} \\
 &\times a^{-2(\beta + d\eta_{G, g} + \sum_{i=1}^s K_i d_i)} b^{-2(\alpha + c\eta_{G, g} + \sum_{i=1}^s K_i c_i)} \\
 &\times I_{U_{21}:W}^{0, n+2; V} \left( \begin{array}{c} \frac{z_1}{a^{2l_1} b^{2h_1}} \\ \frac{z_2}{a^{2l_2} b^{2h_2}} \end{array} \middle| \begin{array}{c} (1 - \alpha - c\eta_{G, g} - \sum_{i=1}^s K_i c_i; h_1, h_2), \\ - \\ (1 - \beta - d\eta_{G, g} - \sum_{i=1}^s K_i d_i; l_1, l_2), A''' : C'''; E''' \\ (1 - \alpha - \beta - (c + d)\eta_{G, g} - \sum_{i=1}^s K(c_i + d_i); h_1 + l_1, h_2 + l_2), B''' : D'''; F''' \end{array} \right),
 \end{aligned}
 \tag{5.1}$$

where  $A''' = (a_j; \alpha'_j, \alpha''_j)_{1, n}$ ,  $(a_{ji}; \alpha'_{ji}, \alpha''_{ji})_{n+1, p_i}$ ,  $B''' = (b_{ji}; \beta'_{ji}, \beta''_{ji})_{1, q_i}$ ,  $C''' = (c_j, \gamma_j)_{1, n_1}$ ,  $(c_{j' i'}, \gamma_{j' i'})_{n_1+1, p_{i'}}$ ;  $D''' = (d_j, \delta_j)_{1, m_1}$ ,  $(d_{j' i'}, \delta_{j' i'})_{m_1+1, q_{i'}}$ ,  $E''' = (e_j, \eta_j)_{1, n_2}$ ,  $(e_{j' i'}, \eta_{j' i'})_{n_2+1, p_{i'}}$ ,  $F''' = (f_j, \zeta_j)_{1, m_2}$ ,  $(f_{j' i'}, \zeta_{j' i'})_{m_2+1, q_{i'}}$ , also satisfy the conditions stated in (2.1).

For more details of  $I$ -function of two variables reader can refer to work Kumari et al. [22].

### 6. CONCLUSION

In this work, an integral involving the multivariable Aleph-function, a class of polynomials with several variables (Srivastava polynomials), and an Aleph-function with one variable was evaluated. The integral derived in this study is of a highly broad character, since it incorporates the multivariable Aleph-function, which is a generic function of multiple variables previously explored. Consequently, the integral produced by this study would serve as a key formula from which, by adjusting the parameters, as many outcomes as required involving the special functions of one and multiple variables may be generated.

**Acknowledgements.** The first author (DK) would like to thank Agriculture University Jodhpur for supporting and encouraging this work.

### REFERENCES

- [1] F. Y. Ayant, *An integral associated with the Aleph-functions of several variables*, Int. J. Math. Trends Tech. **31**(3) (2016), 142–154.
- [2] F. Y. Ayant and D. Kumar, *A unified study of Fourier series involving the Aleph-function and the Kampé de Fériet's function*, Int. J. Math. Trends Tech. **35**(1) (2016), 40–48. <https://doi.org/10.14445/22315373/IJMTT/V35P507>
- [3] F. Y. Ayant and D. Kumar, *Certain finite double integrals involving the hypergeometric function and Aleph-function*, Int. J. Math. Trends Tech. **35**(1) (2016), 49–55. <https://doi.org/10.14445/22315373/IJMTT/V35P508>

- [4] F. Y. Ayant and D. Kumar, *Generating relations and multivariable Aleph-function*, Analysis **38**(3) (2018), 137–143. <https://doi.org/10.1515/anly-2017-0054>
- [5] D. Baleanu, D. Kumar and S. D. Purohit, *Generalized fractional integrals of product of two H-functions and a general class of polynomials*, Int. J. Comput. Math. **93**(8) (2016), 1320–1329. <https://doi.org/10.1080/00207160.2015.1045886>
- [6] V. B. L. Chaurasia and Y. Singh, *New generalization of integral equations of fredholm type using Aleph-function*, International Journal of Modern Mathematical Sciences **9**(3) (2014), 208–220.
- [7] J. Choi and D. Kumar, *Certain unified fractional integrals and derivatives for a product of Aleph function and a general class of multivariable polynomials*, J. Inequal. Appl. **2014** (2014), 15 pages. <https://doi.org/10.1186/1029-242X-2014-499>
- [8] J. Daiya, J. Ram and D. Kumar, *The multivariable H-function and the general class of Srivastava polynomials involving the generalized Mellin-Barnes contour integrals*, FILOMAT **30**(6) (2016), 1457–1464. <https://doi.org/10.2298/FIL1606457D>
- [9] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*, Academic Press, 1996.
- [10] R. K. Gupta, B. S. Shaktawat and D. Kumar, *On generalized fractional differentials involving product of two H-functions and a general class of polynomials*, J. Rajasthan Acad. Phys. Sci. **15**(4) (2016), 327–344.
- [11] R. K. Gupta, B. S. Shaktawat and D. Kumar, *Generalized fractional integrals and derivative formulas for  $\overline{H}$ -function and a general class of polynomials*, Journal of Chemical, Biological and Physical Sciences, Section C **6**(3) (2016), 1081–1097.
- [12] D. Kumar, *Fractional calculus formulas involving  $\overline{H}$ -function and Srivastava polynomials*, Commun. Korean Math. Soc. **31**(4) (2016), 827–844. <https://doi.org/10.4134/CKMS.c150251>
- [13] D. Kumar, *Generalized fractional differintegral operators of the Aleph-function of two variables*, Journal of Chemical, Biological and Physical Sciences, Section C **6**(3) (2016), 1116–1131.
- [14] D. Kumar, P. Agarwal and S. D. Purohit, *Generalized fractional integration of the  $\overline{H}$ -function involving general class of polynomials*, Walailak Journal of Science and Technology **11**(12) (2014), 1019–1030. <https://doi.org/10.14456/WJST.2014.57>
- [15] D. Kumar and F. Y. Ayant, *Application of Jacobi polynomial and multivariable Aleph-function in heat conduction in non-homogeneous moving rectangular parallelepiped*, Kragujevac J. Math. **45**(3) (2021), 439–447. <https://doi.org/10.46793/KgJMat2103.439K>
- [16] D. Kumar and F. Y. Ayant, *Fractional calculus operators pertaining to multivariable Aleph-function*, Bulletin of Parana's Mathematical Society **40** (2022), 1–10. <https://doi.org/10.5269/bspm.42491>
- [17] D. Kumar, F. Y. Ayant, S. Asawasamrit and J. Tariboon, *Certain finite integrals related to the product of special functions*, Symmetry **13** (2021), Article ID 2013, 11 pages. <https://doi.org/10.3390/sym13112013>
- [18] D. Kumar, F. Y. Ayant and J. Choi, *Application of product of the multivariable A-function and the multivariable Srivastava polynomials*, East Asian Math. J. **34**(3) (2018), 295–303. <https://doi.org/10.7858/eamj.2018.021>
- [19] D. Kumar, F. Y. Ayant and D. Kumar, *A new class of integrals involving generalized hypergeometric function and multivariable Aleph-function*, Kragujevac J. Math. **44**(4) (2020), 539–550. <https://doi.org/10.46793/KgJMat2004.539K>
- [20] D. Kumar, F. Y. Ayant and A. Prakash, *Certain integral involving the product of Srivastava polynomials and special functions*, Afr. Mat. **32** (2021), 1–9. <https://doi.org/10.1007/s13370-021-00885-7>
- [21] D. Kumar, S. D. Purohit and J. Choi, *Generalized fractional integrals involving product of multivariable H-function and a general class of polynomials*, J. Nonlinear Sci. Appl. **9** (2016), 8–21. <https://doi.org/10.1007/s13370-021-00885-7>



- [22] S. K. Kumari, V. T. M. Nambisan and A. K. Rathie, *A study of I-functions of two variables*, *Le Matematiche* **69**(1) (2014), 285–305.
- [23] J. Ram and D. Kumar, *Generalized fractional integration of the  $\aleph$ -function*, *J. Rajasthan Acad. Phys. Sci.* **10**(4) (2011), 373–382.
- [24] R. K. Saxena and D. Kumar, *Generalized fractional calculus of the Aleph-function involving a general class of polynomials*, *Acta Math. Sci. Ser. B (Engl. Ed.)* **35**(5) (2015), 1095–1110. [https://doi.org/10.1016/S0252-9602\(15\)30042-4](https://doi.org/10.1016/S0252-9602(15)30042-4)
- [25] R. K. Saxena, J. Ram and D. Kumar, *Generalized fractional integral of the product of two Aleph-functions*, *Appl. Appl. Math.* **8**(2) (2013), 631–646.
- [26] C. K. Sharma and S. S. Ahmad, *On the multivariable I-function*, *Acta Ciencia Indica (Mathematics)* **20**(2) (1994), 113–116.
- [27] C. K. Sharma and P. L. Mishra, *On the I-function of two variables and its properties*, *Acta Ciencia Indica (Mathematics)* **17** (1991), 667–672.
- [28] K. Sharma, *On the integral representation and applications of the generalized function of two variables*, *Int. J. Math. Eng. Sci.* **3** (2014), 1–13.
- [29] H. M. Srivastava, *A multi-linear generating function for the Konhauser set of biorthogonal polynomials suggested by Laguerre polynomial*, *Pacific. J. Math.* **177** (1985), 183–191.
- [30] N. Südländ, B. Baumann and T. F. Nonnenmacher, *Open problem: who knows about the Aleph-functions?*, *Fract. Calc. Appl. Anal.* **1**(4) (1998), 401–402.

<sup>1</sup>DEPARTMENT OF APPLIED SCIENCES,  
COLLEGE OF AGRICULTURE-JODHPUR,  
AGRICULTURE UNIVERSITY JODHPUR, JODHPUR-342304, INDIA  
*Email address:* dinesh\_dino03@yahoo.com

<sup>2</sup>COLLÈGE JEAN L'HERMINIER,  
ALLÉE DES NYMPHÉAS, 83500 LA SEYNE-SUR-MER, FRANCE  
<sup>3</sup>DEPARTMENT VAR, AVENUE JOSEPH RAYNAUD LE PARC FLEURI,  
83140 SIX-FOURS-LES-PLAGES, FRANCE  
*Email address:* fredericayant@gmail.com

<sup>4</sup>DEPARTMENT OF MATHEMATICS & STATISTICS,  
JAI NARAIN VYAS UNIVERSITY, JODHPUR-342005, INDIA  
*Email address:* nareshdudi@yahoo.co.in