

## NOTE ON STRONG PRODUCT OF GRAPHS

M. TAVAKOLI<sup>1</sup>, F. RAHBARNIA<sup>1</sup>, AND A. R. ASHRAFI<sup>2</sup>

*Dedicated to the memory of the late professor Ante Graovac*

ABSTRACT. Let  $G$  and  $H$  be graphs. The strong product  $G \boxtimes H$  of graphs  $G$  and  $H$  is the graph with vertex set  $V(G) \times V(H)$  and  $u = (u_1, v_1)$  is adjacent with  $v = (u_2, v_2)$  whenever  $(v_1 = v_2$  and  $u_1$  is adjacent with  $u_2)$  or  $(u_1 = u_2$  and  $v_1$  is adjacent with  $v_2)$  or  $(u_1$  is adjacent with  $u_2$  and  $v_1$  is adjacent with  $v_2)$ . In this paper, we study some properties of this operation. Also, we obtain lower and upper bounds for Wiener and hyper-Wiener indices of Strong product of graphs.

### 1. INTRODUCTION

Throughout this paper graphs means simple connected graphs. Suppose  $G$  is a graph with vertex set  $V(G)$ . The distance between the vertices  $u$  and  $v$  of  $V(G)$  is defined as the length of a minimal path connecting them, denoted by  $d(u, v)$ . The Wiener index,  $W(G)$ , is equal to the count of all shortest distances in a graph [20]. In other words,  $W(G) = \frac{1}{2} \sum_{u \in V(G)} \sum_{v \in V(G)} d(u, v)$ . We encourage to the interested reader to consult [6][7][9][10][16] for more information on this topic.

The hyper-Wiener index of acyclic graphs was introduced by Milan Randić in 1993. Then Klein et al. [17], generalized Randić's definition for all connected graphs, as a generalization of the Wiener index. It is defined as  $WW(G) = \frac{1}{2}W(G) + \frac{1}{2} \sum_{\{u,v\} \subseteq V(G)} d^2(u, v)$ . The mathematical properties and chemical meaning of this topological index are reported in [4][5][11][15][24].

The degree of a vertex  $v$  in a graph  $G$ ,  $deg_G(v)$ , is the number of edges of  $G$  incident with  $v$ . The eccentricity  $\varepsilon_G(u)$  is defined as the largest distance between  $u$  and other vertices of  $G$ . The eccentric connectivity index of a graph  $G$  is defined as

---

*Key words and phrases.* Strong product, Wiener index, Eulerian graph.

*2010 Mathematics Subject Classification.* Primary: 05C76. Secondary: 05C12, 05C07.

*Received:* March 14, 2012.

*Revised:* November 18, 2012.

$\xi^e(G) = \sum_{u \in V(G)} \deg(u)\varepsilon(u)$  [19]. The investigation of the mathematical properties of  $\xi^e(G)$  started only recently, and has so far resulted in determining the extremal values of the invariant and the extremal graphs where those values are achieved [13][22][23], and also in a number of explicit formulae for the eccentric connectivity index of several classes of graphs [1].

The Strong product  $G \boxtimes H$  of graphs  $G$  and  $H$  has the vertex set  $V(G \boxtimes H) = V(G) \times V(H)$  and  $(a, x)(b, y)$  is an edge of  $G \boxtimes H$  if  $a = b$  and  $xy \in E(H)$ , or  $ab \in E(G)$  and  $x = y$ , or  $ab \in E(G)$  and  $xy \in E(H)$ . Occasionally one also encounters the names strong direct product or symmetric composition for the strong product [12]. It is worthy to mention here that [8] and [21] are the first two papers that considered the problem of distribution of a topological index over a graph operation.

Throughout this paper our notation is standard and taken mainly from [2] and [3].

## 2. MAIN RESULTS

For a connected graph  $G$ , the radius  $r(G)$  and diameter  $D(G)$  are, respectively, the minimum and maximum eccentricity among vertices of  $G$ .

**Lemma 2.1.** *Let  $G$  and  $H$  be graphs. Then for every vertex  $(a, x)$  of  $G \boxtimes H$ , we have*

$$\varepsilon_{G \boxtimes H}((a, x)) = \max\{\varepsilon_G(a), \varepsilon_H(x)\}.$$

*Proof.* Let  $(a, x) \in V(G \boxtimes H)$ . By definition of the eccentricity,

$$\varepsilon_{G \boxtimes H}((a, x)) = \max\{d_{G \boxtimes H}((a, x), (b, y)) \mid (b, y) \in V(G \boxtimes H)\}.$$

On the other hand, it is well-known that  $d_{G \boxtimes H}((a, x), (b, y)) = \max\{d_G(a, b), d_H(x, y)\}$  [12], and so

$$\begin{aligned} \varepsilon_{G \boxtimes H}((a, x)) &= \max\{\max\{d_G(a, b)\}, \max\{d_H(x, y)\} \mid b \in V(G), y \in V(H)\} \\ &= \max\{\varepsilon_G(a), \varepsilon_H(x)\}, \end{aligned}$$

which completes the proof. □

A vertex is called odd vertex if it has odd degree.

**Theorem 2.1.** *Let  $G$  and  $H$  be a nontrivial connected graphs. Then  $G \boxtimes H$  is eulerian if and only if  $G$  and  $H$  are eulerian.*

*Proof.* Let  $G$  and  $H$  are eulerian then clearly  $G \boxtimes H$  is eulerian. Conversely, we assume that  $G \boxtimes H$  is eulerian. Suppose first that one of graphs  $G$  and  $H$  is not eulerian. Without loss of generality, we may assume that  $G$  is not eulerian. Thus,  $G$  has an odd vertex  $u$ . Let  $x$  is a vertex of  $H$ . Then,  $\deg_{G \boxtimes H}((u, x))$  is odd. We conclude that  $G \boxtimes H$  is not eulerian if one of graphs  $G$  and  $H$  is not eulerian.

Assume next that  $G$  and  $H$  are not eulerian. Then  $G$  and  $H$  have odd vertices  $u$  and  $x$ , respectively. Therefore, the vertex  $(u, x)$  of  $G \boxtimes H$  has odd degree and hence  $G \boxtimes H$  is not eulerian in this case, which completes the argument. □

The complement or inverse of a graph  $G$  is a graph  $\bar{G}$  on the same vertices such that two vertices of  $\bar{G}$  are adjacent if and only if they are not adjacent in  $G$ . We denote the complete graph of order  $n$  by  $K_n$ .

**Theorem 2.2.** *Let  $G$  and  $H$  be nontrivial connected graphs. Then*

$$\begin{aligned} W(G \boxtimes H) \geq & (|V(G)| + 2|E(G)|)W(H) + (|V(H)| + 2|E(H)|)W(G) \\ & + |V(G)||V(H)|(|V(G)||V(H)| - |V(G)| - |V(H)| + 1) \\ & - 2|E(G)||V(H)|(|V(H)| - 1) - 2|E(H)|(|V(G)|^2 - |V(G)| - |E(G)|) \end{aligned}$$

with equality if and only if  $\max\{D(G), D(H)\} \leq 2$ , or  $G \cong K_n$ , or  $H \cong K_n$ .

*Proof.* Let  $G$  and  $H$  are nontrivial connected graphs. Suppose  $A$  is the set of vertex-pairs  $(a, x), (a, y)$  that  $a \in V(G)$ ,  $x \neq y$  and  $x, y \in V(H)$ . Then,

$$(2.1) \quad \sum_{(u,v) \in A} d_{G \boxtimes H}(u, v) = |V(G)|W(H).$$

Similarly, for vertex-pairs  $(a, x), (b, x)$  that  $x \in V(H)$ ,  $a \neq b$  and  $a, b \in V(G)$ , we have

$$(2.2) \quad \sum_{\{(a,x),(b,x)\}} d_{G \boxtimes H}((a, x), (b, x)) = |V(H)|W(G).$$

On the other hand, the sum of all distances between vertex-pairs  $(a, x), (b, y)$  that  $(a \neq b \in V(G)$  and  $xy \in E(H))$  or  $(x \neq y \in V(H)$  and  $ab \in E(G))$ , is equal to

$$(2.3) \quad 2|E(G)|W(H) + 2|E(H)|W(G) - 2|E(G)||E(H)|$$

By this fact that for every vertex-pair  $(a, x), (b, y)$  of  $G \boxtimes H$  such that  $ab \in E(\bar{G})$  and  $xy \in E(\bar{H})$ , we have  $d_{G \boxtimes H}((a, x), (b, y)) \geq 2$ , Equations (2.1), (2.2) and (2.3), the result is proved.  $\square$

Using similar arguments as Theorem 2.2 one can prove the following result.

**Theorem 2.3.** *Let  $G$  and  $H$  be nontrivial connected graphs. Then*

$$\begin{aligned} W(G \boxtimes H) \leq & (|V(G)| + 2|E(G)|)W(H) + (|V(H)| + 2|E(H)|)W(G) \\ & + D \left[ \frac{|V(G)||V(H)|}{2} (|V(G)||V(H)| - |V(G)| - |V(H)| + 1) \right. \\ & \left. - 2|E(H)| \binom{|V(G)|}{2} - 2|E(G)| \binom{|V(H)|}{2} \right] + 2|E(G)||E(H)|(D - 1). \end{aligned}$$

where  $D = \max\{D(G), D(H)\}$ . Moreover, the upper bound is attained if and only if  $D \leq 2$ , or  $G \cong K_n$ , or  $H \cong K_n$ .

**Theorem 2.4.** *Let  $G$  and  $H$  be a nontrivial connected graphs. Then*

$$\begin{aligned} WW(G \boxtimes H) &\geq (|V(G)| + 2|E(G)|)WW(H) + (|V(H)| + 2|E(H)|)WW(G) \\ &\quad + \frac{3}{2}|V(G)||V(H)|(|V(G)||V(H)| - |V(G)| - |V(H)| + 1) \\ &\quad - 3|E(G)||V(H)|(|V(H)| - 1) - 3|E(H)|(|V(G)|^2 - |V(G)| \\ &\quad - \frac{4}{3}|E(G)|). \end{aligned}$$

with equality if and only if  $\max\{D(G), D(H)\} \leq 2$ , or  $G \cong K_n$ , or  $H \cong K_n$ .

*Proof.* We split the vertex-pair set of  $G \boxtimes H$  into subsets

$$\begin{aligned} A &= \{(a, x), (b, y) \mid a = b \in V(G) \text{ and } x \neq y, x, y \in V(H)\}, \\ B &= \{(a, x), (b, y) \mid a \neq b, a, b \in V(G) \text{ and } x = y \in V(H)\}, \\ C &= \{(a, x), (b, y) \mid x \neq y, x, y \in V(H) \text{ and } ab \in E(G)\}, \\ D &= \{(a, x), (b, y) \mid a \neq b, a, b \in V(G) \text{ and } xy \in E(H)\}, \\ E &= \{(a, x), (b, y) \mid ab \in E(\bar{G}) \text{ and } xy \in E(\bar{H})\}. \end{aligned}$$

It follows from the edge structure of  $G \boxtimes H$  that, if  $\{(a, x), (b, y)\} \in A \cup C$  then

$$d_{G \boxtimes H}((a, x), (b, y)) + d_{G \boxtimes H}^2((a, x), (b, y)) = d_H(x, y) + d_H^2(x, y),$$

if  $\{(a, x), (b, y)\} \in B \cup D$  then

$$d_{G \boxtimes H}((a, x), (b, y)) + d_{G \boxtimes H}^2((a, x), (b, y)) = d_G(a, b) + d_G^2(a, b),$$

and if  $\{(a, x), (b, y)\} \in E$  then

$$d_{G \boxtimes H}((a, x), (b, y)) + d_{G \boxtimes H}^2((a, x), (b, y)) \geq 6.$$

Therefore,

$$\begin{aligned} WW(G \boxtimes H) &= \frac{1}{2} \sum_{\{u,v\} \in A \cup D} (d_{G \boxtimes H}(u, v) + d_{G \boxtimes H}^2(u, v)) \\ &\quad + \frac{1}{2} \sum_{\{u,v\} \in B \cup C} (d_{G \boxtimes H}(u, v) + d_{G \boxtimes H}^2(u, v)) \\ &\quad + \frac{1}{2} \sum_{\{u,v\} \in E} (d_{G \boxtimes H}(u, v) + d_{G \boxtimes H}^2(u, v)) \\ &= (|V(G)| + 2|E(G)|)WW(H) \\ &\quad + (|V(H)| + 2|E(H)|)WW(G) - 2|E(G)||E(H)| \\ &\quad + \sum_{ab \in E(\bar{G})} \sum_{xy \in E(\bar{H})} (d_{G \boxtimes H}((a, x), (b, y)) + d_{G \boxtimes H}^2((a, x), (b, y))). \end{aligned}$$

On the other hand,

$$\begin{aligned} & \sum_{ab \in E(\bar{G})} \sum_{xy \in E(\bar{H})} (d_{G \boxtimes H}((a, x), (b, y)) + d_{G \boxtimes H}^2((a, x), (b, y))) \\ & \geq \frac{3}{2} |V(G)| |V(H)| (|V(G)| |V(H)| - |V(G)| - |V(H)| + 1) \\ & \quad - 3 |E(G)| |V(H)| (|V(H)| - 1) - 3 |E(H)| |V(G)| (|V(G)| - 1) \\ & \quad + 6 |E(G)| |E(H)|, \end{aligned}$$

which completes the proof.  $\square$

**Theorem 2.5.** *Let  $G$  and  $H$  be nontrivial connected graphs. Then*

$$\begin{aligned} WW(G \boxtimes H) & \leq (|V(G)| + 2|E(G)|) WW(H) + (|V(H)| + 2|E(H)|) WW(G) \\ & \quad + \frac{1}{2} D(D+1) \left[ \frac{|V(G)| |V(H)|}{2} (|V(G)| |V(H)| - |V(G)| - |V(H)| + 1) \right. \\ & \quad \left. - 2|E(H)| \binom{|V(G)|}{2} - 2|E(G)| \binom{|V(H)|}{2} \right] \\ & \quad + |E(G)| |E(H)| (D^2 + D - 2), \end{aligned}$$

where  $D = \max\{D(G), D(H)\}$ . Moreover, the upper bound is attained if and only if  $D \leq 2$ , or  $G \cong K_n$ , or  $H \cong K_n$ .

*Proof.* Using a similar argument as in Theorem 2.4, and the fact that for every  $ab \in E(\bar{G})$  and  $xy \in E(\bar{H})$ ,  $d_{G \boxtimes H}((a, x), (b, y)) \leq D$ , we obtain the result.  $\square$

The first Zagreb index of a graph  $G$  is defined as  $M_1(G) = \sum_{v \in V(G)} \deg^2(v)$  and the second Zagreb of  $G$  is given by  $M_2(G) = \sum_{uv \in E(G)} \deg(u)\deg(v)$ , see [18][14] for details.

**Theorem 2.6.** *For every graph  $G$  and  $H$ , we have*

$$\begin{aligned} M_1(G \boxtimes H) & = (|V(H)| + 4|E(H)|) M_1(G) + (|V(G)| + 4|E(G)|) M_1(H) \\ & \quad + M_1(G) M_1(H) + 8|E(G)| |E(H)|. \end{aligned}$$

*Proof.* It follows from the edge structure of  $G \boxtimes H$  that, for each vertex  $(a, x) \in E(G \boxtimes H)$ , we have

$$\deg_{G \boxtimes H}((a, x)) = \deg_G(a) + \deg_H(x) + \deg_G(a)\deg_H(x),$$

as desired.  $\square$

**Theorem 2.7.** *For every graph  $G$  and  $H$ , we have*

$$\begin{aligned} M_2(G \boxtimes H) & = 3|E(H)| M_1(G) + 3|E(G)| M_1(H) \\ & \quad + 3M_1(G) M_1(H) + 2M_2(G) M_2(H) \\ & \quad + (6|E(H)| + 3M_1(H) + |V(H)|) M_2(G) \\ & \quad + (6|E(G)| + 3M_1(G) + |V(G)|) M_2(H). \end{aligned}$$

*Proof.* Consider the following partition of  $E(G \boxtimes H)$

$$\begin{aligned} A &= \{(a, x)(b, y) \in E(G \boxtimes H) \mid ab \in E(G) \text{ and } xy \in E(H)\}, \\ B &= \{(a, x)(b, y) \in E(G \boxtimes H) \mid a = b \in V(G) \text{ and } xy \in E(H)\}, \\ C &= \{(a, x)(b, y) \in E(G \boxtimes H) \mid ab \in E(G) \text{ and } x = y \in V(H)\}. \end{aligned}$$

The sum of  $\deg_{G \boxtimes H}(u)\deg_{G \boxtimes H}(v)$  over all edges of  $A$ , is equal to

$$(2.4) \quad 2(|E(H)| + M_1(H) + M_2(H))M_2(G) + 2(|E(G)| + M_1(G))M_2(H) + M_1(G)M_1(H).$$

On the other hand, the summation of  $\deg_{G \boxtimes H}(u)\deg_{G \boxtimes H}(v)$  over all edges of  $B$ , is equal to

$$(2.5) \quad |E(H)|M_1(G) + (2|E(G)| + M_1(G))M_1(H) + (|V(G)| + 4|E(G)| + M_1(G))M_2(H),$$

and finally, summing  $\deg_{G \boxtimes H}(u)\deg_{G \boxtimes H}(v)$  over all edges of  $C$  we arrive at

$$(2.6) \quad |E(G)|M_1(H) + (2|E(H)| + M_1(H))M_1(G) + (|V(H)| + 4|E(H)| + M_1(H))M_2(G).$$

Now, by summation of (2.4), (2.5) and (2.6), the result can be proved.  $\square$

A connected graph is called a self-centered graph if all of its vertices have the same eccentricity [3]. Then a connected graph  $G$  is self-centered if and only if  $r(G) = D(G)$ .

**Theorem 2.8.** *Let  $G$  and  $H$  be self-centered graphs that  $D(H) \leq D(G)$ . Then*

$$\xi(G \boxtimes H) = 2r(G)(|E(G)||V(H)| + |E(H)||V(G)| + 2|E(G)||E(H)|).$$

*Proof.* The result follows from Lemma 2.1 and the fact that

$$|E(G \boxtimes H)| = |E(G)||V(H)| + |E(H)||V(G)| + 2|E(G)||E(H)|.$$

$\square$

**Acknowledgment:** The research of the third author is partially supported by the *University of Kashan* under Grant No. 159020/37.

## REFERENCES

- [1] A. R. Ashrafi, M. Saheli, M. Ghorbani, *The eccentric connectivity index of nanotubes and nanotori*, J. Comput. Appl. Math. **235** (2011) 4561–4566.
- [2] J. A. Bondy and U. S. R. Murty, *Graph theory*, Graduate Texts in Mathematics, vol. 244, Springer, New York, 2008.
- [3] F. Buckley, F. Harary, *Distance in Graphs*, Addison-Wesley Publishing Company, New York, 1989.
- [4] G. G. Cash, *Relationship between the Hosoya polynomial and the Hyper-Wiener index*, Appl. Math. Lett. **15** (2002) 893–895.
- [5] G. G. Cash, *Polynomial expressions for the hyper-Wiener index of extended hydrocarbon networks*, Comput. Chem. **25** (2001) 577–582.
- [6] A. A. Dobrynin, R. Entringer, I. Gutman, *Wiener index of trees: Theory and applications*, Acta Appl. Math. **66** (2001) 211–249.

- [7] A. A. Dobrynin, I. Gutman, S. Klavžar, P. Zigert, *Wiener index of hexagonal systems*, Acta Appl. Math. **72** (2002) 247–294.
- [8] A. Graovac, T. Pisanski, *On the Wiener index of a graph*, J. Math. Chem. **8** (1991) 53–62.
- [9] I. Gutman, S. Klavžar, B. Mohar (Eds.), *Fifty years of the Wiener index*, MATCH Commun. Math. Comput. Chem. **35** (1997), 1–259.
- [10] I. Gutman, S. Klavžar, B. Mohar (Eds.), *Fiftieth anniversary of the Wiener index*, Discrete Appl. Math. **80** (1) (1997), 1–113.
- [11] I. Gutman, *Relation between hyper-Wiener and Wiener index*, Chem. Phys. Lett. **364** (2002) 352–356.
- [12] R. Hammack, W. Imrich and S. Klavžar, *Handbook of Product Graphs*, Second edition, Taylor & Francis Group, 2011.
- [13] A. Ilić, I. Gutman, *Eccentric connectivity index of chemical trees*, MATCH Commun. Math. Comput. Chem. **65** (2011) 731–744.
- [14] M.H. Khalifeh, H. Yousefi-Azari, A.R. Ashrafi, *The first and second Zagreb indices of graph operations*, Discrete Appl. Math. **157** (2009) 804–811.
- [15] S. Klavžar, P. Zigert, I. Gutman, *An algorithm for the calculation of the hyper-Wiener index of benzenoid hydrocarbons*, Comput. Chem. **24** (2000) 229–233.
- [16] S. Klavžar, I. Gutman, *A theorem on Wiener-type invariants for isometric subgraphs of hypercubes*, Appl. Math. Lett. **19** (2006) 1129–1133.
- [17] D. J. Klein, I. Lukovits, I. Gutman, *On the definition of the hyper-Wiener index for cycle-containing structures*, J. Chem. Inf. Comput. Sci. **35** (1995) 50–52.
- [18] S. Nikolić, G. Kovačević, A. Miličević, N. Trinajstić, *The Zagreb Indices 30 Years After*, Croat. Chem. Acta **76** (2003) 113–124.
- [19] V. Sharma, R. Goswami, A. K. Madan, *Eccentric connectivity index: a novel highly discriminating topological descriptor for structure property and structure activity studies*, J. Chem. Inf. Comput. Sci. **37** (1997) 273–282.
- [20] H. Wiener, *Structural determination of the paraffin boiling points*, J. Amer. Chem. Soc. **69** (1947) 17–20.
- [21] Y. N. Yeh, I. Gutman, *On the sum of all distances in composite graphs*, Discrete Math. **135** (1994) 359–365.
- [22] G. Yu, L. Feng, A. Ilić, *On the eccentric distance sum of trees and unicyclic graphs*, J. Math. Anal. Appl. **375** (2011) 99–107.
- [23] B. Zhou, Z. Du, *On eccentric connectivity index*, MATCH Commun. Math. Comput. Chem. **63** (2010) 181–198.
- [24] B. Zhou, I. Gutman, *Relations between Wiener, hyper-Wiener and Zagreb indices*, Chem. Phys. Lett. **394** (2004) 93–95.

<sup>1</sup>DEPARTMENT OF MATHEMATICS,  
FERDOWSI UNIVERSITY OF MASHHAD,  
P. O. BOX 1159, MASHHAD 91775, I. R. IRAN

<sup>2</sup>DEPARTMENT OF MATHEMATICS,  
FACULTY OF MATHEMATICS, STATISTICS AND COMPUTER SCIENCE  
UNIVERSITY OF KASHAN,  
KASHAN 87317-51167, I. R. IRAN