

## A NEW OSTROWSKI-GRÜSS TYPE INEQUALITY

YU MIAO<sup>1</sup>, FEI HAN<sup>2,3</sup>, AND JIANYONG MU<sup>1</sup>

ABSTRACT. In the present note, we establish a similar inequality of Ostrowski-Grüss type for functions whose first derivative is absolutely continuous and second derivative is bounded both above and below almost everywhere.

### 1. INTRODUCTION

The following integral inequality which establishes a connection between the integral of the product of two functions and the product of the integrals of the two functions is well known in the literature as Grüss' inequality (see [15, p. 296]).

**Theorem 1.1.** *Let  $f, g : [a, b] \rightarrow R$  be two integrable functions such that  $\phi \leq f(x) \leq \Phi$  and  $\gamma \leq g(x) \leq \Gamma$  for all  $x \in [a, b]$ ,  $\phi, \Phi, \gamma$  and  $\Gamma$  are constants. Then we have*

$$\left| \frac{1}{b-a} \int_a^b f(x)g(x)dx - \frac{1}{b-a} \int_a^b f(x)dx \frac{1}{b-a} \int_a^b g(x)dx \right| \leq \frac{1}{4}(\Gamma - \gamma)(\Phi - \phi),$$

where the constant  $\frac{1}{4}$  is sharp.

Another celebrated integral inequality which provides an approximation of the integral mean  $\frac{1}{b-a} \int_a^b f(t)dt$  in terms of the values of  $f$  at a certain point  $x \in [a, b]$ , is Ostrowski's inequality [16, p. 468].

**Theorem 1.2.** *Let  $f : [a, b] \rightarrow R$  be continuous on  $[a, b]$  and differentiable on  $(a, b)$ , whose derivative  $f' : [a, b] \rightarrow R$  is bounded on  $(a, b)$ , i.e.*

$$\|f'\|_\infty = \sup_{t \in (a,b)} |f'(t)| < \infty.$$

---

*Key words and phrases.* Grüss inequality, Absolutely continuous, Ostrowski-Grüss type inequality.  
*2010 Mathematics Subject Classification.* 26D15.  
*Received:* January 28, 2013.

Then we have the inequality

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[ \frac{1}{4} + \frac{\left(x - \frac{a+b}{2}\right)^2}{(b-a)^2} \right] (b-a) \|f'\|_\infty$$

for all  $x \in [a, b]$ . The constant  $\frac{1}{4}$  is sharp.

In the years thereafter, numerous generalizations, extensions and variants of Grüss inequality have appeared in the literature (see [4–12]). Dragomir and Wang [8] proved the following Ostrowski type inequality in terms of the lower and upper bounds of the first derivative.

**Theorem 1.3.** *Let  $f : [a, b] \rightarrow R$  be continuous on  $[a, b]$  and differentiable on  $(a, b)$ , whose derivative satisfies the condition:  $\gamma \leq f'(x) \leq \Gamma$  for all  $x \in (a, b)$ . Then we have the inequality*

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt - \frac{f(b) - f(a)}{b-a} \left(x - \frac{a+b}{2}\right) \right| \leq \frac{1}{4} (b-a) (\Gamma - \gamma)$$

for all  $x \in [a, b]$ .

In Theorem 1.3, if we take  $x = a$ , then it follows that

$$(1.1) \quad \left| \frac{1}{b-a} \int_a^b f(t) dt - \frac{f(b) + f(a)}{2} \right| \leq \frac{1}{4} (b-a) (\Gamma - \gamma).$$

Cerone et al. [2] obtained the following result for twice differentiable mappings in terms of the upper and lower bounds of the second derivative.

**Theorem 1.4.** *Let  $f : [a, b] \rightarrow R$  be continuous on  $[a, b]$  and twice differentiable on  $(a, b)$ , and assume that the second derivative  $f'' : (a, b) \rightarrow R$  satisfies the condition  $\phi \leq f''(x) \leq \Phi$  for all  $x \in (a, b)$ . If we denote  $t = \frac{a+b}{2}$  then we have the inequality*

$$\begin{aligned} & \left| f(x) - (x-t)f'(x) - \frac{1}{b-a} \int_a^b f(t) dt + \left[ \frac{(b-a)^2}{24} + \frac{1}{2}(x-t)^2 \right] \frac{f'(b) - f'(a)}{b-a} \right| \\ & \leq \frac{1}{8} (\Phi - \phi) \left[ \frac{1}{2}(b-a) + |x-t| \right]^2 \end{aligned}$$

for all  $x \in [a, b]$ .

Motivated by the works of Cerone et al. [2], the purpose of the present note is to establish a similar inequality of Ostrowski-Grüss type to the inequality in Theorem 1.4 for functions whose first derivative is absolutely continuous and second derivative is bounded both above and below almost everywhere. From our inequality, we can obtain an analogue (see Remark 2.1) to (1.1).

2. MAIN RESULTS

**Theorem 2.1.** *Let  $f : [a, b] \rightarrow (-\infty, \infty)$  be a function such that the derivative  $f'$  is absolutely continuous on  $[a, b]$ . Assume that there exist constants  $\gamma, \Gamma \in (-\infty, \infty)$  such that  $\gamma \leq f''(x) \leq \Gamma$  a.e. on  $[a, b]$ . Then we have*

$$\begin{aligned} & \left| [f'(b) + f'(a)] \left[ \frac{1}{6}(b+a) + \frac{1}{4} \right] + \frac{bf'(b) + af'(a)}{6} + \frac{f(a)(1+2a) - f(b)(1+2b)}{2(b-a)} \right. \\ & \left. + \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \begin{cases} \frac{\Gamma-\gamma}{2(b-a)} \left( \frac{1}{6} + \frac{2}{3}C \right) \sqrt{1+4C}, & -\frac{3}{2} - 2b \leq a \leq -\frac{3}{4} - \frac{b}{2}; \\ -\frac{\Gamma-\gamma}{2(b-a)} \int_a^{x_2} (x^2 + x - C) dx, & -\frac{3}{4} - \frac{b}{2} \leq a \leq b; \\ -\frac{\Gamma-\gamma}{2(b-a)} \int_{x_1}^b (x^2 + x - C) dx, & a \leq -\frac{3}{2} - 2b, \end{cases} \end{aligned}$$

where

$$C := \frac{1}{b-a} \int_a^b (x + x^2) dx = \frac{1}{2}(b+a) + \frac{1}{3}(b^2 + ab + a^2),$$

and

$$x_1 = \frac{-1 - \sqrt{1+4C}}{2}, \quad x_2 = \frac{-1 + \sqrt{1+4C}}{2}.$$

*Proof.* It is not difficult to check that

$$\begin{aligned} & [f'(b) + f'(a)] \left[ \frac{1}{6}(b+a) + \frac{1}{4} \right] + \frac{bf'(b) + af'(a)}{6} + \frac{1}{b-a} \int_a^b f(x) dx \\ & + \frac{f(a)(1+2a) - f(b)(1+2b)}{2(b-a)} \\ & = \frac{1}{2} \left\{ \frac{1}{b-a} \int_a^b (x + x^2) f''(x) dx - \frac{1}{(b-a)^2} \int_a^b (x + x^2) dx \int_a^b f''(x) dx \right\} \\ (2.1) \quad & = \frac{1}{2(b-a)} \int_a^b \left\{ (x + x^2) - \frac{1}{b-a} \int_a^b (x + x^2) dx \right\} f''(x) dx. \end{aligned}$$

Let

$$\begin{aligned} A &= \left\{ x \in [a, b] : x + x^2 \geq \frac{1}{b-a} \int_a^b (x + x^2) dx \right\}; \\ A^c &= \left\{ x \in [a, b] : x + x^2 < \frac{1}{b-a} \int_a^b (x + x^2) dx \right\}. \end{aligned}$$

Then we have

$$\begin{aligned} & \int_a^b \left\{ (x+x^2) - \frac{1}{b-a} \int_a^b (x+x^2) dx \right\} f''(x) dx \\ & \leq \Gamma \int_A \left\{ (x+x^2) - \frac{1}{b-a} \int_a^b (x+x^2) dx \right\} dx \\ & \quad + \gamma \int_{A^c} \left\{ (x+x^2) - \frac{1}{b-a} \int_a^b (x+x^2) dx \right\} dx \end{aligned}$$

and

$$\begin{aligned} & \int_a^b \left\{ (x+x^2) - \frac{1}{b-a} \int_a^b (x+x^2) dx \right\} f''(x) dx \\ & \geq \gamma \int_A \left\{ (x+x^2) - \frac{1}{b-a} \int_a^b (x+x^2) dx \right\} dx \\ & \quad + \Gamma \int_{A^c} \left\{ (x+x^2) - \frac{1}{b-a} \int_a^b (x+x^2) dx \right\} dx. \end{aligned}$$

Since

$$\begin{aligned} & \int_A \left\{ (x+x^2) - \frac{1}{b-a} \int_a^b (x+x^2) dx \right\} dx \\ & = - \int_{A^c} \left\{ (x+x^2) - \frac{1}{b-a} \int_a^b (x+x^2) dx \right\} dx, \end{aligned}$$

it follows that

$$\begin{aligned} & \frac{1}{b-a} \left| \int_a^b \left\{ (x+x^2) - \frac{1}{b-a} \int_a^b (x+x^2) dx \right\} f''(x) dx \right| \\ & \leq \frac{\Gamma-\gamma}{b-a} \int_A \left\{ (x+x^2) - \frac{1}{b-a} \int_a^b (x+x^2) dx \right\} dx \\ (2.2) \quad & = -\frac{\Gamma-\gamma}{b-a} \int_{A^c} \left\{ (x+x^2) - \frac{1}{b-a} \int_a^b (x+x^2) dx \right\} dx. \end{aligned}$$

Putting

$$C := \frac{1}{b-a} \int_a^b (x+x^2) dx = \frac{1}{2}(b+a) + \frac{1}{3}(b^2+ab+a^2).$$

Therefore, from (2.2), it is enough to discuss the following integral,

$$(2.3) \quad \int_{A^c} (x^2+x-C) dx.$$

It is easy to see that  $1+4C < 0$  does not hold in  $x \in [a, b]$ , thus we only consider the case  $1+4C \geq 0$ . Denote the solutions of the equation  $x^2+x-C=0$  by  $x_1$  and  $x_2$ , then

$$x_1 = \frac{-1 - \sqrt{1+4C}}{2}, \quad x_2 = \frac{-1 + \sqrt{1+4C}}{2}.$$

**Case 1.**  $-\frac{3}{2} - 2b \leq a \leq -\frac{3}{4} - \frac{b}{2}$ .

This case implies  $b \geq -\frac{1}{2}$  and the claim

$$(2.4) \quad a \leq x_1 \leq x_2 \leq b.$$

Since  $-\frac{3}{2} - 2b \leq a \leq -\frac{3}{4} - \frac{b}{2}$ , we have

$$b \geq -\frac{1}{2}, \quad (b + 2a)(b - a) \leq \frac{3}{2}(b - a), \quad (a + 2b)(a - b) \leq \frac{3}{2}(b - a),$$

which yield

$$\sqrt{1 + 4C} \leq -2a - 1, \quad \sqrt{1 + 4C} \leq 2b + 1.$$

The claim (2.4) can be obtained. Then we have

$$\int_{A^c} (x^2 + x - C)dx = \int_{x_1}^{x_2} (x^2 + x - C)dx = -\left(\frac{1}{6} + \frac{2}{3}C\right)\sqrt{1 + 4C}.$$

**Case 2.**  $-\frac{3}{4} - \frac{b}{2} \leq a \leq b$ .

With the same proof of Case 1, it is easy to check

$$x_1 \leq a \leq x_2 \leq b.$$

Then we have

$$\begin{aligned} \int_{A^c} (x^2 + x - C)dx &= \int_a^{x_2} (x^2 + x - C)dx \\ &= -\frac{1}{6} [x_2(4C + 1) + 2a^3 + 3a^2 - 2aC + C]. \end{aligned}$$

**Case 3.**  $a \leq -\frac{3}{2} - 2b$ .

With the same proof of Case 1, it is easy to check

$$a \leq x_1 \leq b \leq x_2.$$

Then we have

$$\begin{aligned} \int_{A^c} (x^2 + x - C)dx &= \int_{x_1}^b (x^2 + x - C)dx \\ &= \frac{1}{6} [x_1(4C + 1) + 2b^3 + 3b^2 - 6bC - C]. \end{aligned}$$

From (2.1), (2.2) and above discussion, the desired result can be obtained. □

Next we shall show the following similar Ostrowski-Grüss type inequality.

**Theorem 2.2.** *Let  $f : [a, b] \rightarrow (-\infty, \infty)$  be a function such that the derivative  $f'$  is absolutely continuous on  $[a, b]$ . Assume that there exist constants  $\gamma, \Gamma \in (-\infty, \infty)$*

such that  $\gamma \leq f''(x) \leq \Gamma$  a.e. on  $[a, b]$ . Then for  $t \in [a, b]$ , we have

$$(2.5) \quad \left| f(t) + \frac{[f'(b) - f'(a)](t-a)(b-t)}{2(b-a)} + \frac{(b+a)(f'(b) + f'(a))}{2} + \frac{(t+a)f(a) - (b+t)f(b)}{b-a} \right| \leq \frac{\Gamma - \gamma}{b-a} G(t, a, b)$$

where if  $t+a > 0$ , then

$$G(t, a, b) = \frac{(t+a)}{2}(t^2 - C_1^2) - (at+C)(t-C_1) + \frac{(b+t)}{2}(b^2 - C_2^2) - (bt+C)(b-C_2);$$

if  $t+a < 0$ ,  $t+b < 0$ , then

$$G(t, a, b) = \frac{(t+a)}{2}(C_3^2 - a^2) - (at+C)(C_3 - a) + \frac{(b+t)}{2}(C_4^2 - t^2) - (bt+C)(C_4 - t);$$

if  $t+a < 0$ ,  $t+b > 0$ , then

$$G(t, a, b) = \frac{(t+a)}{2}(C_3^2 - a^2) - (at+C)(C_3 - a) + \frac{(b+t)}{2}(b^2 - C_2^2) - (bt+C)(b-C_2);$$

if  $t+a = 0$ ,  $t \neq b$ , then

$$G(t, a, b) = \frac{(b+t)}{2}(b^2 - C_2^2) - (bt+C)(b-C_2);$$

if  $t+b = 0$ ,  $t \neq a$ , then

$$G(t, a, b) = \frac{(t+a)}{2}(C_3^2 - a^2) - (at+C)(C_3 - a);$$

if  $t+a = 0$ ,  $t = b$  or  $t+b = 0$ ,  $t = a$ , then

$$G(t, a, b) = 0,$$

and where

$$\begin{aligned} C &:= \frac{1}{2}[t^2 - (b+a)t + (b^2 + ab + a^2)], \\ C_1 &= \left( \frac{at+C}{t+a} \vee a \right) \wedge t, \quad C_2 = \left( \frac{bt+C}{t+b} \vee t \right) \wedge b, \\ C_3 &= a \vee \left( \frac{at+C}{t+a} \wedge t \right), \quad C_4 = t \vee \left( \frac{bt+C}{t+b} \wedge b \right). \end{aligned}$$

*Proof.* Let

$$(2.6) \quad K(t, x) = \begin{cases} (t-x)(x-a), & a \leq x \leq t; \\ (x-t)(b-x), & t \leq x \leq b. \end{cases}$$

Then it is easy to see that

$$(2.7) \quad \frac{1}{b-a} \int_a^b (K(t, x) + x^2) dx = \frac{1}{2}[t^2 - (b+a)t + (b^2 + ab + a^2)],$$

and

$$(2.8) \quad \begin{aligned} & \frac{1}{b-a} \int_a^b (K(t, x) + x^2) f''(x) dx \\ & = f(t) + \frac{f'(b)b^2 - f'(a)a^2}{b-a} + \frac{(t+a)f(a) - (b+t)f(b)}{b-a}. \end{aligned}$$

Therefore, from (2.7) and (2.8), we get

$$(2.9) \quad \begin{aligned} & f(t) + \frac{[f'(b) - f'(a)](t-a)(b-t)}{2(b-a)} + \frac{(b+a)(f'(b) + f'(a))}{2} \\ & + \frac{(t+a)f(a) - (b+t)f(b)}{b-a} \\ & = \frac{1}{b-a} \int_a^b \left[ (K(t, x) + x^2) - \frac{1}{b-a} \int_a^b (K(t, x) + x^2) dx \right] f''(x) dx. \end{aligned}$$

Let

$$\begin{aligned} B & = \left\{ x \in [a, b], (K(t, x) + x^2) \geq \frac{1}{b-a} \int_a^b (K(t, x) + x^2) dx \right\}; \\ B^c & = \left\{ x \in [a, b], (K(t, x) + x^2) < \frac{1}{b-a} \int_a^b (K(t, x) + x^2) dx \right\}. \end{aligned}$$

As the proof of Theorem 2.1, we only need to estimate the bound of the following inequality

$$(2.10) \quad \begin{aligned} & \frac{1}{b-a} \left| \int_a^b \left[ (K(t, x) + x^2) - \frac{1}{b-a} \int_a^b (K(t, x) + x^2) dx \right] f''(x) dx \right| \\ & \leq \frac{\Gamma - \gamma}{b-a} \int_B \left[ (K(t, x) + x^2) - \frac{1}{b-a} \int_a^b (K(t, x) + x^2) dx \right] dx. \end{aligned}$$

Next we shall divide our proof into five cases.

**Case 1:**  $t + a > 0$ . Under this case, it follows that  $b + t > 0$ . Then if  $a \leq x \leq t$ , we have

$$K(t, x) + x^2 \geq C \iff x \geq \frac{at + C}{t + a},$$

and if  $t \leq x \leq b$ , we obtain

$$K(t, x) + x^2 \geq C \iff x \geq \frac{bt + C}{t + b}.$$

Therefore

$$\begin{aligned} & \int_B \left[ (K(t, x) + x^2) - \frac{1}{b-a} \int_a^b (K(t, x) + x^2) dx \right] dx \\ & = \frac{(t+a)}{2} (t^2 - C_1^2) - (at + C)(t - C_1) + \frac{(b+t)}{2} (b^2 - C_2^2) - (bt + C)(b - C_2), \end{aligned}$$

where

$$C_1 = \left( \frac{at + C}{t + a} \vee a \right) \wedge t, \quad C_2 = \left( \frac{bt + C}{t + b} \vee t \right) \wedge b.$$

**Case 2:**  $t + a < 0, t + b < 0$ . Then if  $a \leq x \leq t$ , we have

$$K(t, x) + x^2 \geq C \iff x \leq \frac{at + C}{t + a},$$

and if  $t \leq x \leq b$ , we obtain

$$K(t, x) + x^2 \geq C \iff x \leq \frac{bt + C}{t + b},$$

Therefore

$$\begin{aligned} & \int_B \left[ (K(t, x) + x^2) - \frac{1}{b - a} \int_a^b (K(t, x) + x^2) dx \right] dx \\ &= \frac{(t + a)}{2} (C_3^2 - a^2) - (at + C)(C_3 - a) + \frac{(b + t)}{2} (C_4^2 - t^2) - (bt + C)(C_4 - t), \end{aligned}$$

where

$$C_3 = a \vee \left( \frac{at + C}{t + a} \wedge t \right), \quad C_4 = t \vee \left( \frac{bt + C}{t + b} \wedge b \right).$$

**Case 3:**  $t + a < 0, t + b > 0$ . Then if  $a \leq x \leq t$ , we have

$$K(t, x) + x^2 \geq C \iff x \leq \frac{at + C}{t + a},$$

and if  $t \leq x \leq b$ , we obtain

$$K(t, x) + x^2 \geq C \iff x \geq \frac{bt + C}{t + b}.$$

Therefore

$$\begin{aligned} & \int_B \left[ (K(t, x) + x^2) - \frac{1}{b - a} \int_a^b (K(t, x) + x^2) dx \right] dx \\ &= \frac{(t + a)}{2} (C_3^2 - a^2) - (at + C)(C_3 - a) + \frac{(b + t)}{2} (b^2 - C_2^2) - (bt + C)(b - C_2). \end{aligned}$$

**Case 4:**  $t + a = 0$ . This case implies  $a \leq 0$  and  $t = -a \geq 0$ . Furthermore, it is easy to see that

$$at + C = \frac{1}{2} [t^2 - (b - a)t + (b^2 + ab + a^2)] \geq 0, \quad \forall t \in [a, b].$$

Then if  $a \leq x \leq t$ , we have

$$\begin{aligned} \{x : K(t, x) + x^2 \geq C\} &= \emptyset, \quad \text{for } t \neq b \\ \{x : K(t, x) + x^2 = C\} &= [a, t], \quad \text{for } t = b \end{aligned}$$

and if  $t \leq x \leq b$ , we obtain

$$K(t, x) + x^2 \geq C \iff x \geq \frac{bt + C}{t + b} \quad \text{for } t \neq b.$$

Therefore,

$$\begin{aligned} & \int_B \left[ (K(t, x) + x^2) - \frac{1}{b-a} \int_a^b (K(t, x) + x^2) dx \right] dx \\ &= \begin{cases} \frac{(b+t)}{2}(b^2 - C_2^2) - (bt + C)(b - C_2), & \text{for } t \neq b; \\ 0, & \text{for } t = b. \end{cases} \end{aligned}$$

**Case 5:**  $t + b = 0$ . This case implies  $b \geq 0$  and  $t = -b \leq 0$ . Furthermore, it is easy to see that

$$bt + C = \frac{1}{2}[t^2 - (a - b)t + (b^2 + ab + a^2)] \geq 0, \quad \forall t \in [a, b].$$

Then if  $a \leq x \leq t$ , we have

$$K(t, x) + x^2 \geq C \iff x \leq \frac{at + C}{t + a} \quad \text{for } t \neq a,$$

and if  $t \leq x \leq b$ , we obtain

$$\begin{aligned} \{x : K(t, x) + x^2 \geq C\} &= \emptyset \quad \text{for } t \neq a, \\ \{x : K(t, x) + x^2 = C\} &= [a, b] \quad \text{for } t = a. \end{aligned}$$

Therefore,

$$\begin{aligned} & \int_B \left[ (K(t, x) + x^2) - \frac{1}{b-a} \int_a^b (K(t, x) + x^2) dx \right] dx \\ &= \begin{cases} \frac{(t+a)}{2}(C_3^2 - a^2) - (at + C)(C_3 - a), & \text{for } t \neq a; \\ 0 & \text{for } t = a. \end{cases} \end{aligned}$$

□

*Remark 2.1.* We give an example for the case  $t = a$  to show a perturbed two points inequality. Let  $a \geq 0$ , then  $b > 0$ . It is easy to get

$$C = \frac{1}{2}(a^2 + b^2), \quad C_1 = a, \quad C_2 = \frac{1}{2}(a + b), \quad G(t, a, b) = \frac{1}{8}(b - a)^2(b + a).$$

Similarly, if  $a < 0$  and  $a + b > 0$ , then

$$C = \frac{1}{2}(a^2 + b^2), \quad C_3 = a, \quad C_2 = \frac{1}{2}(a + b), \quad G(t, a, b) = \frac{1}{8}(b - a)^2(b + a);$$

if  $a < 0$  and  $a + b < 0$ , then

$$C = \frac{1}{2}(a^2 + b^2), \quad C_3 = a, \quad C_4 = \frac{1}{2}(a + b), \quad G(t, a, b) = \frac{1}{8}(b - a)^2(b + a);$$

Hence, we have

$$\left| \frac{1}{b-a}(f(b) - f(a)) - \frac{1}{2}(f'(b) + f'(a)) \right| \leq \frac{1}{8}(\Gamma - \gamma)(b - a).$$

If we rewrite the above inequality as follows

$$\left| \frac{1}{b-a} \int_a^b f'(x) dx - \frac{1}{2}(f'(b) + f'(a)) \right| \leq \frac{1}{8}(\Gamma - \gamma)(b-a),$$

then, the bound is formally better than the inequality (1.1).

*Remark 2.2.* If we take  $b > 0$ ,  $a = -b$  and  $t = 0$ , then we have

$$C = \frac{1}{2}b^2, \quad C_3 = -\frac{1}{2}b, \quad C_2 = \frac{1}{2}b, \quad G(t, a, b) = \frac{1}{4}b^3,$$

which implies

$$\left| f(0) + \frac{b}{4}(f'(b) - f'(a)) - \frac{1}{2}(f(b) + f(a)) \right| \leq \frac{b^2(\Gamma - \gamma)}{8}.$$

#### REFERENCES

- [1] K. Alam and K. M. L. Saxena, *Positive dependence in multivariate distributions*, Comm. Statist. A-Theory Methods, **10** (1981), no. 12, 1183–1196.
- [2] P. Cerone, S. S. Dragomir and J. Roumeliotis, *An inequality of Ostrowski-Grüss type for twice differentiable mappings and applications in numerical integration*. Kyungpook Math. J. **39** (1999), no. 2, 333-341.
- [3] X. K. Chai and Y. Miao, *Note on the Grüss Inequality*. Gen. Math. **19** (2011), no. 3, 93–99.
- [4] X. L. Cheng and J. Sun, *A note on the perturbed trapezoid inequality*, JIPAM. J. Inequal. Pure Appl. Math. **3** (2002), no. 2, Article 29.
- [5] S. S. Dragomir, *A companion of the Grüss inequality and applications*, Appl. Math. Lett., **17** (2004), no. 4, 429-435.
- [6] S. S. Dragomir, P. Cerone and A. Sofo, *Some remarks on the trapezoid rule in numerical integration*, Indian J. Pure Appl. Math., **31** (2000), no. 5, 475-494.
- [7] S. S. Dragomir, Y. J. Cho and S. S. Kim, *Inequalities of Hadamard's type for Lipschitzian mappings and their applications*. J. Math. Anal. Appl., **245** (2000), no. 2, 489-501.
- [8] S. S. Dragomir and S. Wang, *An inequality of Ostrowski-Grüss type and its applications to the estimation of error bounds for some special means and for some numerical quadrature rules*, Comput. Math. Appl., **33** (1997), no. 11, 15-20.
- [9] N. Elezović, Lj. Marangunić and J. Pečarić, *Some improvements of Grüss type inequality*. J. Math. Inequal. **1** (2007), no. 3, 425-436.
- [10] Z. Liu, *A sharp integral inequality of Ostrowski-Grüss type*, Soochow J. Math., **32** (2006), no. 2, 223-231.
- [11] M. Matić, J. Pečarić and N. Ujević, *Improvement and further generalization of inequalities of Ostrowski-Grüss type*, Comput. Math. Appl., **39** (2000), no. 3-4, 161-175.
- [12] A. M. Mercer, *An improvement of the Grüss inequality*. JIPAM. J. Inequal. Pure Appl. Math., **6** (2005), no. 4, Article 93.

- [13] Y. Miao, *Note on the discrete Ostrowski-Grüss type inequality*. J. Math. Inequal. **3** (2009), no. 1, 93-98.
- [14] Y. Miao and L. M. Shen, *Some general Ostrowski-Grüss type inequalities*. Acta Math. Acad. Paedagog. Nyhazi. (N.S.), **24** (2008), no. 2, 243–248.
- [15] D. S. Mitrinović, J. E. Pečarić and A. M. Fink, *Classical and new inequalities in analysis*. Mathematics and its Applications (East European Series), 61. Kluwer Academic Publishers Group, Dordrecht, 1993.
- [16] D. S. Mitrinović, J. E. Pečarić and A. M. Fink, *Inequalities for Functions and Their Integrals and Derivatives*. Kluwer Academic Publishers, Dordrecht, 1994.
- [17] G. H. Peng and Y. Miao, *A note on Grüss type inequality*. Appl. Math. Sci. (Ruse) **3** (2009), no. 8, 399-402.

<sup>1</sup>COLLEGE OF MATHEMATICS AND INFORMATION SCIENCE,  
HENAN NORMAL UNIVERSITY,  
HENAN PROVINCE, 453007, CHINA.  
*E-mail address:* yumiao728@gmail.com  
*E-mail address:* jianyongmu@163.com

<sup>2</sup>SCHOOL OF MANAGEMENT,  
UNIVERSITY OF SHANGHAI FOR SCIENCE AND TECHNOLOGY,  
SHANGHAI, 200093, CHINA.

<sup>3</sup>DEPARTMENT OF MATHEMATICS AND INFORMATION SCIENCE,  
XINXIANG UNIVERSITY,  
HENAN PROVINCE, 453000, CHINA.  
*E-mail address:* tomcumt@126.com