

CHARACTERISTIC POLYNOMIAL OF SOME CLUSTER GRAPHS

PRABHAKAR R. HAMPIHOLI¹ AND BASAVRAJ S. DURGI²

ABSTRACT. The characteristic polynomial of a graph G with p vertices is defined as $\phi(G : \lambda) = \det(\lambda I - A(G))$, where A is the adjacency matrix of G and I is the unit matrix. The roots of the characteristic equation $\phi(G : \lambda) = 0$, denoted by $\lambda_1, \lambda_2, \dots, \lambda_p$ are the eigenvalues of G . The graphs with large number of edges are referred as graph representations of inorganic clusters, called as Cluster graphs. In this paper we obtain the characteristic polynomial of class of cluster graphs.

1. INTRODUCTION

Let G be a simple undirected graph with p vertices and q edges. Let $V(G) = \{v_1, v_2, \dots, v_p\}$ be the vertex set of G . The adjacency matrix of G is defined as $A(G) = [a_{ij}]$, where $a_{ij} = 1$ if v_i is adjacent to v_j and $a_{ij} = 0$, otherwise. The characteristic polynomial of G is $\phi(G : \lambda) = \det(\lambda I - A(G))$, where I is an unit matrix of order p . The roots of an equation $\phi(G : \lambda) = 0$ denoted by $\lambda_1, \lambda_2, \dots, \lambda_p$ are the eigenvalues of G and their collection is the spectrum of G [1]. Characteristic polynomial of some cluster graphs are obtained in [2–5]. In this paper we obtain the characteristic polynomial of some more cluster graphs.

2. SOME EDGE DELETED CLUSTER GRAPHS

I. Gutman and L. Pavlović [2] introduced four classes of graphs obtained from complete graph by deleting edges. For the sake of continuity we produce these here.

Definition 2.1. [2] Let v be a vertex of the complete graph K_p , $p \geq 3$ and let e_i , $i = 1, 2, \dots, k$, $1 \leq k \leq p - 1$, be its distinct edges, all being incident to v . The graph $Ka_p(k)$ or $Cl(p, k)$ is obtained by deleting e_i , $i = 1, 2, \dots, k$ from K_p . In addition $Ka_p(0) \cong K_p$.

Key words and phrases. Spectra, Complete graph, Cluster graphs.
2010 Mathematics Subject Classification. 05C50.
Received: January 13, 2013.
Revised: June 21, 2013.

Definition 2.2. [2] Let $f_i, i = 1, 2, \dots, k, 1 \leq k \leq \lfloor p/2 \rfloor$ be independent edges of the complete graph $K_p, p \geq 3$. The graph Kb_p is obtained by deleting $f_i, i = 1, 2, \dots, k$ from K_p . The graph $Kb_p(0) \cong K_p$.

Definition 2.3. [2] Let V_k be a k - element subset of the vertex set of the complete graph $K_p, 2 \leq k \leq p, p \geq 3$. The graph $Kc_p(k)$ is obtained by deleting from K_p all the edges connecting pairs of vertices from V_k . In addition $Kc_p(0) \cong Kc_p(1) \cong K_p$.

Definition 2.4. [2] Let $3 \leq k \leq p, p \geq 3$. The graph $Kd_p(k)$ obtained by deleting from K_p , the edges belonging to a k - membered cycle.

Theorem 2.1. [2] For $p \geq 3$ and $3 \leq k \leq p - 1$, it holds that

$$\phi(Ka_p(k) : \lambda) = (\lambda + 1)^{p-3}[\lambda^3 - (p - 3)\lambda^2 - (2p - k - 3)\lambda + (k - 1)(p - 1 - k)].$$

Theorem 2.2. [2] For $p \geq 3$ and $0 \leq k \leq \lfloor p/2 \rfloor$ it holds that

$$\phi(Kb_p(k) : \lambda) = \lambda^k(\lambda + 1)^{p-2k-1}(\lambda + 2)^{k-1}[\lambda^2 - (p - 3)\lambda - 2(p - k - 1)].$$

Theorem 2.3. [2] For $p \geq 3$ and $0 \leq k \leq p$ it holds that

$$\phi(Kc_p(k) : \lambda) = \lambda^{k-1}(\lambda + 1)^{p-k-1}[\lambda^2 - (p - k - 1)\lambda - k(p - k)].$$

Theorem 2.4. [2] For $p \geq 3$ and $3 \leq k \leq p$ it holds that

$$\begin{aligned} \phi(Kd_p(k) : \lambda) &= (\lambda + 1)^{p-k-1}[\lambda^2 - (p - 4)\lambda - (3p - 2k - 3)] \\ &\times \prod_{i=1}^{k-1} \left(\lambda + 2 \cos \left(\frac{2\pi i}{k} \right) + 1 \right). \end{aligned}$$

We introduce here another class of graphs obtained from K_p and we denote it by $Ka_p(p, m, k)$. Two subgraphs G_1 and G_2 of graph G are independent subgraphs if $V(G_1) \cap V(G_2)$ is an empty set, where $V(G)$ is the vertex set of the graph G .

Definition 2.5. [4] Let $(K_m)_i, i = 1, 2, \dots, k, 1 \leq k \leq \lfloor p/m \rfloor, 1 \leq m \leq p$, be independent complete subgraphs with m vertices of the complete graph $K_p, p \geq 3$. The graph $Ka_p(m, k)$ obtained from K_p by deleting all the edges of k independent complete subgraphs $(K_m)_i, i = 1, 2, \dots, k$. In addition $Ka_p(m, 0) \cong Ka_p(0, k) \cong Ka_p(0, 0) \cong K_p$.

We note that for $k = 1, Ka_p(p, m, k) \cong Ka_p(p, m)$.

Theorem 2.5. [4] Let p, m, k be positive integers. Let $mk \leq p$. Then for $p \geq 3, 0 \leq k \leq \lfloor p/m \rfloor, 1 \leq m \leq p$, the characteristic polynomial of $Ka_p(p, m, k)$ is

$$\phi(Ka_p(p, m, k) : \lambda) = \lambda^{mk-k}(\lambda + 1)^{p-mk-1}(\lambda + m)^{k-1}[\lambda^2 - (p - m - 1)\lambda - m(p + k - mk - 1)].$$

Definition 2.6. [5] Let K_{m_1} and K_{m_2} be independent complete subgraphs of K_p . The graph obtained by deleting all the edges of $k_1 + k_2$ such that $m_1k_1 + m_2k_2 \leq p$ independent complete subgraphs K_{m_1} (k_1 copies) and K_{m_2} (k_2 copies) is denoted by $K_{a_p}(p, (m_1, k_1), (m_2, k_2))$.

Theorem 2.6. [5] Let $p, m_i, k_i, i = 1, 2$ are positive integers with $\sum_{i=1}^2 m_i k_i \leq p$. Then for $p \geq 3$, the characteristic polynomial of $K_{a_p}(p, (m_1, k_1), (m_2, k_2))$ is

$$\begin{aligned} \phi(K_{a_p}(p, (m_1, k_1), (m_2, k_2)) : \lambda) &= \lambda^{\sum_{i=1}^2 m_i k_i - \sum_{i=1}^2 k_i} (\lambda + 1)^{p - \sum_{i=1}^2 m_i k_i - 1} \\ &\times \left\{ \left[(\lambda + m_1 - p) + \frac{k_2(m_2^2 - m_1 m_2)}{\lambda + m_2} \right] (\lambda + 1) + (m_1 - 1) \left(\sum_{i=1}^2 m_i k_i - p \right) \right\} \\ &\times (\lambda + m_1)^{k_1 - 1} (\lambda + m_2)^{k_2}. \end{aligned}$$

Now we proceed to prove the main result.

Definition 2.7. Let $K_{m_i}, i = 1, 2, \dots, l$ are the independent complete subgraphs of K_p . The graph obtained by deleting all the edges of $\sum_{i=1}^l k_i$, such that $\sum_{i=1}^l m_i k_i \leq p$ independent complete subgraphs K_{m_1} (k_1 copies), K_{m_2} (k_2 copies), ... , K_{m_l} (k_l copies) is denoted by $K_{a_p}(p, (m_1, k_1), (m_2, k_2), \dots, (m_l, k_l))$.

We note that for $k_1 = k, m_1 = m, k_i = 0$ for $i = 2, 3, \dots, l$ it holds that

$$K_{a_p}(p, (m_1, k_1), (m_2, k_2), \dots, (m_l, k_l)) \cong K_{a_p}(p, m, k).$$

Theorem 2.7. Let $p, m_i, k_i, i = 1, 2, \dots, l$ are positive integers with $\sum_{i=1}^l m_i k_i \leq p$. Then for $p \geq 3$, the characteristic polynomial of $K_{a_p}(p, (m_1, k_1), (m_2, k_2), \dots, (m_l, k_l))$ is

$$\begin{aligned} \phi(K_{a_p}(p, (m_1, k_1), (m_2, k_2), \dots, (m_l, k_l)) : \lambda) &= \lambda^{\sum_{i=1}^l m_i k_i - \sum_{i=1}^l k_i} (\lambda + 1)^{p - \sum_{i=1}^l m_i k_i - 1} \\ &\times \left\{ \left[(\lambda + m_1 - p) + \sum_{i=2}^l \frac{k_i(m_i^2 - m_i m_1)}{\lambda + m_i} \right] (\lambda + 1) + (m_1 - 1) \left(\sum_{i=1}^l m_i k_i - p \right) \right\} \\ &\times (\lambda + m_1)^{k_1 - 1} \prod_{i=2}^l (\lambda + m_i)^{k_i}. \end{aligned}$$

Proof. Without loss of generality we assume that the vertices of $K_{a_p}(p, (m_1, k_1), (m_2, k_2), \dots, (m_l, k_l))$ are the vertices of K_p and the edges are the edges of K_p that are not there in K_{m_1} (k_1 copies), K_{m_2} (k_2 copies), ... , K_{m_l} (k_l copies). The characteristic polynomial of $K_{a_p}(p, (m_1, k_1), (m_2, k_2), \dots, (m_l, k_l))$ is

$$\begin{vmatrix}
 \lambda & 0 & \cdots & 0 & -1 & -1 & \cdots & -1 & -1 & -1 & \cdots & -1 \\
 0 & \lambda & \cdots & 0 & -1 & -1 & \cdots & -1 & -1 & -1 & \cdots & -1 \\
 \vdots & \vdots & & \vdots & & & & \vdots & & & & \vdots & & & \vdots & & & \vdots & & & \vdots \\
 0 & 0 & \cdots & \lambda & -1 & -1 & \cdots & -1 & -1 & -1 & \cdots & -1 \\
 -1 & -1 & \cdots & -1 & \lambda & 0 & \cdots & 0 & -1 & -1 & \cdots & -1 \\
 -1 & -1 & \cdots & -1 & 0 & \lambda & \cdots & 0 & -1 & -1 & \cdots & -1 \\
 \vdots & \vdots & & \vdots & & & & \vdots & & & & \vdots & & & \vdots & & & \vdots & & & \vdots \\
 -1 & -1 & \cdots & -1 & 0 & 0 & \cdots & \lambda & -1 & -1 & \cdots & -1 \\
 -1 & -1 & \cdots & -1 & -1 & -1 & \cdots & -1 & \lambda & 0 & \cdots & 0 & -1 & \cdots & -1 & -1 & \cdots & -1 & -1 & \cdots & -1 \\
 -1 & -1 & \cdots & -1 & -1 & -1 & \cdots & -1 & 0 & \lambda & \cdots & 0 & -1 & \cdots & -1 & -1 & \cdots & -1 & -1 & \cdots & -1 \\
 \vdots & \vdots & & \vdots & & & & \vdots & & & & \vdots & & & \vdots & & & \vdots & & & \vdots \\
 -1 & -1 & \cdots & -1 & -1 & -1 & \cdots & -1 & 0 & 0 & \cdots & \lambda & -1 & \cdots & -1 & -1 & \cdots & -1 & -1 & \cdots & -1 \\
 -1 & -1 & \cdots & -1 & -1 & -1 & \cdots & -1 & -1 & -1 & \cdots & -1 & \lambda & \cdots & 0 & -1 & \cdots & -1 & -1 & \cdots & -1 \\
 \vdots & \vdots & & \vdots & & & & \vdots & & & & \vdots & & & \vdots & & & \vdots & & & \vdots \\
 -1 & -1 & \cdots & -1 & -1 & -1 & \cdots & -1 & -1 & -1 & \cdots & -1 & 0 & \cdots & \lambda & -1 & \cdots & -1 & -1 & \cdots & -1 \\
 -1 & -1 & \cdots & -1 & -1 & -1 & \cdots & -1 & -1 & -1 & \cdots & -1 & -1 & \cdots & -1 & \lambda & \cdots & 0 & -1 & \cdots & -1 \\
 \vdots & \vdots & & \vdots & & & & \vdots & & & & \vdots & & & \vdots & & & \vdots & & & \vdots \\
 -1 & -1 & \cdots & -1 & -1 & -1 & \cdots & -1 & -1 & -1 & \cdots & -1 & -1 & \cdots & -1 & 0 & \cdots & \lambda & -1 & \cdots & -1 \\
 -1 & -1 & \cdots & -1 & -1 & -1 & \cdots & -1 & -1 & -1 & \cdots & -1 & -1 & \cdots & -1 & -1 & \cdots & -1 & \lambda & \cdots & -1 \\
 \vdots & \vdots & & \vdots & & & & \vdots & & & & \vdots & & & \vdots & & & \vdots & & & \vdots \\
 -1 & -1 & \cdots & -1 & -1 & -1 & \cdots & -1 & -1 & -1 & \cdots & \lambda
 \end{vmatrix}$$

By elementary calculations on the above determinant, we get the result. □

Here we give some examples in Table 1.

$p, (m_1, k_1), (m_2, k_2), \dots, (m_l, k_l)$	The characteristic equations of graphs K_{a_p}
$9, (2,1), (3,1), (4,1)$	$\lambda^6(\lambda^3 - 26\lambda - 48)$
$11, (2,2), (3,1), (4,1)$	$\lambda^7(\lambda^4 - 44\lambda^2 - 152\lambda - 144)$
$12, (2,1), (3,1), (4,1)$	$\lambda^6(\lambda^6 - 56\lambda^4 - 260\lambda^3 - 477\lambda^2 - 392\lambda - 120)$
$15, (3,2), (4,1), (5,1)$	$\lambda^{11}(\lambda^4 - 83\lambda^2 - 402\lambda - 540)$
$14, (2,1), (3,1), (4,1), (5,1)$	$\lambda^{10}(\lambda^4 - 71\lambda^2 - 308\lambda - 360)$

TABLE 1. The characteristic equations of the graphs denoted by $K_{a_p}(p, (m_1, k_1), (m_2, k_2), \dots, (m_l, k_l))$

Acknowledgement: Authors are thankful to the referees for suggestions.

REFERENCES

[1] D. M. Cvetković, M. Doob and H. Sachs, *Spectra of Graphs*, Academic Press, New York, 1980.

[2] I. Gutman and L. Pavlović, *The energy of some graphs with large number of edges*, Bull. Acad. Serbe Sci. Arts (Cl. Math. Natur.) **118** (1999), 35.

[3] P. R. Hampiholi, H. B. Walikar and B. S. Durgi, *Energy of complement of stars*, J. Indones. Math. Soc. **19(1)** (2013), 15–21.

- [4] H. B. Walikar and H. S. Ramane, *Energy of some cluster graphs*, Kragujevac J. Sci. **23** (2001), 51–62.
- [5] P. R. Hampiholi and B. S. Durgi, *On the spectra and energy of some cluster graphs*, Int. J. Math. Sci. Engg. Appl. **7(2)** (2013), 219–228.

¹DEPARTMENT OF MASTER OF COMPUTER APPLICATIONS,
GOGTE INSTITUTE OF TECHNOLOGY,
BELGAUM - 590008,
INDIA
E-mail address: prhampi@yahoo.in

²DEPARTMENT OF MATHEMATICS,
KLE COLLEGE OF ENGINEERING AND TECHNOLOGY,
BELGAUM - 590008,
INDIA
E-mail address: durgibs@yahoo.com