

## INTEGRAL BOUNDARY VALUE PROBLEMS FOR IMPLICIT FRACTIONAL DIFFERENTIAL EQUATIONS INVOLVING HADAMARD AND CAPUTO-HADAMARD FRACTIONAL DERIVATIVES

P. KARTHIKEYAN<sup>1</sup> AND R. ARUL<sup>1</sup>

ABSTRACT. In this paper, we examine the existence and uniqueness of integral boundary value problem for implicit fractional differential equations (IFDE's) involving Hadamard and Caputo-Hadamard fractional derivative. We prove the existence and uniqueness results by utilizing Banach and Schauder's fixed point theorem. Finally, examples are introduced of our results.

### 1. INTRODUCTION

FDE's are considered to be a different model to integer differential equations. It has been proved by applying importance in the modeling of various fields of physical sciences, medicine, electronics and wave transformation [8, 16, 21, 23, 26]. The dominant techniques are the method of introducing a parameter for solving an implicit differential equations. In past three years, the most of research paper to developed existence and uniqueness of implicit FDE's involving various derivatives like the Caputo, Riemann-Liouville, Caputo-Hadamard, Hadamard, Hilfer-Hadamard fractional derivatives etc., (see [4–7, 9, 14, 15, 19, 20, 24]).

Caputo Hadamard fractional derivatives were studied in [12] by the authors F. Jarad, T. Abdeljawad and D. Baleanu, where a Caputo-type modification for Hadamard derivatives was introduced and studied. Later, more properties of Hadamard fractional derivatives were investigated in [1, 2, 10, 13].

---

*Key words and phrases.* Implicit fractional differential equations, Hadamard fractional operators, boundary condition, fixed point theorem, existence and uniqueness.

2010 *Mathematics Subject Classification.* Primary: 34A09. Secondary: 34A08, 26A33, 34A12, 34B10.

DOI 10.46793/KgJMat2103.331K

*Received:* August 01, 2018.

*Accepted:* January 08, 2019.

The applications of Hadamard fractional differential equations in mathematical physics could be found in [11, 17, 18, 22, 25]. In [3] the authors have studied Hilfer-Hadamard FDE's with variable-order fractional integral and fractional derivative. Motivated by the above cited work, we studies the solutions of existence and uniqueness results to the following implicit fractional differential equations with integral boundary conditions of the form

$$(1.1) \quad {}^{\mathcal{H}}\mathcal{D}^{\vartheta}x(t) = g(t, x(t), {}^{\mathcal{H}}\mathcal{D}^{\vartheta}x(t)), \quad t \in \mathcal{J} := (b, \mathcal{T}),$$

$$(1.2) \quad x(b) = 0, x(\mathcal{T}) = \lambda \int_0^{\sigma} x(s)ds, \quad b < \sigma < \mathcal{T}, \lambda \in \mathbb{R},$$

where  ${}^{\mathcal{H}}\mathcal{D}^{\vartheta}$  is the Hadamard fractional derivative of order  $1 < \vartheta \leq 2$ ,

$$(1.3) \quad {}^{\text{c}\mathcal{H}}\mathcal{D}^{\vartheta}x(t) = g(t, x(t), {}^{\text{c}\mathcal{H}}\mathcal{D}^{\vartheta}x(t)), \quad t \in \mathcal{J} := [b, \mathcal{T}],$$

$$(1.4) \quad x(b) = 0, x(\mathcal{T}) = \lambda \int_0^{\sigma} x(s)ds, \quad b < \sigma < \mathcal{T}, \lambda \in \mathbb{R},$$

where  ${}^{\text{c}\mathcal{H}}\mathcal{D}^{\vartheta}$  is the Caputo-Hadamard fractional derivative of order  $1 < \vartheta \leq 2$  and  $g : \mathcal{J} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  is a continuous function.

In this paper, Section 2, has definitions and some of the most important basic concepts of the fractional calculus. In Section 3, existence and uniqueness of solutions for integral boundary conditions of implicit fractional differential equations involving Hadamard fractional derivative and Caputo-Hadamard fractional derivatives are proved by utilizing Banach and Schauder's fixed point theorems. In Section 4, an illustrative examples are provided to explain of the results of the problem (1.1)–(1.4).

## 2. BASIC RESULTS

In this section, the some most important basic concepts, definitions and some supporting results are used in this paper. By  $\mathcal{C}(\mathcal{J}, \mathbb{R})$  we denote the Banach space of all continuous functions form  $\mathcal{J}$  into  $\mathbb{R}$  with the norm  $\|x\|_{\infty} = \sup\{|x(t)| : t \in \mathcal{J}\}$ .

**Definition 2.1** ([15]). The derivative of fractional order  $\vartheta > 0$  of a function  $g : (0, \infty) \rightarrow \mathbb{R}$  is given by

$$\mathcal{D}_{0+}^{\vartheta}x(t) = \frac{1}{\Gamma(n - \vartheta)} \left( \frac{d}{dt} \right)^n \int_0^t \frac{g(s)}{(t - s)^{\vartheta - n + 1}} ds,$$

where  $n = [\vartheta] + 1$ , provided the right side is pointwise defined on  $(0, \infty)$ .

**Definition 2.2** ([15]). The **Hadamard fractional integral** of  $g$  is defined by

$${}^{\mathcal{H}}\mathcal{I}^{\vartheta}x(t) = \frac{1}{\Gamma(\vartheta)} \int_b^t \left( \log \frac{t}{s} \right)^{\vartheta - 1} \frac{g(s)}{s} ds, \quad \vartheta > 0.$$

**Definition 2.3** ([15]). The **Hadamard fractional derivative** of  $g$  is continuous function and further,  $\log(\cdot) = \log_e(\cdot)$  is defined as

$${}^{\mathcal{H}}\mathcal{D}^{\vartheta}x(t) = \frac{1}{\Gamma(n - \vartheta)} \left( t \frac{d}{dt} \right)^n \int_b^t \left( \log \frac{t}{s} \right)^{n - \vartheta - 1} \frac{g(s)}{s} ds,$$

where  $n - 1 < \vartheta < n$ ,  $n = [\vartheta] + 1$  and  $[\vartheta]$  denotes the integer part of the real number  $\vartheta$ .

**Definition 2.4** ([12]). For at least  $n$ -times differentiable function  $g$ , the **Caputo-Hadamard fractional derivative** of order  $\vartheta$  is defined as

$${}^{\mathcal{C}\mathcal{H}}\mathcal{D}^\vartheta x(t) = \frac{1}{\Gamma(n - \vartheta)} \int_a^t \left(\ln \frac{t}{s}\right)^{n-\vartheta-1} \delta^n \frac{g(s)}{s} ds.$$

**Lemma 2.1** (Hadamard fractional derivative). *Let  $v \in \mathcal{C}([b, \mathcal{T}], \mathbb{R})$  and  $x \in \mathcal{C}_s^2([b, \mathcal{T}], \mathbb{R})$ . Then*

$$(2.1) \quad \begin{aligned} &{}^{\mathcal{H}}\mathcal{D}^\vartheta x(t) = v(t), \quad t \in \mathcal{J} := [b, \mathcal{T}], \\ &x(b) = 0, x(\mathcal{T}) = \lambda \int_0^\sigma x(s) ds, \quad b < \sigma < \mathcal{T}, \lambda \in \mathbb{R}, \end{aligned}$$

is equivalent to the integral equation given by

$$(2.2) \quad \begin{aligned} x(t) = &\frac{1}{\Gamma(\vartheta)} \int_b^t \left(\ln \frac{t}{s}\right)^{\vartheta-1} \frac{v(s)}{s} ds + \frac{\left(\ln \frac{t}{s}\right)^{\vartheta-1}}{\Gamma(\vartheta) \left[ \left(\ln \frac{\mathcal{T}}{s}\right)^{\vartheta-1} - \lambda \left[ \int_b^\sigma \left(\ln \frac{s}{b}\right)^{\vartheta-1} ds \right] \right]} \\ &\times \left[ \lambda \int_b^\sigma \int_b^s \left(\ln \frac{s}{r}\right)^{\vartheta-1} \frac{v(r)}{r} dr ds - \int_b^\mathcal{T} \left(\ln \frac{\mathcal{T}}{s}\right)^{\vartheta-1} \frac{v(s)}{s} ds \right] \end{aligned}$$

**Lemma 2.2** (Caputo-Hadamard fractional derivative). *Let  $v \in \mathcal{C}([b, \mathcal{T}], \mathbb{R})$  and  $x \in \mathcal{C}_s^2([b, \mathcal{T}], \mathbb{R})$ .*

$$(2.3) \quad \begin{aligned} &{}^{\mathcal{C}\mathcal{H}}\mathcal{D}^\vartheta x(t) = v(t), \quad t \in \mathcal{J} := [b, \mathcal{T}], \\ &x(b) = 0, x(\mathcal{T}) = \lambda \int_0^\sigma x(s) ds, \quad b < \sigma < \mathcal{T}, \lambda \in \mathbb{R}, \end{aligned}$$

is equivalent to the integral equation given by

$$(2.4) \quad \begin{aligned} x(t) = &\frac{1}{\Gamma(\vartheta)} \int_b^t \left(\ln \frac{t}{s}\right)^{\vartheta-1} \frac{v(s)}{s} ds + \frac{\left(\ln \frac{t}{s}\right)}{\Gamma(\vartheta) \left[ \left(\ln \frac{\mathcal{T}}{s}\right) - \lambda \left[ \sigma \left(\ln \frac{\sigma}{b} - 1\right) + b \right] \right]} \\ &\times \left[ \lambda \int_b^\sigma \int_b^s \left(\ln \frac{s}{r}\right)^{\vartheta-1} \frac{v(r)}{r} dr ds - \int_b^\mathcal{T} \left(\ln \frac{\mathcal{T}}{s}\right)^{\vartheta-1} \frac{v(s)}{s} ds \right]. \end{aligned}$$

**Lemma 2.3** (Nonlinear alternative of Lerary-Schauder type, [7]). *Let  $\mathcal{B}$  be a Banach space,  $\mathcal{C}$  a closed, convex subset of  $\mathcal{B}$ ,  $\mathcal{U}$  an open subset of  $\mathcal{C}$  and  $0 \in \mathcal{U}$ . Suppose that  $F : \overline{\mathcal{U}} \rightarrow \mathcal{C}$  is a continuous, compact map. Then either (i)  $F$  has a fixed point in  $\overline{\mathcal{U}}$ , or (ii) there is a  $u \in \partial\mathcal{U}$  and  $\lambda \in (0, 1)$ , with  $u = \lambda F(u)$ .*

### 3. MAIN RESULTS

To prove the existence and uniqueness results we need the following assumptions.

*Assumption 3.1.* The function  $g : \mathcal{J} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  is a continuous function.

*Assumption 3.2.* There exists constants  $K_g > 0$  and  $0 < L_g < 1$  such that

$$|g(t, u, v) - g(t, u_1, v_1)| \leq K_g|u - u_1| + L_g|v - v_1|, \quad \text{for any } u, v, u_1, v_1 \in \mathbb{R}.$$

*Assumption 3.3.* There exist a continuous nondecreasing function  $\varphi$  on  $[0, \infty) \rightarrow (0, \infty)$  and a function  $p(t) \in \mathcal{C}^1([b, \mathcal{J}], \mathbb{R}^+)$  such that

$$\|g(t, u, v)\| \leq p(t)\varphi(\|u\| + \|v\|).$$

The integral boundary conditions for implicit fractional differential equations with Hadamard fractional derivative (1.1)–(1.2) is equivalent to the integral equation

$$\begin{aligned} x(t) = & \frac{1}{\Gamma(\vartheta)} \int_b^t \left(\ln \frac{t}{s}\right)^{\vartheta-1} \frac{g(s, x(s), {}^{\mathcal{H}}\mathcal{D}^\vartheta x(s))}{s} ds + \frac{\left(\ln \frac{t}{b}\right)^{\vartheta-1}}{\Gamma(\vartheta) \left[\left(\ln \frac{\mathcal{J}}{b}\right)^{\vartheta-1} - \lambda N_1\right]} \\ & \times \left[ \lambda \int_b^\sigma \int_b^s \left(\ln \frac{s}{r}\right)^{\vartheta-1} \frac{g(r, x(r), {}^{\mathcal{H}}\mathcal{D}^\vartheta x(r))}{r} dr ds \right. \\ & \left. - \int_b^{\mathcal{J}} \left(\ln \frac{\mathcal{J}}{s}\right)^{\vartheta-1} \frac{g(s, x(s), {}^{\mathcal{H}}\mathcal{D}^\vartheta x(s))}{s} ds \right], \end{aligned}$$

where  $N_1 = \int_b^\sigma \left(\ln \frac{s}{b}\right)^{\vartheta-1} ds$ .

The integral boundary conditions for implicit fractional differential equations with Caputo-Hadamard fractional derivative (1.3)–(1.4) is equivalent to the integral equation

$$\begin{aligned} x(t) = & \frac{1}{\Gamma(\vartheta)} \int_b^t \left(\ln \frac{t}{s}\right)^{\vartheta-1} \frac{g(s, x(s), {}^{\text{CH}}\mathcal{D}^\vartheta x(s))}{s} ds + \frac{\left(\ln \frac{t}{b}\right)}{\Gamma(\vartheta) \left[\left(\ln \frac{\mathcal{J}}{b}\right) - \lambda N_2\right]} \\ & \times \left[ \lambda \int_b^\sigma \int_r^s \left(\ln \frac{s}{r}\right)^{\vartheta-1} \frac{g(r, x(r), {}^{\text{CH}}\mathcal{D}^\vartheta x(r))}{r} dr ds \right. \\ & \left. - \int_b^{\mathcal{J}} \left(\ln \frac{\mathcal{J}}{s}\right)^{\vartheta-1} \frac{g(s, x(s), {}^{\text{CH}}\mathcal{D}^\vartheta x(s))}{s} ds \right] \end{aligned}$$

or

$$x(t) = I^\vartheta f(s) + \left( \frac{\left(\ln \frac{t}{b}\right)}{\Gamma(\vartheta) \left[\left(\ln \frac{\mathcal{J}}{b}\right) - \lambda N_2\right]} \right) \left[ \lambda \int_b^\sigma I^\vartheta f_1(r) ds - I^\vartheta f_2(s) \right],$$

where  $N_2 = \sigma \left(\ln \frac{\sigma}{b} - 1\right) + b$  and  $f, f_1, f_2 \in \mathcal{C}(\mathcal{J}, \mathbb{R})$  satisfies the functional equations

$$\begin{aligned} f(s) &= g(s, I^\vartheta f(s), f(s)), \\ f_1(r) &= g(r, I^\vartheta f_1(r), f_1(r)), \\ f_2(s) &= g(s, I^\vartheta f_2(s), f_2(r)), \\ I^\vartheta f(s) &= \frac{1}{\Gamma(\vartheta)} \int_b^t \left(\ln \frac{t}{s}\right)^{\vartheta-1} \frac{g(s, x(s), {}^{\text{CH}}\mathcal{D}^\vartheta x(s))}{s} ds, \end{aligned}$$

$$I^\vartheta f_1(r) = \int_b^s \left(\ln \frac{s}{r}\right)^{\vartheta-1} \frac{g(r, x(r), {}^{\mathcal{C}\mathcal{H}}\mathcal{D}^\vartheta x(r))}{r} dr,$$

$$I^\vartheta f_2(s) = \int_b^{\mathcal{J}} \left(\ln \frac{\mathcal{J}}{s}\right)^{\vartheta-1} \frac{g(s, x(s), {}^{\mathcal{C}\mathcal{H}}\mathcal{D}^\vartheta x(s))}{s} ds.$$

**Theorem 3.1.** Assume that assumptions 3.1 and 3.2 hold. If

$$\left[ \frac{1}{\Gamma(\vartheta+1)} \left(\ln \frac{\mathcal{J}}{b}\right)^\vartheta + \frac{\left(\ln \frac{\mathcal{J}}{b}\right)^{2\vartheta-1}}{\Gamma(\vartheta+1) \left| \left(\ln \frac{\mathcal{J}}{b}\right)^{\vartheta-1} - \lambda N_1 \right|} (|\lambda|(\sigma-b) - 1) \right] \frac{K_g}{(1-L_g)} < 1,$$

then there exists a unique solution for (1.1)–(1.2) on  $\mathcal{J} := [b, \mathcal{J}]$ .

*Proof.* Let  $B_r = \{x \in \mathcal{C}([b, \mathcal{J}], \mathbb{R}) : \|x\| \leq r\}$ . Consider the operator  $\mathcal{H} : \mathcal{C}([b, \mathcal{J}], \mathbb{R}) \rightarrow \mathcal{C}([b, \mathcal{J}], \mathbb{R})$  defined by

$$(3.1) \quad \mathcal{H}(x)(t) = I^\vartheta f(s) + \left( \frac{\left(\ln \frac{t}{b}\right)^{\vartheta-1}}{\Gamma(\vartheta) \left[ \left(\ln \frac{\mathcal{J}}{b}\right)^{\vartheta-1} - \lambda N_1 \right]} \right) \left( \lambda \int_b^\sigma I^\vartheta f_1(r) ds - I^\vartheta f_2(s) \right),$$

where  $f, f_1, f_2 \in \mathcal{C}(\mathcal{J}, \mathbb{R})$  satisfies the functional equations

$$f(s) = f(s, I^\vartheta f(s), f(s)),$$

$$f_1(r) = f(r, I^\vartheta f_1(r), f_1(r)),$$

$$f_2(s) = f(s, I^\vartheta f_2(s), f_2(s)),$$

where  $N_1 = \int_b^\sigma \left(\ln \frac{s}{b}\right)^{\vartheta-1} ds$  and

$$I^\vartheta f(s) = \frac{1}{\Gamma(\vartheta)} \int_b^t \left(\ln \frac{t}{s}\right)^{\vartheta-1} \frac{g(s, x(s), {}^{\mathcal{H}}\mathcal{D}^\vartheta x(s))}{s} ds,$$

$$I^\vartheta f_1(r) = \int_b^s \left(\ln \frac{s}{r}\right)^{\vartheta-1} \frac{g(r, x(r), {}^{\mathcal{H}}\mathcal{D}^\vartheta x(r))}{r} dr,$$

$$I^\vartheta f_2(s) = \int_b^{\mathcal{J}} \left(\ln \frac{\mathcal{J}}{s}\right)^{\vartheta-1} \frac{g(s, x(s), {}^{\mathcal{H}}\mathcal{D}^\vartheta x(s))}{s} ds.$$

Clearly, the fixed point of operator  $\mathcal{H}$  is solution of problem (1.1)–(1.2). Let  $x_1, x_2 \in \mathcal{C}([b, \mathcal{J}], \mathbb{R})$ . Then

$$(\mathcal{H}x_1)(t) - (\mathcal{H}x_2)(t) = \frac{1}{\Gamma(\vartheta)} \int_b^t \left(\ln \frac{t}{s}\right)^{\vartheta-1} \frac{f(s) - h(s)}{s} ds$$

$$+ \left( \frac{\left(\ln \frac{t}{b}\right)^{\vartheta-1}}{\Gamma(\vartheta) \left[ \left(\ln \frac{\mathcal{J}}{b}\right)^{\vartheta-1} - \lambda N_1 \right]} \right)$$

$$\times \left[ \lambda \int_b^\sigma \int_b^s \left(\ln \frac{s}{r}\right)^{\vartheta-1} \frac{f(r) - h(r)}{r} dr ds \right]$$

$$- \int_b^{\mathcal{J}} \left( \ln \frac{\mathcal{J}}{b} \right)^{\vartheta-1} \frac{f(s) - h(s)}{s} ds \Big],$$

where  $f(s), h(s), f(r), h(r) \in \mathcal{C}([b, \mathcal{J}], \mathbb{R})$  are such that

$$\begin{aligned} f(s) &= f(s, x_1(s), f(s)), & f(r) &= f(r, x_2(r), f(r)), \\ h(s) &= h(s, x_1(s), h(s)), & h(r) &= h(r, x_2(r), h(r)). \end{aligned}$$

Now,

$$\begin{aligned} |(\mathcal{H}x_1)(t) - (\mathcal{H}x_2)(t)| &\leq \frac{1}{\Gamma(\vartheta)} \int_b^t \left( \ln \frac{t}{s} \right)^{\vartheta-1} \frac{|f(s) - h(s)|}{s} ds \\ &\quad + \left( \frac{\left( \ln \frac{t}{s} \right)^{\vartheta-1}}{\Gamma(\vartheta) \left| \left( \ln \frac{\mathcal{J}}{s} \right)^{\vartheta-1} - \lambda N_1 \right|} \right) \\ &\quad \times \left[ |\lambda| \int_b^\sigma \int_b^s \left( \ln \frac{s}{r} \right)^{\vartheta-1} \frac{|f(r) - h(r)|}{r} dr ds \right. \\ &\quad \left. - \int_b^{\mathcal{J}} \left( \ln \frac{\mathcal{J}}{b} \right)^{\vartheta-1} \frac{|f(s) - h(s)|}{s} ds \right], \end{aligned} \tag{3.2}$$

and, by Assumption 3.2, we have

$$\begin{aligned} |(f(s) - h(s))| &= |g(s, x_1(s), f(s)) - g(s, x_2(s), h(s))|, \\ |(f(s) - h(s))| &\leq K_g |x_1(s) - x_2(s)| + L_g |x_1(s) - x_2(s)| \leq \frac{K_g}{1 - L_g} |x_1(s) - x_2(s)|, \\ |(f(s) - h(s))| &\leq \frac{K_g}{1 - L_g} |x_1(s) - x_2(s)|. \end{aligned}$$

Similarly,

$$|(f(r) - h(r))| \leq \frac{K_g}{1 - L_g} |x_1(r) - x_2(r)|.$$

The equation (3.2) implies

$$\begin{aligned} |(\mathcal{H}x_1)(t) - (\mathcal{H}x_2)(t)| &\leq \frac{1}{\Gamma(\vartheta + 1)} \left( \frac{K_g}{1 - L_g} \right) \|x_1 - x_2\| \left( \ln \frac{\mathcal{J}}{b} \right)^\vartheta \\ &\quad + \frac{\left( \ln \frac{\mathcal{J}}{b} \right)^{2\vartheta-1}}{\Gamma(\vartheta + 1) \left| \left( \ln \frac{\mathcal{J}}{b} \right)^{\vartheta-1} - \lambda N_1 \right|} \\ &\quad \times \left( (|\lambda|(\sigma - b) - 1) \left( \frac{K_g}{1 - L_g} \right) \right) \|x_1 - x_2\| \end{aligned}$$

$$\begin{aligned} &\leq \left( \frac{1}{\Gamma(\vartheta + 1)} \left( \ln \frac{\mathcal{J}}{b} \right)^\vartheta + \frac{\left( \ln \frac{\mathcal{J}}{b} \right)^{2\vartheta-1}}{\Gamma(\vartheta + 1) \left| \left( \ln \frac{\mathcal{J}}{b} \right)^{\vartheta-1} - \lambda N_1 \right|} \right) \\ &\quad \times (|\lambda|(\sigma - b) - 1) \left( \frac{K_g}{1 - L_g} \right) \|x_1 - x_2\|_\infty. \end{aligned}$$

Thus,

$$\begin{aligned} |(\mathcal{H}x_1)(t) - (\mathcal{H}x_2)(t)| &\leq \left( \frac{1}{\Gamma(\vartheta + 1)} \left( \ln \frac{\mathcal{J}}{b} \right)^\vartheta + \frac{\left( \ln \frac{\mathcal{J}}{b} \right)^{2\vartheta-1}}{\Gamma(\vartheta + 1) \left| \left( \ln \frac{\mathcal{J}}{b} \right)^{\vartheta-1} - \lambda N_1 \right|} \right) \\ &\quad \times (|\lambda|(\sigma - b) - 1) \left( \frac{K_g}{1 - L_g} \right) \|x_1 - x_2\|_\infty. \end{aligned}$$

By (3.1), the operator  $\mathcal{H}$  is continuous. Hence, by Banach's contraction principle,  $\mathcal{H}$  has a unique fixed point which is a unique solution of the problem (1.1)–(1.2) on  $\mathcal{J} := [b, \mathcal{J}]$ .  $\square$

**Theorem 3.2.** *Assume that assumptions 3.1 and 3.2 hold. If*

$$\left[ \frac{1}{\Gamma(\vartheta + 1)} \left( \ln \frac{\mathcal{J}}{b} \right)^\vartheta + \frac{\left( \ln \frac{\mathcal{J}}{b} \right)^{\vartheta+1}}{\Gamma(\vartheta + 1) \left| \left( \ln \frac{\mathcal{J}}{b} \right) - \lambda N_2 \right|} (|\lambda|(\sigma - b) - 1) \right] \left( \frac{K_g}{1 - L_g} \right) < 1,$$

*then there exists a unique solution for (1.3)–(1.4) on  $\mathcal{J} := [b, \mathcal{J}]$ .*

The proof of Theorem 3.2 is similar to the Theorem 3.1.

**Theorem 3.3.** *Assume that assumptions 3.1 and 3.3 hold. Then there is at least one solution for the problem (1.1)–(1.2) on  $\mathcal{J} := [b, \mathcal{J}]$ .*

*Proof. Step 1.* Show that  $\mathcal{H}$  maps bounded sets (balls) into bounded sets in  $\mathcal{C}([b, \mathcal{J}], \mathbb{R})$ .

For a positive number  $r_1$ , let  $B_{r_1} = \{x \in \mathcal{C}([b, \mathcal{J}], \mathbb{R}) : \|\mathcal{Z}^*\| \leq r_1\}$  be a bounded ball in  $\mathcal{C}([b, \mathcal{J}], \mathbb{R})$ , where

$$\|\mathcal{Z}^*\| = \sup_{t \in [b, \mathcal{J}]} (|x| + |g|).$$

Then

$$\begin{aligned} |\mathcal{H}(x)(t)| &\leq \frac{1}{\Gamma(\vartheta)} \int_b^t \left( \ln \frac{t}{s} \right)^{\vartheta-1} \frac{|g(s, x(s), {}^c \mathcal{D}^\vartheta x(s))|}{s} ds + \frac{\left( \ln \frac{t}{b} \right)^{\vartheta-1}}{\Gamma(\vartheta) \left| \left( \ln \frac{\mathcal{J}}{b} \right)^{\vartheta-1} - \lambda N_1 \right|} \\ &\quad \times \left[ |\lambda| \int_b^\sigma \int_b^s \left( \ln \frac{s}{r} \right)^{\vartheta-1} \frac{|g(r, x(r), {}^c \mathcal{D}^\vartheta x(r))|}{r} dr ds \right. \\ &\quad \left. - \int_b^\mathcal{J} \left( \ln \frac{\mathcal{J}}{s} \right)^{\vartheta-1} \frac{|g(s, x(s), {}^c \mathcal{D}^\vartheta x(s))|}{s} ds \right] \end{aligned}$$

$$\begin{aligned} &\leq \frac{1}{\Gamma(\vartheta)} \int_b^t \left(\ln \frac{t}{s}\right)^{\vartheta-1} \frac{\varphi(\|\mathcal{Z}^*\|)\|p\|}{s} ds + \frac{\left(\ln \frac{t}{b}\right)^{\vartheta-1}}{\Gamma(\vartheta) \left| \left(\ln \frac{\mathcal{T}}{b}\right)^{\vartheta-1} - \lambda N_1 \right|} \\ &\quad \times \left[ |\lambda| \int_b^\sigma \int_b^s \left(\ln \frac{s}{r}\right)^{\vartheta-1} \frac{\varphi(\|\mathcal{Z}^*\|)\|p\|}{r} dr ds \right. \\ &\quad \left. - \int_b^{\mathcal{T}} \left(\ln \frac{\mathcal{T}}{s}\right)^{\vartheta-1} \frac{\varphi(\|\mathcal{Z}^*\|)\|p\|}{s} ds \right], \end{aligned}$$

i.e.,

$$\begin{aligned} |\mathcal{H}(x)(t)| &\leq \left( \frac{1}{\Gamma(\vartheta+1)} \left(\ln \frac{\mathcal{T}}{s}\right)^\vartheta + \frac{\left(\ln \frac{\mathcal{T}}{b}\right)^{2\vartheta-1}}{\Gamma(\vartheta+1) \left| \left(\ln \frac{\mathcal{T}}{b}\right)^{\vartheta-1} - \lambda N_1 \right|} \right) \\ &\quad \times (|\lambda|(\sigma - b) - 1) \varphi(r)\|p\|. \end{aligned}$$

**Step 2.** Show that  $\mathcal{H}$  maps bounded sets (balls) into equicontinuous sets in  $\mathcal{C}([b, \mathcal{T}], \mathbb{R})$ . Let  $\mu_1, \mu_2 \in [b, \mathcal{T}]$ ,  $\mu_1 < \mu_2$ . Then, we have

$$\begin{aligned} \|\mathcal{H}(x)(\mu_1) - \mathcal{H}(x)(\mu_2)\| &\leq \frac{1}{\Gamma(\vartheta)} \left[ \int_b^{\mu_1} \left[ \left(\ln \frac{\mu_2}{r}\right)^{\vartheta-1} - \left(\ln \frac{\mu_1}{r}\right)^{\vartheta-1} \right] \frac{\varphi(\|\mathcal{Z}^*\|)\|p\|}{s} ds \right. \\ &\quad \left. + \int_{\mu_1}^{\mu_2} \left(\ln \frac{\mu_2}{s}\right)^{\vartheta-1} \frac{\varphi(\|\mathcal{Z}^*\|)\|p\|}{s} ds \right] \\ &\quad + \frac{\left(\ln \frac{\mu_2}{b}\right)^{\vartheta-1} - \left(\ln \frac{\mu_1}{b}\right)^{\vartheta-1}}{\Gamma(\vartheta) \left| \left(\ln \frac{\mathcal{T}}{b}\right)^{\vartheta-1} - \lambda N_1 \right|} \\ &\quad \times \left[ |\lambda| \int_b^\sigma \int_b^s \left(\ln \frac{s}{r}\right)^{\vartheta-1} \frac{\varphi(\|\mathcal{Z}^*\|)\|p\|}{r} dr ds \right. \\ &\quad \left. - \int_b^{\mathcal{T}} \left(\ln \frac{\mathcal{T}}{s}\right)^{\vartheta-1} \frac{\varphi(\|\mathcal{Z}^*\|)\|p\|}{s} ds \right]. \end{aligned}$$

Obviously, the right-hand side of the above inequality tends to zero independently of  $u, v \in B_{r_1}$  as  $\mu_2 - \mu_1 \rightarrow 0$ . As  $\mathcal{H}$  satisfies the above assumptions, therefore, by the Arzela-Ascoli theorem, it follows that  $\mathcal{H} : \mathcal{C}([b, \mathcal{T}], \mathbb{R}) \rightarrow \mathcal{C}([b, \mathcal{T}], \mathbb{R})$  is completely continuous. Let  $x$  be a solution. Then, for  $t \in [b, \mathcal{T}]$  and following the similar computations as in the first step, we have

$$\begin{aligned} |x(t)| &= \lambda |\mathcal{H}(x)(t)| \\ &\leq \left( \frac{1}{\Gamma(\vartheta+1)} \left(\ln \frac{\mathcal{T}}{s}\right)^\vartheta + \frac{\left(\ln \frac{\mathcal{T}}{b}\right)^{2\vartheta-1}}{\Gamma(\vartheta+1) \left| \left(\ln \frac{\mathcal{T}}{b}\right)^{\vartheta-1} - \lambda N_1 \right|} \right) (|\lambda|(\sigma - b) - 1) \end{aligned}$$



$$\times \varphi(\|x\|)\|p\|.$$

Consequently, we have

$$\frac{\|x(t)\|}{\left(\frac{1}{\Gamma(\vartheta+1)} \left(\ln \frac{\mathcal{T}}{s}\right)^\vartheta + \frac{\left(\ln \frac{\mathcal{T}}{b}\right)^{2\vartheta-1}}{\Gamma(\vartheta+1) \left| \left(\ln \frac{\mathcal{T}}{b}\right)^{\vartheta-1} - \lambda N_1 \right|} (|\lambda|(\sigma - b) - 1)\right) \varphi(\|x\|)\|p\|} \leq 1.$$

There exists  $\mathcal{M}^*$  such that  $\|x\| \neq \mathcal{M}^*$ . Let us set

$$U = \{x \in \mathcal{C}([b, \mathcal{T}], \mathbb{R}) : \|x\| < \mathcal{M}^*\}.$$

Note that the operator  $\mathcal{H} : \bar{U} \rightarrow \mathcal{C}([b, \mathcal{T}], \mathbb{R})$  is continuous and completely continuous. From the choice of  $\mathcal{U}$ , there is no  $x \in \partial\mathcal{U}$  such that  $x = \lambda\mathcal{H}x$  for some  $0 \leq \lambda \leq 1$ . Consequently, by the nonlinear alternative of Lerary-Schauder type (Lemma 2.3), we deduce that  $\mathcal{H}$  has fixed point  $x \in \bar{U}$  which is a solution of the problem (1.1)–(1.2).  $\square$

**Theorem 3.4.** *Assume that assumptions 3.1 and 3.3 hold and there exists a constant  $\mathcal{M}^* > 0$ , such that*

$$\mathcal{M}^* > \left(\frac{1}{\Gamma(\vartheta + 1)} \left(\ln \frac{\mathcal{T}}{s}\right)^\vartheta + \frac{\left(\ln \frac{\mathcal{T}}{b}\right)^{\vartheta+1}}{\Gamma(\vartheta + 1) \left| \left(\ln \frac{\mathcal{T}}{b}\right) - \lambda N_1 \right|} (|\lambda|(\sigma - b) - 1)\right) \|p\| \varphi(\|x\|).$$

Then, there is at least one solution for the problem (1.3)–(1.4) on  $\mathcal{J} =: [b, \mathcal{T}]$ .

The proof of Theorem 3.4 is similar to the Theorem 3.3.

#### 4. EXAMPLES

In this section, some examples are introduced for Hadamard and Caputo-Hadamard fractional derivatives of implicit fractional differential equations with integral boundary conditions.

*Example 4.1.* Consider the implicit Hadamard FDE's with three point integral boundary conditions of the form

$$(4.1) \quad {}^{\mathcal{H}}\mathcal{D}^{\frac{10}{7}} x(t) = \frac{|x|}{(t + 6)^2 (|1 + |x| + |{}^{\mathcal{H}}\mathcal{D}^{\frac{10}{7}} x(t)|)}, \quad 1 < \vartheta \leq 2,$$

$$(4.2) \quad x(1) = 0, \quad x(b) = \lambda \int_b^\sigma x(s) ds.$$

Here  $\vartheta = \frac{10}{7}$ ,

$$g(t, x(t), {}^{\mathcal{H}}\mathcal{D}^\vartheta x(t)) = \frac{|x|}{(t + 6)^2 (|1 + |x| + |{}^{\mathcal{H}}\mathcal{D}^{\frac{10}{7}} x(t)|)},$$

$\sigma = 3, \lambda = 5$ . Hence, the Assumption 3.2 holds, with  $K_g = L_g = \frac{1}{49}$  and we will check that

$$\left[ \frac{1}{\Gamma(\vartheta + 1)} \left( \ln \frac{\mathcal{J}}{b} \right)^\vartheta + \frac{\left( \ln \frac{\mathcal{J}}{b} \right)^{2\vartheta-1}}{\Gamma(\vartheta + 1) \left| \left( \ln \frac{\mathcal{J}}{b} \right)^{\vartheta-1} - \lambda N_1 \right|} (|\lambda|(\sigma - b) - 1) \right] \frac{K_g}{(1 - L_g)} < 1.$$

Thus, the Theorem 3.1 is satisfied and shows that the problem (4.1)–(4.2) has a unique solution on  $\mathcal{J} =: [b, \mathcal{J}]$ .

*Example 4.2.* Consider the implicit Caputo-Hadamard FDE's with three point integral boundary conditions of the form

$$(4.3) \quad {}^{\text{CH}}\mathcal{D}^{\frac{10}{7}} x(t) = \frac{|x|}{(t+6)^2(|1+|x|+|{}^{\text{CH}}\mathcal{D}^{\frac{10}{7}} x(t)|)}, \quad 1 < \vartheta \leq 2,$$

$$(4.4) \quad x(1) = 0, x(b) = \lambda \int_b^\sigma x(s) ds.$$

Here  $\vartheta = \frac{10}{7}$ ,

$$g(t, x(t), {}^{\text{CH}}\mathcal{D}^\vartheta x(t)) = \frac{|x|}{(t+6)^2(|1+|x|+|{}^{\text{CH}}\mathcal{D}^{\frac{10}{7}} x(t)|)},$$

$\sigma = 3, \lambda = 5$ . Hence, the Assumption 3.2 holds, with  $K_g = L_g = \frac{1}{49}$  and we will check that

$$\left[ \frac{1}{\Gamma(\vartheta + 1)} \left( \ln \frac{\mathcal{J}}{b} \right)^\vartheta + \frac{\left( \ln \frac{\mathcal{J}}{b} \right)^{\vartheta+1}}{\Gamma(\vartheta + 1) \left| \left( \ln \frac{\mathcal{J}}{b} \right) - \lambda N_2 \right|} (|\lambda|(\sigma - b) - 1) \right] \frac{K_g}{(1 - L_g)} < 1.$$

Thus, the Theorem 3.2 is satisfied and shows that the problem (4.3)–(4.4) has a unique solution on  $\mathcal{J} =: [b, \mathcal{J}]$ .

## REFERENCES

- [1] Y. Adjabi, F. Jarad and T. Abdeljawad, *On generalized fractional operators and a Gronwall type Inequality with applications*, Filomat **31**(17) (2017), 5457–5473.
- [2] Y. Adjabi, F. Jarad, D. Baleanu and T. Abdeljawad, *On Cauchy problems with Caputo Hadamard fractional derivatives*, J. Comput. Anal. Appl. **21**(1) (2016), 661–681.
- [3] R. Almeida, D. Tavares and D. F. M. Torres, *The Variable-Order Fractional Calculus of Variations*, Springer Briefs in Applied Sciences and Technology, Springer, Cham, Switzerland, 2018.
- [4] A. Anguraj, P. Karthikeyan and J. J. Trujillo, *Existence of solutions to fractional mixed integro-differential equations with nonlocal initial condition*, Adv. Difference Equ. **2011** (2011), 1–12.
- [5] A. Babakhani and T. Abdeljawad, *A Caputo fractional order boundary value problem with integral boundary conditions*, J. Comput. Anal. Appl. **15**(4) (2013), 753–763.
- [6] T. D. Benavides, *An existence theorem for implicit differential equations in a Banach space*, Ann. Mat. Pura Appl. **4** (1978), 119–130.
- [7] M. Benchohra and J. E. Lazreg, *Existence results for nonlinear implicit fractional differential equations*, Surv. Math. Appl. **9** (2014), 79–92.
- [8] Z. Dahmani and L. Tabharit, *Fractional order differential equations involving Caputo derivative*, Comput. Math. Appl. **4** (2014), 40–55.

- [9] D. B. Dhaigude and S. P. Bhairat, *Local existence and uniqueness of solution for Hilfer-Hadamard fractional differential problem*, Nonlinear Dyn. Syst. Theory **18**(2) (2018), 144–153.
- [10] Y. Y. Gambo, F. Jarad, D. Baleanu and T. Abdeljawad, *On Caputo modification of the Hadamard fractional derivatives*, Adv. Difference Equ. **2014**(10) (2014), 1–12.
- [11] R. Hilfer, *Applications of Fractional Calculus in Physics*, World Scientific, Singapore, 2000.
- [12] F. Jarad, T. Abdeljawad and D. Baleanu, *Caputo-type modification of the Hadamard fractional derivatives*, Adv. Difference Equ. **2012**(142) (2012), 1–8.
- [13] F. Jarad, T. Abdeljawad and D. Baleanu, *On the generalized fractional derivatives and their Caputo modification*, J. Nonlinear Sci. Appl. **10**(5) (2017), 2607–2619.
- [14] P. Karthikeyan and R. Arul, *Existence of solutions for Hadamard fractional hybrid differential equations with impulsive and nonlocal conditions*, J. Fract. Calc. Appl. **9**(1) (2018), 232–240.
- [15] P. Karthikeyan and R. Arul, *Stability for impulsive implicit Hadamard fractional differential equations*, Malaya J. Mat. **6**(1) (2018), 28–33.
- [16] A. A. Kilbas, H. M. Srivastava and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, North-Holland Mathematics Studies **204**, Elsevier Science, Amsterdam, 2006.
- [17] A. A. Kilbas, *Hadamard-type fractional calculus*, J. Korean Math. Soc. **38** (2001), 1191–1204.
- [18] A. A. Kilbas and J. J. Trujillo, *Hadamard-type integrals as G-transforms*, Integral Transforms Spec. Funct. **14** (2003), 413–427.
- [19] N. I. Mahmudov, M. Awadalla and K. Abuassba, *Hadamard and Caputo-Hadamard fractional differential equations with three point integral boundary conditions*, Nonlinear Analysis and Differential Equations **5**(6) (2017), 271–282.
- [20] S. K. Ntouyas and J. Tariboon, *Fractional boundary value problems with multiple order of fractional derivatives and integrals*, Electron. J. Differential Equations **100** (2017), 1–18.
- [21] K. B. Oldham and J. Spanier, *The Fractional Calculus*, Academic Press, New York, London, 1974.
- [22] P. D. Phung and L. X. Truong, *Existence of solutions to three-point boundary-value problems at resonance*, Electron. J. Differential Equations **2016**(115) (2016), 1–13.
- [23] I. Podlubny, *Fractional Differential Equations*, Academic Press, San Diego, 1999.
- [24] D. Vivek, K. Kanagarajan and E. M. Elsayed, *Some existence and stability results for Hilfer-fractional implicit differential equations with nonlocal conditions*, Mediterr. J. Math. **15**(1) (2018), 1–21.
- [25] J. R. Wang, Y. Zhou and M. Feckan, *On recent development in the theory of boundary value problems for impulsive fractional differential equations*, Comput. Math. Appl. **64**(10) (2012), 3008–3020.
- [26] Y. Zhou, *Basic Theory of Fractional Differential Equations*, World Scientific, Singapore, 2014.

<sup>1</sup>DEPARTMENT OF MATHEMATICS,  
 SRI VASAVI COLLEGE, ERODE,  
 TAMILNADU, INDIA-638 316  
 Email address: pkarthisvc@gmail.com  
 Email address: mathematicsarul@gmail.com