

## APPLICATIONS POISSON DISTRIBUTION AND RUSCHEWEYH DERIVATIVE OPERATOR FOR BI-UNIVALENT FUNCTIONS

ABBAS KAREEM WANAS<sup>1</sup> AND JANUSZ SOKÓŁ<sup>2</sup>

ABSTRACT. In this paper we establish upper bounds for the second and third coefficients of holomorphic and bi-univalent functions in a new family which involve the Bazilevič functions and  $\beta$ -pseudo-starlike functions under a new operator joining Poisson distribution with Ruscheweyh derivative operator. Also, we discuss Fekete-Szegő problem of functions in this family.

### 1. INTRODUCTION

Let  $\mathcal{A}$  be the collection of functions  $f$  that are holomorphic in the unit disk  $\mathbb{D} = \{|z| < 1\}$  in the complex plane  $\mathbb{C}$  and that have the form:

$$(1.1) \quad f(z) = z + \sum_{n=2}^{\infty} a_n z^n, \quad z \in \mathbb{D}.$$

Further, let  $\mathcal{S}$  be the sub-collection of  $\mathcal{A}$  containing of functions which are univalent in  $\mathbb{D}$ . According to the Koebe one-quarter theorem (see [3]), every function  $f \in \mathcal{S}$  has an inverse  $f^{-1}$  such that  $f^{-1}(f(z)) = z$ ,  $z \in \mathbb{D}$ , and  $f(f^{-1}(w)) = w$ ,  $|w| < r_0(f)$ ,  $r_0(f) \geq \frac{1}{4}$ . If  $f$  is of the form (1.1), then

$$(1.2) \quad f^{-1}(w) = w - a_2 w^2 + (2a_2^2 - a_3) w^3 - (5a_2^3 - 5a_2 a_3 + a_4) w^4 + \cdots, \quad |w| < r_0(f).$$

A function  $f \in \mathcal{A}$  is said to be bi-univalent in  $\mathbb{D}$  if both  $f$  and  $f^{-1}$  are univalent in  $\mathbb{D}$ . We denote by  $\Sigma$  the set of bi-univalent functions in  $\mathbb{D}$ . Srivastava et al. [19] have apparently resuscitated the study of holomorphic and bi-univalent functions in

---

*Key words and phrases.* Bi-univalent function,  $(M, N)$ -Lucas polynomial, coefficient bound, Fekete-Szegő problem, Poisson distribution, subordination, Ruscheweyh derivative.

2010 *Mathematics Subject Classification.* Primary: 30C45. Secondary: 30C80.

DOI 10.46793/KgJMat2401.089W

*Received:* May 03, 2020.

*Accepted:* March 05, 2021.

recent years. It was followed by such works as those by Frasin and Aouf [5], Goyal and Goswami [6], Srivastava and Bansal [15] and others (see, for example [2, 16–18, 20]).

For the polynomials  $M(x)$  and  $N(x)$  with real coefficients, the  $(M, N)$ -Lucas Polynomials  $L_{M,N,k}(x)$  are defined by the following recurrence relation (see [8]):

$$L_{M,N,k}(x) = M(x)L_{M,N,k-1}(x) + N(x)L_{M,N,k-2}(x), \quad k \geq 2,$$

with

$$(1.3) \quad L_{M,N,0}(x) = 2, \quad L_{M,N,1}(x) = M(x) \quad \text{and} \quad L_{M,N,2}(x) = M^2(x) + 2N(x).$$

The Lucas Polynomials play an important role in a diversity of disciplines in the mathematical, statistical, physical and engineering sciences (see, for example [4, 9, 21]). The generating function of the  $(M, N)$ -Lucas Polynomial  $L_{M,N,k}(x)$  (see [9]) is given by

$$(1.4) \quad T_{M(x),N(x)}(z) = \sum_{k=2}^{\infty} L_{M,N,k}(x)z^k = \frac{2 - M(x)z}{1 - M(x)z - N(x)z^2}.$$

Let the functions  $f$  and  $g$  be holomorphic in  $\mathbb{D}$ , we say that the function  $f$  is subordinate to  $g$ , if there exists a function  $w$ , holomorphic in  $\mathbb{D}$  with  $w(0) = 0$  and  $|w(z)| < 1$ ,  $z \in \mathbb{D}$ , such that  $f(z) = g(w(z))$ . This subordination is indicated by  $f \prec g$  or  $f(z) \prec g(z)$ ,  $z \in \mathbb{D}$ . Furthermore, if the function  $g$  is univalent in  $\mathbb{D}$ , then we have the following equivalence (see [10])

$$f(z) \prec g(z) \Leftrightarrow f(0) = g(0) \quad \text{and} \quad f(\mathbb{D}) \subset g(\mathbb{D}).$$

A function  $f \in \mathcal{A}$  is called Bazilevič function of order  $\alpha$ ,  $\alpha \geq 0$ , if (see [14])

$$\operatorname{Re} \left\{ \frac{z^{1-\alpha} f'(z)}{(f(z))^{1-\alpha}} \right\} > 0, \quad z \in \mathbb{D}.$$

A function  $f \in \mathcal{A}$  is called  $\beta$ -pseudo-starlike function of order  $\beta$ ,  $\beta \geq 1$ , if (see [1])

$$\operatorname{Re} \left\{ \frac{z (f'(z))^\beta}{f(z)} \right\} > 0, \quad z \in \mathbb{D}.$$

Recall that a random variable  $X$  has the Poisson distribution with parameter  $\theta$ , if

$$P(X = r) = \frac{\theta^r e^{-\theta}}{r!}, \quad r = 0, 1, 2, 3, \dots$$

Recently, Porwal [11] introduced a power series whose coefficients are probabilities of "Poisson distribution"

$$K(\theta, z) = z + \sum_{n=2}^{\infty} \frac{\theta^{n-1}}{(n-1)!} e^{-\theta} z^n, \quad z \in \mathbb{D},$$

where  $\theta > 0$ . By ratio test the radius of convergence of the above series is infinity.

In 2016, Porwal and Kumar [12] introduced and investigated a linear operator  $I(\theta, z) : \mathcal{A} \rightarrow \mathcal{A}$ ,  $\theta > 0$ , by using the Hadamard product (or convolution) and defined as follows

$$I(\theta, z)f(z) = K(\theta, z) * f(z) = z + \sum_{n=2}^{\infty} \frac{\theta^{n-1}}{(n-1)!} e^{-\theta} a_n z^n, \quad z \in \mathbb{D},$$

where "\*" indicate the Hadamard product (or convolution) of two power series.

In this paper, for  $f \in \mathcal{A}$  we introduce a new linear operator  $\mathcal{J}_\theta^\delta : \mathcal{A} \rightarrow \mathcal{A}$  by

$$(1.5) \quad \mathcal{J}_\theta^\delta f(z) = I(\theta, z) * \mathcal{R}^\delta,$$

where  $\mathcal{R}^\delta$ ,  $\delta \in \mathbb{N}_0 = \{0, 1, 2, \dots\}$ , denote the Ruscheweyh derivative operator [13] given by

$$\mathcal{R}^\delta f(z) = z + \sum_{n=2}^{\infty} \frac{\Gamma(\delta + n)}{\Gamma(\delta + 1)\Gamma(n)} a_n z^n, \quad z \in \mathbb{D}.$$

It is easy to obtain from (1.5) that

$$\mathcal{J}_\theta^\delta f(z) = z + \sum_{n=2}^{\infty} \frac{\theta^{n-1}\Gamma(\delta + n)}{\Gamma(\delta + 1)(\Gamma(n))^2} e^{-\theta} a_n z^n, \quad z \in \mathbb{D},$$

where  $\theta > 0$ ,  $\delta \in \mathbb{N}_0 = \{0, 1, 2, \dots\}$ .

## 2. MAIN RESULTS

We begin this section by defining the family  $\Upsilon_\Sigma(\lambda, \alpha, \beta, \delta, \theta; h)$  as follows.

**Definition 2.1.** Assume that  $\alpha \geq 0$ ,  $\beta \geq 1$ ,  $\delta \in \mathbb{N}_0$ ,  $0 \leq \lambda \leq 1$ ,  $\theta > 0$  and  $h$  is analytic in  $\mathbb{D}$ ,  $h(0) = 1$ . The function  $f \in \Sigma$  is in the family  $\Upsilon_\Sigma(\alpha, \beta, \delta, \lambda, \theta; h)$  if it fulfills the subordinations:

$$(1 - \lambda) \frac{z^{1-\alpha} \left(\mathcal{J}_\theta^\delta f(z)\right)'}{\left(\mathcal{J}_\theta^\delta f(z)\right)^{1-\alpha}} + \lambda \frac{z \left(\left(\mathcal{J}_\theta^\delta f(z)\right)'\right)^\beta}{\mathcal{J}_\theta^\delta f(z)} \prec h(z)$$

and

$$(1 - \lambda) \frac{w^{1-\alpha} \left(\mathcal{J}_\theta^\delta f^{-1}(w)\right)'}{\left(\mathcal{J}_\theta^\delta f^{-1}(w)\right)^{1-\alpha}} + \lambda \frac{w \left(\left(\mathcal{J}_\theta^\delta f^{-1}(w)\right)'\right)^\beta}{\mathcal{J}_\theta^\delta f^{-1}(w)} \prec 1 + e_1 z + e_2 z^2 + \dots,$$

where  $f^{-1}$  is given by (1.2).

In particular, if we choose  $\lambda = 1$  in Definition 2.1, the family  $\Upsilon_\Sigma(\alpha, \beta, \delta, \lambda, \theta; h)$  reduces to the family  $\mathcal{L}_\Sigma(\beta, \delta, \theta; h)$  of  $\beta$ -pseudo bi-starlike functions which satisfying the following subordinations:

$$\frac{z \left(\left(\mathcal{J}_\theta^\delta f(z)\right)'\right)^\beta}{\mathcal{J}_\theta^\delta f(z)} \prec h(z)$$

and

$$\frac{w \left( \left( \mathcal{J}_\theta^\delta f^{-1}(w) \right)' \right)^\beta}{\mathcal{J}_\theta^\delta f^{-1}(w)} \prec h(w).$$

If we choose  $\lambda = 0$  in Definition 2.1, the family  $\Upsilon_\Sigma(\alpha, \beta, \delta, \lambda, \theta; h)$  reduces to the family  $\mathcal{B}_\Sigma(\alpha, \delta, \theta; h)$  of Bazilevič bi-univalent functions which satisfying the following subordinations:

$$\frac{z^{1-\alpha} \left( \mathcal{J}_\theta^\delta f(z) \right)'}{\left( \mathcal{J}_\theta^\delta f(z) \right)^{1-\alpha}} \prec h(z)$$

and

$$\frac{w^{1-\alpha} \left( \mathcal{J}_\theta^\delta f^{-1}(w) \right)'}{\left( \mathcal{J}_\theta^\delta f^{-1}(w) \right)^{1-\alpha}} \prec h(w).$$

If we choose  $\lambda = \beta = 1$  in Definition 2.1, the family  $\Upsilon_\Sigma(\alpha, \beta, \delta, \lambda, \theta; h)$  reduces to the family  $\mathcal{S}_\Sigma(\delta, \theta; h)$  of bi-starlike functions which satisfying the following subordinations:

$$\frac{z \left( \mathcal{J}_\theta^\delta f(z) \right)'}{\mathcal{J}_\theta^\delta f(z)} \prec h(z)$$

and

$$\frac{w \left( \mathcal{J}_\theta^\delta f^{-1}(w) \right)'}{\mathcal{J}_\theta^\delta f^{-1}(w)} \prec h(w).$$

**Theorem 2.1.** *Assume that  $\alpha \geq 0$ ,  $\beta \geq 1$ ,  $\delta \in \mathbb{N}_0$ ,  $0 \leq \lambda \leq 1$  and  $\theta > 0$ . If  $f \in \Sigma$  of the form (1.1) is in the class  $\Upsilon_\Sigma(\alpha, \beta, \delta, \lambda, \theta; h)$ , with  $h(z) = 1 + e_1 z + e_2 z^2 + \dots$ , then*

$$(2.1) \quad |a_2| \leq \frac{|e_1|}{[(1-\lambda)(\alpha+1) + \lambda(2\beta-1)](\delta+1)\theta e^{-\theta}} = \frac{|e_1|}{A}$$

and

$$(2.2) \quad |a_3| \leq \min \left\{ \max \left\{ \left| \frac{e_1}{B} \right|, \left| \frac{e_2}{B} - \frac{C e_1^2}{A^2 B} \right| \right\}, \max \left\{ \left| \frac{e_1}{B} \right|, \left| \frac{e_2}{B} - \frac{(2B+C)e_1^2}{A^2 B} \right| \right\} \right\},$$

where

$$(2.3) \quad \begin{aligned} A &= [(1-\lambda)(\alpha+1) + \lambda(2\beta-1)](\delta+1)\theta e^{-\theta}, \\ B &= \frac{1}{4} [(1-\lambda)(\alpha+2) + \lambda(3\beta-1)](\delta^2 + 3\delta + 2)\theta^2 e^{-\theta}, \\ C &= \left[ \frac{1}{2}(1-\lambda)(\alpha+2)(\alpha-1) + \lambda(2\beta(\beta-2) + 1) \right] (\delta+1)^2 \theta^2 e^{-2\theta}. \end{aligned}$$

*Proof.* Suppose that  $f \in \Upsilon_\Sigma(\alpha, \beta, \delta, \lambda, \theta; ; e_1; e_2)$ . Then there exist two holomorphic functions  $\phi, \psi : \mathbb{D} \rightarrow \mathbb{D}$  given by

$$(2.4) \quad \phi(z) = r_1 z + r_2 z^2 + r_3 z^3 + \dots, \quad z \in \mathbb{D},$$

and

$$(2.5) \quad \psi(w) = s_1 w + s_2 w^2 + s_3 w^3 + \dots, \quad w \in \mathbb{D},$$

with  $\phi(0) = \psi(0) = 0$ ,  $|\phi(z)| < 1$ ,  $|\psi(w)| < 1$ ,  $z, w \in \mathbb{D}$  such that

$$(2.6) \quad (1 - \lambda) \frac{z^{1-\alpha} (\mathcal{J}_\theta^\delta f(z))'}{(\mathcal{J}_\theta^\delta f(z))^{1-\alpha}} + \lambda \frac{z \left( (\mathcal{J}_\theta^\delta f(z))' \right)^\beta}{\mathcal{J}_\theta^\delta f(z)} = 1 + e_1 \phi(z) + e_2 \phi^2(z) + \dots$$

and

$$(2.7) \quad (1 - \lambda) \frac{w^{1-\alpha} (\mathcal{J}_\theta^\delta f^{-1}(w))'}{(\mathcal{J}_\theta^\delta f^{-1}(w))^{1-\alpha}} + \lambda \frac{w \left( (\mathcal{J}_\theta^\delta f^{-1}(w))' \right)^\beta}{\mathcal{J}_\theta^\delta f^{-1}(w)} = 1 + e_1 \psi(w) + e_2 \psi^2(w) + \dots$$

Combining (2.4), (2.5), (2.6) and (2.7), yield

$$(2.8) \quad (1 - \lambda) \frac{z^{1-\alpha} (\mathcal{J}_\theta^\delta f(z))'}{(\mathcal{J}_\theta^\delta f(z))^{1-\alpha}} + \lambda \frac{z \left( (\mathcal{J}_\theta^\delta f(z))' \right)^\beta}{\mathcal{J}_\theta^\delta f(z)} = 1 + e_1 r_1 z + [e_1 r_2 + e_2 (x) r_1^2] z^2 + \dots$$

and

$$(2.9) \quad (1 - \lambda) \frac{w^{1-\alpha} (\mathcal{J}_\theta^\delta f^{-1}(w))'}{(\mathcal{J}_\theta^\delta f^{-1}(w))^{1-\alpha}} + \lambda \frac{w \left( (\mathcal{J}_\theta^\delta f^{-1}(w))' \right)^\beta}{\mathcal{J}_\theta^\delta f^{-1}(w)} = 1 + e_1 s_1 w + [e_1 s_2 + e_2 s_1^2] w^2 + \dots$$

It is quite well-known that if  $|\phi(z)| < 1$  and  $|\psi(w)| < 1$ ,  $z, w \in \mathbb{D}$ , we get

$$(2.10) \quad |r_j| \leq 1 \quad \text{and} \quad |s_j| \leq 1, \quad j \in \mathbb{N}.$$

In the light of (2.8) and (2.9), after simplifying, we find that

$$(2.11) \quad [(1 - \lambda)(\alpha + 1) + \lambda(2\beta - 1)] (\delta + 1) \theta e^{-\theta} a_2 = e_1 r_1,$$

$$\frac{1}{4} [(1 - \lambda)(\alpha + 2) + \lambda(3\beta - 1)] (\delta^2 + 3\delta + 2) \theta^2 e^{-\theta} a_3$$

$$+ \left[ \frac{1}{2} (1 - \lambda)(\alpha + 2)(\alpha - 1) + \lambda(2\beta(\beta - 2) + 1) \right] (\delta + 1)^2 \theta^2 e^{-2\theta} a_2^2$$

$$(2.12) \quad = e_1 r_2 + e_2 r_1^2,$$

$$(2.13) \quad - [(1 - \lambda)(\alpha + 1) + \lambda(2\beta - 1)] (\delta + 1) \theta e^{-\theta} a_2 = e_1 s_1$$

and

$$(2.14) \quad \begin{aligned} & \frac{1}{4} [(1-\lambda)(\alpha+2) + \lambda(3\beta-1)] (\delta^2 + 3\delta + 2) \theta^2 e^{-\theta} (2a_2^2 - a_3) \\ & + \left[ \frac{1}{2} (1-\lambda)(\alpha+2)(\alpha-1) + \lambda(2\beta(\beta-2) + 1) \right] (\delta+1)^2 \theta^2 e^{-2\theta} a_2^2 \\ & = e_1 s_2 + e_2 s_1^2. \end{aligned}$$

Inequality (2.1) follows from (2.11) and (2.13). If we apply notation (2.3), then (2.11) and (2.12) become

$$(2.15) \quad Aa_2 = e_1 r_1, \quad Ba_3 + Ca_2^2 = e_1 r_2 + e_2 r_1^2.$$

This gives

$$(2.16) \quad \frac{B}{e_1} a_3 = r_2 + \left( \frac{e_2}{e_1} - \frac{Ce_1}{A^2} \right) r_1^2,$$

and on using the known sharp result [7, page 10]:

$$(2.17) \quad |r_2 - \mu r_1^2| \leq \max \{1, |\mu|\},$$

for all  $\mu \in \mathbb{C}$ , we obtain

$$(2.18) \quad \left| \frac{B}{e_1} \right| |a_3| \leq \max \left\{ 1, \left| \frac{e_2}{e_1} - \frac{Ce_1}{A^2} \right| \right\}.$$

In the same way, (2.13) and (2.14) become

$$(2.19) \quad -Aa_2 = e_1 s_1, \quad B(2a_2^2 - a_3) + Ca_2^2 = e_1 s_2 + e_2 s_1^2.$$

This gives

$$(2.20) \quad -\frac{B}{e_1} a_3 = s_2 + \left( \frac{e_2}{e_1} - \frac{(2B+C)e_1}{A^2} \right) s_1^2.$$

Applying (2.17), we obtain

$$(2.21) \quad \left| \frac{B}{e_1} \right| |a_3| \leq \max \left\{ 1, \left| \frac{e_2}{e_1} - \frac{(2B+C)e_1}{A^2} \right| \right\}.$$

Inequality (2.2) follows from (2.18) and (2.21).  $\square$

If we take the generating function (1.4) of the  $(M, N)$ -Lucas polynomials  $L_{M,N,k}(x)$  as  $h(z) + 1$ , then from (1.3), we have  $e_1 = M(x)$  and  $e_2 = M^2(x) + 2N(x)$  and Theorem 2.1 becomes the following corollary.

**Corollary 2.1.** *If  $f \in \Sigma$  of the form (1.1) is in the class  $\Upsilon_{\Sigma}(\alpha, \beta, \delta, \lambda, \theta; T_{M(x), N(x)} - 1)$ , then*

$$|a_2| \leq \frac{|M(x)|}{[(1-\lambda)(\alpha+1) + \lambda(2\beta-1)] (\delta+1) \theta e^{-\theta}}$$

and

$$|a_3| \leq \min \left\{ \max \left\{ \left| \frac{M(x)}{B} \right|, \left| \frac{M^2(x) + 2N(x)}{B} - \frac{CM^2(x)}{A^2B} \right| \right\}, \max \left\{ \left| \frac{M(x)}{B} \right|, \left| \frac{M^2(x) + 2N(x)}{B} - \frac{(2B+C)M^2(x)}{A^2B} \right| \right\} \right\},$$

for all  $\alpha, \beta, \delta, \lambda, \theta, x$  such that  $\alpha \geq 0$ ,  $\beta \geq 1$ ,  $\delta \in \mathbb{N}_0$ ,  $0 \leq \lambda \leq 1$ ,  $\theta > 0$  and  $x \in \mathbb{R}$ , where  $A, B, C$  are given by (2.3) and  $T_{M(x), N(x)}$  is given by (1.4).

In the next theorem, we discuss a bound for  $|a_3 - \eta a_2^2|$  called "the Fekete-Szegő problem".

**Theorem 2.2.** *If  $f \in \Sigma$  of the form (1.1) is in the class  $\Upsilon_\Sigma(\alpha, \beta, \delta, \lambda, \theta; h)$ , then*

(2.22)

$$\begin{aligned} & |a_3 - \eta a_2^2| \\ & \leq \frac{|e_1|}{B} \min \left\{ \max \left\{ 1, \left| \frac{e_2}{e_1} - \frac{(C - \eta B)e_1}{A^2} \right| \right\}, \max \left\{ 1, \left| \frac{e_2}{e_1} - \frac{(2B + C - \eta B)e_1}{A^2} \right| \right\} \right\}, \end{aligned}$$

for all  $\alpha, \beta, \delta, \lambda, \theta, \eta$  such that  $\alpha \geq 0$ ,  $\beta \geq 1$ ,  $\delta \in \mathbb{N}_0$ ,  $0 \leq \lambda \leq 1$ ,  $\theta > 0$  and  $\eta \in \mathbb{C}$ , where  $A, B, C$  are given by (2.3).

*Proof.* We apply the notations from the proof of Theorem 2.1. From (2.15) and from (2.16), we have

$$(2.23) \quad a_3 - \eta a_2^2 = \frac{e_1}{B} \left( r_2 + \left( \frac{e_2}{e_1} - \frac{(C - \eta B)e_1}{A^2} \right) r_1^2 \right)$$

and on using the known sharp result  $|r_2 - \mu r_1^2| \leq \max \{1, |\mu|\}$ , we get

$$(2.24) \quad |a_3 - \eta a_2^2| \leq \frac{|e_1|}{B} \max \left\{ 1, \left| \frac{e_2}{e_1} - \frac{(C - \eta B)e_1}{A^2} \right| \right\}.$$

In the same way, from (2.19) and from (2.20), we have

$$(2.25) \quad a_3 - \eta a_2^2 = -\frac{e_1}{B} \left( s_2 + \left( \frac{e_2}{e_1} - \frac{(2B + C - \eta B)e_1}{A^2} \right) s_1^2 \right)$$

and on using  $|s_2 - \mu s_1^2| \leq \max \{1, |\mu|\}$ , we get

$$(2.26) \quad |a_3 - \eta a_2^2| \leq \frac{|e_1|}{B} \max \left\{ 1, \left| \frac{e_2}{e_1} - \frac{(2B + C - \eta B)e_1}{A^2} \right| \right\}.$$

Inequality (2.22) follows from (2.24) and (2.26). □

**Corollary 2.2.** *If  $f \in \Sigma$  of the form (1.1) is in the class  $\Upsilon_{\Sigma}(\alpha, \beta, \delta, \lambda, \theta; T_{M(x), N(x)} - 1)$ , then*

$$\left| a_3 - \eta a_2^2 \right| \leq \frac{|M(x)|}{B} \min \left\{ \max \left\{ 1, \left| \frac{M^2(x) + 2N(x)}{M(x)} - \frac{(C - \eta B)M(x)}{A^2} \right| \right\}, \right. \\ \left. \max \left\{ 1, \left| \frac{M^2(x) + 2N(x)}{M(x)} - \frac{(2B + C - \eta B)M(x)}{A^2} \right| \right\} \right\},$$

for all  $\alpha, \beta, \delta, \lambda, \theta, \eta, x$  such that  $\alpha \geq 0$ ,  $\beta \geq 1$ ,  $\delta \in \mathbb{N}_0$ ,  $0 \leq \lambda \leq 1$ ,  $\theta > 0$ ,  $\eta \in \mathbb{C}$  and  $x \in \mathbb{R}$ , where  $A, B, C$  are given by (2.3) and  $T_{M(x), N(x)}$  is given by (1.4).

#### REFERENCES

- [1] K. O. Babalola, *On  $\lambda$ -pseudo-starlike functions*, J. Class. Anal. **3**(2) (2013), 137–147. <https://doi.org/10.7153/jca-03-12>
- [2] M. Caglar, E. Deniz and H. M. Srivastava, *Second Hankel determinant for certain subclasses of bi-univalent functions*, Turkish J. Math. **41** (2017), 694–706. <https://doi.org/10.3906/mat-1602-25>
- [3] P. L. Duren, *Univalent Functions*, Springer Verlag, New York, Berlin, Heidelberg, Tokyo, 1983.
- [4] P. Filippini and A. F. Horadam, *Derivative sequences of Fibonacci and Lucas polynomials*, Applications of Fibonacci Numbers **4** (1991), 99–108. [https://doi.org/10.1007/978-94-011-3586-3\\_12](https://doi.org/10.1007/978-94-011-3586-3_12)
- [5] B. A. Frasin and M. K. Aouf, *New subclasses of bi-univalent functions*, Appl. Math. Lett. **24** (2011), 1569–1573. <https://doi.org/10.1016/j.aml.2011.03.048>
- [6] S. P. Goyal and P. Goswami, *Estimate for initial Maclaurin coefficients of bi-univalent functions for a class defined by fractional derivatives*, J. Egyptian Math. Soc. **20** (2012), 179–182. <https://doi.org/10.1016/j.joems.2012.08.020>
- [7] F. R. Keogh and E. P. Merkes, *A coefficient inequality for certain classes of analytic functions*, Proc. Amer. Math. Soc. **20** (1969), 8–12. <https://doi.org/10.2307/2035949>
- [8] G. Y. Lee and M. Asci, *Some properties of the  $(p, q)$ -Fibonacci and  $(p, q)$ -Lucas polynomials*, J. Appl. Math. **2012**, Article ID 264842. <https://doi.org/10.1155/2012/264842>
- [9] A. Lupas, *A guide of Fibonacci and Lucas polynomials*, Octagon Mathematical Magazine **7** (1999), 2–12.
- [10] S. S. Miller and P. T. Mocanu, *Differential Subordinations: Theory and Applications*, Series on Monographs and Textbooks in Pure and Applied Mathematics **225**, Marcel Dekker Inc., New York, Basel, 2000.
- [11] S. Porwal, *An application of a Poisson distribution series on certain analytic functions*, J. Complex Anal. **2014** (2014), Article ID 984135, 1–3. <https://doi.org/10.1155/2014/984135>
- [12] S. Porwal and M. Kumar, *A unified study on starlike and convex functions associated with Poisson distribution series*, Afrika Math. **27** (2016), 1021–1027. <https://doi.org/10.1007/s13370-016-0398-z>
- [13] S. Ruscheweyh, *New criteria for univalent functions*, Proc. Amer. Math. Soc. **49** (1975), 109–115. <https://doi.org/10.2307/2039801>
- [14] R. Singh, *On Bazilevič functions*, Proc. Amer. Math. Soc. **38**(2) (1973), 261–271. <https://doi.org/10.1090/S0002-9939-1973-0311887-9>
- [15] H. M. Srivastava and D. Bansal, *Coefficient estimates for a subclass of analytic and bi-univalent functions*, J. Egyptian Math. Soc. **23** (2015), 242–246. <https://doi.org/10.1016/j.joems.2014.04.002>
- [16] H. M. Srivastava, S. S. Eker and R. M. Ali, *Coefficient bounds for a certain class of analytic and bi-univalent functions*, Filomat **29** (2015), 1839–1845. <https://doi.org/10.2298/FIL1508839S>



- [17] H. M. Srivastava, S. S. Eker, S. G. Hamidi and J. M. Jahangiri, *Faber polynomial coefficient estimates for bi-univalent functions defined by the Tremblay fractional derivative operator*, Bull. Iranian Math. Soc. **44**(1) (2018), 149–157. <https://doi.org/10.1007/s41980-018-0011-3>
- [18] H. M. Srivastava, S. Gaboury and F. Ghanim, *Coefficient estimates for some general subclasses of analytic and bi-univalent functions*, Afrika Math. **28** (2017), 693–706. <https://doi.org/10.1007/s13370-016-0478-0>
- [19] H. M. Srivastava, A. K. Mishra and P. Gochhayat, *Certain subclasses of analytic and bi-univalent functions*, Appl. Math. Lett. **23** (2010), 1188–1192. <https://doi.org/10.1016/j.aml.2010.05.009>
- [20] A. K. Wanas and A. L. Alina, *Applications of Horadam polynomials on Bazilevič bi-univalent function satisfying subordinate conditions*, Journal of Physics: Conference Series **1294** (2019), 1–6. <https://doi.org/10.1088/1742-6596/1294/3/032003>
- [21] T. Wang and W. Zhang, *Some identities involving Fibonacci, Lucas polynomials and their applications*, Bull. Math. Soc. Sci. Math. Roumanie **55** (2012), 95–103. <https://doi.org/10.3390/math6120334>

<sup>1</sup>DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCE,  
UNIVERSITY OF AL-QADISIYAH,  
IRAQ  
*Email address:* abbas.kareem.w@qu.edu.iq

<sup>2</sup>COLLEGE OF NATURAL SCIENCES,  
UNIVERSITY OF RZESZÓW,  
UL. PROF. PIGONIA 1, 35-310 RZESZÓW, POLAND  
*Email address:* jsokol@ur.edu.pl