

ON SIMULTANEOUS APPROXIMATION AND COMBINATIONS OF LUPAS TYPE OPERATORS

T. A. K. SINHA¹, K. K. SINGH², AND AVINASH K. SHARMA³

ABSTRACT. The purpose of the present paper is to study a sequence of linear and positive operators which was introduced by A. Lupas. First, we obtain estimate of moments of the operators and then prove a basic convergence theorem for simultaneous approximation. Further, we find error in approximation in terms of modulus of continuity of function. Finally, we establish a Voronovskaja asymptotic formula for linear combination of the above operators.

1. INTRODUCTION

At the International Dortmund Meeting held in Written (Germany, March, 1995), A. Lupas [11] introduced the following Linear positive operators for $f : [0, \infty) \rightarrow \mathbb{R}$ as

$$(1.1) \quad L_n(f, x) = (1 - a)^{nx} \sum_{\nu=0}^{\infty} \frac{(nx)_{\nu}}{\nu!} a^{\nu} f\left(\frac{\nu}{n}\right), \quad x \geq 0,$$

$$(1.2) \quad (1 - a)^{-\alpha} = \sum_{\nu=0}^{\infty} \frac{(\alpha)_{\nu}}{\nu!} a^{\nu},$$

where

$$|a| < 1, \quad (\alpha)_0 = 1, \quad (\alpha)_{\nu} = \alpha(\alpha + 1) \cdots (\alpha + \nu - 1), \quad \nu \geq 1.$$

If we impose that $L_n(t, x) = x$, we find that $a = 1/2$. Therefore, operator (1.1) becomes

$$L_n(f, x) = 2^{-nx} \sum_{\nu=0}^{\infty} \frac{(nx)_{\nu}}{2^{\nu} \nu!} f\left(\frac{\nu}{n}\right), \quad x \geq 0.$$

Key words and phrases. Lupas operators, simultaneous approximation, modulus of continuity, Voronovskaja asymptotic formula, linear combinations,

2020 *Mathematics Subject Classification.* Primary: 41A25. Secondary: 41A28, 41A36.

DOI 10.46793/KgJMat2404.619S

Received: April 14, 2021.

Accepted: July 24, 2021.

It was seen that these operators are positive and linear and preserve linear functions. Bernstein polynomials [10] exhibit the property of simultaneous approximation. Simultaneous approximation for Baskakov operators, modified by Durrmeyer, was studied by Heilmann and Müller [8]. Another modification of Baskakov operators for simultaneous approximation was investigated by Sinha et al. [18]. Yet another modification of Baskakov operators viz., integral modification of Baskakov operators shows simultaneous approximation property in Thamer et al. [20]. This was studied for Durrmeyer modification of Bernstein polynomials by Gonska and Zhou [4]. In the summation-integral type operators Gupta et al. [7] explored the simultaneous approximation. So far research work was done for linear positive operators ([3],[6],[9],[12]-[15], [19]). Singh and Agrawal [17] proved simultaneous approximation by a linear combination of Bernstein-Durrmeyer type polynomials. Gupta [5] studied the differences of operators of Lupas type. So, the Lupas operators play very important role to approximate functions for $f \in C[0, \infty)$.

It turns out that the order of approximation by these operators is at best $O(n^{-1})$, however smooth the function may be. Therefore, in order to improve the order of approximation by the operators (1.1), we apply the technique of linear combination introduced by Butzer [2] and Rathore [16].

The approximation process for linear combination is defined as follows.

Let d_0, d_1, \dots, d_k be $(k+1)$ arbitrary but fixed distinct positive integers. Then, the linear combination $L_n(f, k, x)$ of $L_{d_j n}(f, x), j = 0, 1, 2, \dots, k$, is given by

$$L_n(f, k, x) = \frac{1}{\Delta} \begin{vmatrix} L_{d_0 n}(f, x) & d_0^{-1} & d_0^{-2} & \cdots & d_0^{-k} \\ L_{d_1 n}(f, x) & d_1^{-1} & d_1^{-2} & \cdots & d_1^{-k} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ L_{d_k n}(f, x) & d_k^{-1} & d_k^{-2} & \cdots & d_k^{-k} \end{vmatrix},$$

where Δ is the Vandermonde determinant defined as

$$\Delta = \begin{vmatrix} 1 & d_0^{-1} & d_0^{-2} & \cdots & d_0^{-k} \\ 1 & d_1^{-1} & d_1^{-2} & \cdots & d_1^{-k} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & d_k^{-1} & d_k^{-2} & \cdots & d_k^{-k} \end{vmatrix}.$$

On simplification, we have

$$(1.3) \quad L_n(f, k, x) = \sum_{j=0}^k C(j, k) L_{d_j n}(f, x),$$

where

$$C(j, k) = \prod_{i=0, i \neq j}^k \frac{d_j}{d_j - d_i}, \quad k \neq 0 \text{ and } C(0, 0) = 1.$$

2. MOMENT ESTIMATES

Lemma 2.1 ([5]). *The following relations hold:*

$$L_n(1, x) = 1, \quad L_n(t - x, x) = \frac{2a - 1}{1 - a}x, \quad L_n((t - x)^2, x) = \frac{n^2x^2(2a - 1)^2 + nax}{n^2(1 - a)^2}.$$

Now, we define m th order moment

$$\mu_m(x) = L_n((t - x)^m, x) = (1 - a)^{nx} \left\{ \sum_{\nu=0}^{\infty} \frac{(nx)_{\nu}}{\nu!} a^{\nu} \left(\frac{\nu}{n} - x \right)^m \right\}.$$

Lemma 2.2. $\mu_m(x)$ is a polynomial in x of degree $[m/2]$. Moreover

$$\mu_m(x) = O\left(\frac{1}{n^{\lfloor \frac{m+1}{2} \rfloor}}\right), \quad n \rightarrow \infty.$$

Proof. By definition of moments of m th order, we have

$$\begin{aligned} \mu_m(x) &= (1 - a)^{nx} \sum_{\nu=0}^{\infty} (-1)^m a^{\nu} \frac{(nx)_{\nu}}{\nu!} \left\{ \sum_{r=0}^m \binom{m}{r} (-1)^r x^{m-r} \left(\frac{\nu}{n}\right)^r \right\} \\ (2.1) \quad &= (-1)^m (1 - a)^{nx} \sum_{\nu=0}^{\infty} \frac{(nx)_{\nu}}{\nu!} a^{\nu} \left\{ \sum_{r=0}^m \binom{m}{r} \left(-\frac{1}{n}\right)^r x^{m-r} \right. \\ &\quad \left. \times (\nu^{(r)} + p_2\nu^{(r-1)} + p_4\nu^{(r-2)} + \dots) \right\}, \end{aligned}$$

where $\nu^{(r)} = \nu(\nu - 1)(\nu - 2) \dots (\nu - r + 1)$, p_2 is a polynomial in r of second degree, p_4 is a polynomial in r of fourth degree and so on.

It follows from (1.2) upon s times differentiation in a that

$$(2.2) \quad \sum_{\nu=0}^{\infty} \frac{(nx)_{\nu}}{\nu!} \nu(\nu - 1) \dots (\nu - s + 1) a^{\nu-s} = (nx)_s (1 - a)^{-nx-s}.$$

Making an use of (2.2) in (2.1)

$$\begin{aligned} \mu_m(x) &= (-1)^m \sum_{r=0}^m \binom{m}{r} \left(-\frac{1}{n}\right)^r x^{m-r} \left\{ \frac{a^r}{(1 - a)^r} (nx)_r + p_2 \frac{a^{r-1}}{(1 - a)^{r-1}} (nx)_{r-1} \right. \\ &\quad \left. + p_4 \frac{a^{r-2}}{(1 - a)^{r-2}} (nx)_{r-2} + \dots \right\}. \end{aligned}$$

Again,

$$\frac{(nx)_r}{n^r x^r} = 1 + \frac{q_2}{nx} + \frac{q_4}{(nx)^2} + \frac{q_6}{(nx)^3} + \dots,$$

where q_j as before is a polynomial in r of degree j .

Therefore, taking $a = 1/2$ and using fact that $\sum_{r=0}^m \binom{m}{r} (-1)^r r^s = 0$, $s < m$, we find that

$$\mu_m(x) = (-1)^m x^m \left\{ \frac{C}{(nx)^{\lfloor \frac{m+1}{2} \rfloor}} + \cdots \text{higher order terms} \right\}.$$

Therefore, $\mu_m(x)$ is a polynomial in x of degree $\lfloor m/2 \rfloor$. This completes the proof of lemma. \square

3. SIMULTANEOUS APPROXIMATION

Theorem 3.1. *Let $f' \in C_B[0, \infty)$. Then, sequence $\left\{ \frac{d}{dx}(L_n(f, x)) \right\}_{n=1}^{\infty}$ converges to $f'(x)$ pointwise on $[0, \infty)$. Moreover, if S is a compact subset of $[0, \infty)$ then sequence $\left\{ \frac{d}{dx}(L_n(f, x)) \right\}_{n=1}^{\infty}$ converges to $f'(x)$ uniformly on S .*

Proof. We expand

$$f(w) = f(x) + (w - x)f'(x) + \int_x^w (f'(t) - f'(x))dt.$$

Operating $L_n(\cdot, y)$ on both sides of above equation and in view of Lemma 2.1, we obtain

$$L_n(f, y) = f(x) + \left(\frac{ay}{1-a} - x \right) f'(x) + (1-a)^{ny} \left\{ \sum_{\nu=0}^{\infty} \frac{(ny)_{\nu}}{\nu!} a^{\nu} R_{\nu} \right\},$$

where $R_{\nu} = \int_x^{\nu/n} (f'(t) - f'(x))dt$. Thus,

$$(3.1) \quad \frac{d}{dx} L_n(f, x) = \frac{a}{1-a} f'(x) + n(1-a)^{nx} \\ \times \left\{ \ln(1-a) \sum_{\nu=0}^{\infty} \frac{(nx)_{\nu}}{\nu!} a^{\nu} R_{\nu} + \sum_{\nu=1}^{\infty} \frac{d(nx)_{\nu}}{d(nx)} \frac{a^{\nu}}{\nu!} R_{\nu} \right\}.$$

We put $nx = \alpha$ and differentiate (1.2) w.r.t. α . Further, we equate coefficient of a^{ν} on both sides, we get

$$(3.2) \quad \frac{1}{\nu!} \cdot \frac{d(\alpha)_{\nu}}{d\alpha} = \frac{\alpha_{\nu-1}}{(\nu-1)!} + \frac{1}{2} \cdot \frac{\alpha_{\nu-2}}{(\nu-2)!} + \frac{1}{3} \cdot \frac{\alpha_{\nu-3}}{(\nu-3)!} + \cdots + \frac{1}{\nu} \cdot \frac{\alpha_0}{0!}.$$

Using (3.2) in (3.1), we get

$$\begin{aligned}
 & \frac{d}{dx}L_n(f, x) - \frac{a}{1-a}f'(x) \\
 &= n(1-a)^\alpha \left[a(R_1 - R_0) + a^2 \left\{ \frac{(\alpha)_1}{1!}(R_2 - R_1) + \frac{(\alpha)_0}{2}(R_2 - R_0) \right\} \right. \\
 & \quad + a^3 \left\{ \frac{(\alpha)_2}{2!}(R_3 - R_2) + \frac{1}{2} \cdot \frac{(\alpha)_1}{1!}(R_3 - R_1) + \frac{1}{3}(\alpha)_0(R_3 - R_0) \right\} + \dots \\
 & \quad + a^\nu \left\{ \frac{(\alpha)_{\nu-1}}{(\nu-1)!}(R_\nu - R_{\nu-1}) + \frac{(\alpha)_{\nu-2}}{(\nu-2)!} \cdot \frac{1}{2}(R_\nu - R_{\nu-2}) \right. \\
 & \quad + \frac{(\alpha)_{\nu-3}}{(\nu-3)!} \cdot \frac{1}{3}(R_\nu - R_{\nu-3}) + \dots + \frac{(\alpha)_1}{1!} \cdot \frac{1}{\nu-1}(R_\nu - R_1) \\
 & \quad \left. + \frac{1}{\nu} \cdot \frac{(\alpha)_0}{1}(R_\nu - R_0) \right\} + \dots \Big] \\
 &= n(1-a)^\alpha \left[a \left\{ (\alpha)_0(R_1 - R_0) + \frac{a(\alpha)_1}{1!}(R_2 - R_1) + \frac{a^2(\alpha)_2}{2!}(R_3 - R_2) \right. \right. \\
 & \quad + \left. \left. \frac{a^3(\alpha)_3}{3!}(R_4 - R_3) + \dots \right\} + a^2 \left\{ \frac{(\alpha)_0}{2}(R_2 - R_0) + \frac{a(\alpha)_1}{1!} \cdot \frac{1}{2}(R_3 - R_1) \right. \right. \\
 & \quad + \left. \left. \frac{a^2(\alpha)_2}{2!} \cdot \frac{1}{2}(R_4 - R_2) + \frac{a^3(\alpha)_3}{3!} \cdot \frac{1}{2}(R_5 - R_3) + \dots \right\} \right. \\
 & \quad + a^3 \left\{ (\alpha)_0 \frac{1}{3}(R_3 - R_0) + \frac{a(\alpha)_1}{1!} \cdot \frac{1}{3}(R_4 - R_1) + \frac{a^2(\alpha)_2}{2!} \cdot \frac{1}{3}(R_5 - R_2) + \dots \right\} \\
 & \quad \left. + \dots \right] \\
 (3.3) \quad &= n(1-a)^\alpha [\Sigma_1 + \Sigma_2 + \Sigma_3 + \dots], \quad \text{say.}
 \end{aligned}$$

The continuity of $f'(\cdot)$ at point x implies that for a given $\epsilon > 0$ there exists a $\delta = \delta(x)$, (depending on x) such that $|f'(t) - f'(x)| < \epsilon$ if $|t - x| < \delta$. We break $R_p - R_q$ in two parts depending upon $|t - x| < \delta$ and $|t - x| \geq \delta$. In the second part, there may be two terms, where $|f'(t) - f'(x)| \leq 2\|f'\|_{C_B[0,\infty)} \cdot \frac{1}{\delta^2}(t - x)^2$.

Using Lemma 2.1, we get

$$\begin{aligned}
 |\Sigma_1| &\leq a \frac{\epsilon}{n} \left(\sum_{k=0}^{\infty} \frac{a^k}{k!} (\alpha)_k \right) + \frac{2 \cdot 2 \|f'\|_{C_B[0,\infty)}}{\delta^2} \cdot \frac{a}{n} \left\{ \sum_{k=0}^{\infty} \frac{a^k}{k!} (\alpha)_k \left(\frac{k}{n} - x \right)^2 \right\} \\
 (3.4) \quad &= a \frac{\epsilon}{n} (1-a)^{-\alpha} + \frac{4 \|f'\|_{C_B[0,\infty)}}{\delta^2} \cdot \frac{a}{n} \cdot \left\{ \frac{nx^2(2a-1)^2 + ax}{n(1-a)^2} \right\} (1-a)^{-\alpha}.
 \end{aligned}$$

Now,

$$(3.5) \quad \begin{aligned} |\Sigma_2| &\leq a^2 \frac{\epsilon}{n} \left\{ \sum_{k=0}^{\infty} \frac{a^k}{k!} (\alpha)_k \right\} + \frac{4 \|f'\|_{C_B[0,\infty)}}{\delta^2} \cdot \frac{a^2}{n} \left\{ \sum_{k=0}^{\infty} \frac{a^k}{k!} (\alpha)_k \left(\frac{k}{n} - x \right)^2 \right\} \\ &= a^2 \frac{\epsilon}{n} (1-a)^{-\alpha} + \frac{4 \|f'\|_{C_B[0,\infty)}}{\delta^2} \cdot \frac{a^2}{n} \left\{ \frac{nx^2(2a-1)^2 + ax}{n(1-a)^2} \right\} (1-a)^{-\alpha}. \end{aligned}$$

The similar estimates for $\Sigma_3, \Sigma_4, \dots$ are combined in (3.3) and we take $a = \frac{1}{2}$ due to Agratini [1]. Finally,

$$\left| \frac{d}{dx} L_n(f, x) - f'(x) \right| \leq Cn \left(\frac{\epsilon}{n} + \frac{1}{n^2} \right).$$

This completes the proof of the first part.

Proof of second part of Theorem 3.1. Let S be a compact subset of $[0, \infty)$. The pointwise continuity of function $f'(\cdot)$ at points of S , imply, by virtue of compactness of S , that $f'(\cdot)$ is now uniformly continuous on S . Thus, δ is now independent of x . Moreover S , being compact, is a bounded subset of $[0, \infty)$. Thus $x \in S$ implies $|x| < C_1$, a constant. This implies by (3.4) and (3.5) that convergence is uniform. \square

Theorem 3.2. Let $f' \in C_B[0, \infty)$. Then for $\delta > 0$ and $[a, b] \subset (a_1, b_1)$ we have

$$\sup_{x \in [a, b]} |L'_n(f, x) - f'(x)| \leq \omega(f', \delta, [a_1, b_1]) + \frac{C}{n} \|f'\|_{C_B[0, \infty)}.$$

Proof. We proceed in similar way as in the proof of Theorem 3.1. In the steps following (3.3) if $|t - x| < \delta$, then $|f'(t) - f'(x)| \leq \omega(f', \delta, [a_1, b_1])$. When $|t - x| \geq \delta$, using boundedness of f' the total contribution is of order $\|f'\|_{C_B[0, \infty)} O\left(\frac{1}{n}\right)$ as $n \rightarrow \infty$, by Lemma 2.1. Hence, the proof follows. \square

4. LINEAR COMBINATIONS

Theorem 4.1. Let $f^{(2k+2)} \in C_B[0, \infty)$. Then there holds for each $x \in [0, \infty)$, point-wise:

$$(4.1) \quad \frac{d}{dx} L_n(f, k, x) - f'(x) = \frac{1}{n^{k+1}} \left\{ \sum_{j=k+2}^{2k+2} q_j(x) f^{(j)}(x) \right\} + o\left(\frac{1}{n^{k+1}}\right), \quad n \rightarrow \infty.$$

Moreover, if S is a compact subset of $[0, \infty)$, then convergence (4.1) is uniform on S .

Proof. Using Taylor’s series expansion, we write

$$\begin{aligned} f(w) = & f(x) + (w - x)f'(x) + \frac{(w - x)^2}{2!}f^{(2)}(x) + \dots \\ & + \frac{(w - x)^{2k+2}}{(2k + 2)!}f^{(2k+2)}(x) \\ & + \int_x^w \int_x^{t_1} \int_x^{t_2} \dots \int_x^{t_{2k+1}} (f^{(2k+2)}(u) - f^{(2k+2)}(x)) dt_{2k+1} dt_{2k} \dots dt_1 du. \end{aligned}$$

Operating $L_n(\cdot, y)$ on both sides of above equation and in view of Lemma 2.1, we obtain

$$\begin{aligned} L_n(f, y) = & f(x) + \left(\frac{ay}{1 - a} - x\right) f'(x) + \frac{f^{(2)}(x)}{2!}p_2(1/n, y) \\ & + \frac{f^{(3)}(x)}{3!}p_3(1/n, y) + \dots + \frac{f^{(2k+2)}(x)}{(2k + 2)}p_{2k+2}(1/n, y) \\ & + (1 - a)^{ny} \left(\sum_{\nu=0}^{\infty} \frac{(ny)_{\nu}}{\nu!} a^{\nu} R_{\nu} \right), \end{aligned}$$

where

$$R_{\nu} = \int_x^{\nu/n} \int_x^{t_1} \int_x^{t_2} \dots \int_x^{t_{2k+1}} (f^{(2k+2)}(u) - f^{(2k+2)}(x)) dt_{2k+1} dt_{2k} \dots dt_1 du$$

and $p_j\left(\frac{1}{n}, y\right)$ is a polynomial in y of degree j and in $\frac{1}{n}$ of degree $(j - 1)$.

This implies that

$$\begin{aligned} (4.2) \quad \frac{d}{dx}L_n(f, x) = & \frac{a}{1 - a}f'(x) + \frac{f^{(2)}(x)}{2!}p'_2(1/n, x) \\ & + \frac{f^{(3)}(x)}{3!}p'_3(1/n, x) + \dots + \frac{f^{(2k+2)}(x)}{(2k + 2)}p'_{2k+2}(1/n, x) \\ & + n(1 - a)^{nx} \left\{ \log(1 - a) \sum_{\nu=0}^{\infty} \frac{(nx)_{\nu}}{\nu!} a^{\nu} R_{\nu} + \sum_{\nu=1}^{\infty} \frac{d(nx)_{\nu}}{d(nx)} \cdot \frac{a^{\nu}}{\nu!} R_{\nu} \right\}. \end{aligned}$$

Let $\phi(n, x) = n(1 - a)^{nx} \left\{ \log(1 - a) \sum_{\nu=0}^{\infty} \frac{(nx)_{\nu}}{\nu!} a^{\nu} R_{\nu} + \sum_{\nu=1}^{\infty} \frac{d(nx)_{\nu}}{d(nx)} \cdot \frac{a^{\nu}}{\nu!} R_{\nu} \right\}$. Now, taking linear combinations on (4.2) and using their properties (1.3), we have

$$\frac{d}{dx}L_n(f, k, x) - \left(\frac{a}{1 - a}\right) f'(x) = \left\{ \sum_{j=k+2}^{2k+2} q_j(x)f^{(j)}(x) \right\} \frac{1}{n^{k+1}} + \sum_{j=0}^k C(j, k)\phi(d_j n, x).$$

We analyze last term as in (3.1) and obtain the required result.

The proof of the second part of theorem follows from the proof of the second part of Theorem 3.1. □

Acknowledgements. The authors are thankful to the reviewers for helpful remarks and suggestions which lead to essential improvement of the whole manuscript.

REFERENCES

- [1] O. Agratini, *On a sequence of linear and positive operators*, Facta Univ. Ser. Math. Inform. **14** (1999), 41–48.
- [2] P. L. Butzer, *Linear combinations of Bernstein polynomials*, Canadian J. Math. **5** (1953), 559–567.
- [3] F. Özger, H. M. Srivastava and S. A. Mohiuddine, *Approximation of functions by a new class of generalized Bernstein-Schurer operators*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Math. RACSAM **114** (2020), Article ID 173. <https://doi.org/10.1007/s13398-020-00903-6>
- [4] H. H. Gonska and X. L. Zhou, *A global Inverse theorem on simultaneous approximation by Bernstein Durrmeyer operators*, J. Approx. Theory **67** (1991), 284–302. [https://doi.org/10.1016/0021-9045\(91\)90004-T](https://doi.org/10.1016/0021-9045(91)90004-T)
- [5] V. Gupta, *Differences of operators of Lupas type*, Constructive Mathematical Analysis **1**(1) (2018), 9–14. <https://doi.org/10.33205/cma.452962>
- [6] V. Gupta and H. M. Srivastava, *A general family of the Srivastava-Gupta operators preserving linear functions*, Eur. J. Pure Appl. Math. **11**(3) (2018), 575–579. <https://doi.org/10.29020/nybg.ejpam.v11i3.3314>
- [7] V. Gupta, M. K. Gupta and V. Vasishtha, *Simultaneous approximation by summation integral type operators*, Journal of Nonlinear Functional Analysis **8**(3) (2003), 399–412.
- [8] M. Heilmann and M. W. Müller, *On simultaneous approximation by the method of Baskakov-Durrmeyer operators*, Numer. Funct. Anal. Optim. **10**(1–2) (1989), 127–138. <https://doi.org/10.1080/01630568908816295>
- [9] A. Kajla and T. Acar, *Modified α -Bernstein operators with better approximation properties*, Ann. Funct. Anal. **10**(4) (2019), 570–582. <https://doi.org/10.1215/20088752-2019-0015>
- [10] G. G. Lorentz, *Bernstein Polynomials*, Chelsea Publishing Company, New York, 1986.
- [11] A. Lupas, *The approximation by some positive linear operators*, in: M. W. Müller et al. (Eds.), *Proceedings of the International Dortmund Meeting on Approximation Theory*, Akademie Verlag, Berlin, 1995, 201–229.
- [12] A. Kajla, S. A. Mohiuddine and A. Alotaibi, *Blending-type approximation by Lupas-Durrmeyer-type operators involving Pólya distribution*, Math. Methods Appl. Sci. **44** (2021), 9407–9418. <https://doi.org/10.1002/mma.7368>
- [13] S. A. Mohiuddine and F. Özger, *Approximation of functions by Stancu variant of Bernstein-Kantorovich operators based on shape parameter α* , Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Math. RACSAM **114** (2020), Article ID 70. <https://doi.org/10.1007/s13398-020-00802-w>
- [14] S. A. Mohiuddine, T. Acar and A. Alotaibi, *Construction of a new family of Bernstein-Kantorovich operators*, Math. Methods Appl. Sci. **40** (2017), 7749–7759. <https://doi.org/10.1002/mma.4559>
- [15] S. A. Mohiuddine, N. Ahmad, F. Özger, A. Alotaibi and B. Hazarika, *Approximation by the parametric generalization of Baskakov-Kantorovich operators linking with Stancu operators*, Iran. J. Sci. Technol. Trans. A Sci. **45** (2021), 593–605. <https://doi.org/10.1007/s40995-020-01024-w>
- [16] R. K. S. Rathore, *Linear combination of linear positive operators and generating relations in special functions*, Ph. D. Thesis, I. I. T. Delhi, India, 1973.
- [17] K. K. Singh and P. N. Agrawal, *Simultaneous approximation by a linear combination of Bernstein-Durrmeyer type polynomials*, Bull. Math. Anal. Appl. **3**(2) (2011), 70–82.

- [18] R. P. Sinha, P. N. Agrawal and V. Gupta, *On simultaneous approximation by modified Baskakov operators*, Bull. Soc. Math. Belg. Ser. B. **43**(2) (1991), 217–231.
- [19] H. M. Srivastava and V. Gupta, *A certain family of summation-integral type operators*, Math. Comput. Modelling **37** (2003), 1307–1315. [https://doi.org/10.1016/S0895-7177\(03\)90042-2](https://doi.org/10.1016/S0895-7177(03)90042-2)
- [20] K. J. Thamer and A. I. Ibrahim, *Simultaneous approximation with linear combination of integral Baskakov type operators*, Revista De La Union Matematica Argentina **46**(1) (2005), 1–10.

¹DEPARTMENT OF MATHEMATICS,
 PATLIPUTRA UNIVERSITY,
 BODH-GAYA-824234(BIHAR), INDIA
Email address: thakurashok1212@gmail.com

²DEPARTMENT OF APPLIED SCIENCES AND HUMANITIES,
 INSTITUTE OF ENGINEERING AND TECHNOLOGY,
 DR. A.P.J. ABDUL KALAM TECHNICAL UNIVERSITY,
 LUCKNOW-226021(UTTAR PRADESH), INDIA
Email address: kksiitr.singh@gmail.com

³PG DEPARTMENT OF MATHEMATICS,
 MAGADH UNIVERSITY,
 BODH-GAYA-824234(BIHAR), INDIA
Email address: ak95257016@gmail.com