

EXISTENCE OF CLASSICAL SOLUTIONS FOR BROER-KAUP EQUATIONS

DALILA BOURENI¹, SVETLIN G. GEORGIEV², AREZKI KHELOUFI³,
 AND KARIMA MEBARKI⁴

ABSTRACT. In this paper we investigate the Cauchy problem for one dimensional Broer-Kaup equations for existence of global classical solutions. We give conditions under which the considered equations have at least one and at least two classical solutions. To prove our main results we propose a new approach based upon recent theoretical results.

1. INTRODUCTION

Study of existence of global classical solutions of nonlinear models is one of the important works in nonlinear science. In this paper, we investigate the Cauchy problem for a model describing the bi-directional propagation of long waves in shallow water which was proposed by Broer and Kaup [2, 9] and called Broer-Kaup (BK) equations. Namely, we are concerned with the following system:

$$\begin{aligned}
 (1.1) \quad & u_t + uu_x + v_x = 0, \quad t \in (0, \infty), x \in \mathbb{R}, \\
 & v_t + u_x + 2(uv)_x + u_{xxx} = 0, \quad t \in (0, \infty), x \in \mathbb{R}, \\
 & u(0, x) = u_0(x), \quad x \in \mathbb{R}, \\
 & v(0, x) = v_0(x), \quad x \in \mathbb{R},
 \end{aligned}$$

where

(H1): $u_0, v_0 \in \mathcal{C}^1(\mathbb{R})$, $0 \leq u_0, v_0 \leq B$ on \mathbb{R} for some positive constant B .

Here the unknowns $u = u(t, x)$ and $v = v(t, x)$ denote respectively, the horizontal velocity and the elevation of the water wave. The Broer-Kaup equations of system

Key words and phrases. Broer-Kaup equations, classical solution, fixed point, initial value problem.
 2020 *Mathematics Subject Classification.* Primary: 35Q35. Secondary: 35A09, 35E15.

DOI 10.46793/KgJMat2501.125B

Received: September 09, 2021.

Accepted: January 09, 2022.

(1.1) can be obtained from the symmetry constraints of the Kadomtsev-Petviashvili (KP) equation and are a mathematical model of many nonlinear waves, see [12]. More precisely, they describe the evolution of the horizontal velocity component $u(t, x)$ of water waves of height $v(t, x)$ propagating in both directions in an infinite narrow channel of finite constant depth. Several methods have been used to capture different nature of solutions contained in Broer-Kaup equations like traveling wave solutions, periodic wave solutions, dromion solutions, solitary wave solutions and soliton-like solutions, see [21] and [29]. By qualitative analysis method, a sufficient condition for the existence of peaked periodic wave solutions to the Broer-Kaup equations was given in [8] and some exact explicit expressions of peaked periodic wave solutions were also presented. In [16], fission and fusion phenomena were revealed and soliton solutions were obtained. A family of traveling wave solutions is given in [18, 19] and [7]. Solitary wave solutions to the Broer-Kaup equations are considered in [14] by using the first integral method. By application of the sub-ode method [25], new and more general form solutions are obtained for the Broer-Kaup equations. Using a consistent tanh expansion method, Chen et al. [1] gave the interaction solutions between the solitons and other different types of nonlinear waves. In [13], some smooth and peaked solitary wave solutions have been constructed by the bifurcation method of dynamical system. By using a Darboux transformation, Zhou et al. [27] obtained new exact solutions for Broer-Kaup system. In [6], new type of solitary wave solutions for the Broer-Kaup equations were presented by using the He's variational principle.

Various algebraic aspects of BK equations solutions have been studied. Kupershmidt [11] showed that BK equations are integrable and possess infinite number of conservation laws and tri-Hamiltonian structure. In [4], The geometric properties of non-Noether symmetries as well as their applications were discussed.

The analysis by many methods of the (2+1)-dimensional BK system can be found in [26] and [28] and the references therein. The (1+1)-dimensional and the (2 + 1)-dimensional higher order Broer-Kaup equation was considered for example in [23] and [20], respectively. Concerning generalized and variable coefficient Broer-Kaup equations, see for example [22] and [10]. Recently, fractional and stochastic Broer-Kaup system, were studied in [3] and [24].

The aim of this paper is to investigate the initial value problem (1.1) for existence and nonuniqueness of global classical solutions. For goal, a new topological approach which uses the abstract theory of the sum of two operators is used for investigations of existence of at least one and at least two classical solutions. This basic and new idea can be used for investigations for existence of global classical solutions for many of the interesting equations of mathematical physics. Here, by a classical solution to the Broer-Kaup equations we mean a solution at least three times continuously differentiable in x and once in t for any $t \geq 0$. In other words, (u, v) belongs to the space $\mathcal{C}^1([0, \infty), \mathcal{C}^3(\mathbb{R})) \times \mathcal{C}^1([0, \infty), \mathcal{C}^3(\mathbb{R}))$ of continuously differentiable functions on $[0, \infty)$ with values in the Banach space $\mathcal{C}^3(\mathbb{R})$.

The paper is organized as follows. In the next section, we give some properties of solutions of problem (1.1). First, we give an integral representation of these solutions, then we prove some *a priori* estimates in a sense that will be defined later on. In Section 3 we prove our main results about existence and multiplicity of solutions for the Broer-Kaup system (1.1). Finally, in Section 4 we give an example to illustrate our main results.

2. SOME PROPERTIES OF SOLUTIONS OF PROBLEM (1.1)

Let $X = X^1 \times X^1$, where $X^1 = \mathcal{C}^1([0, \infty), \mathcal{C}^3(\mathbb{R}))$. For $(u, v) \in X$, define the operators S_1^1 , S_1^2 and S_1 as follows.

$$\begin{aligned} S_1^1(u, v)(t, x) &= u(t, x) - u_0(x) + \int_0^t (u(t_1, x)u_x(t_1, x) + v_x(t_1, x))dt_1, \\ S_1^2(u, v)(t, x) &= v(t, x) - v(0, x) + \int_0^t \left(u_x(t_1, x) + 2u_x(t_1, x)v(t_1, x) \right. \\ &\quad \left. + 2u(t_1, x)v_x(t_1, x) + u_{xxx}(t_1, x) \right) dt_1, \\ S_1(u, v)(t, x) &= \left(S_1^1(u, v)(t, x), S_1^2(u, v)(t, x) \right), \quad (t, x) \in [0, \infty) \times \mathbb{R}. \end{aligned}$$

2.1. Integral representation of the solutions.

Lemma 2.1. *Suppose that (H1) is satisfied. If $(u, v) \in X$ satisfies the equation*

$$(2.1) \quad S_1(u, v)(t, x) = 0, \quad (t, x) \in [0, \infty) \times \mathbb{R},$$

then (u, v) is a solution of the IVP (1.1).

Proof. Let $(u, v) \in X$ be a solution of the equation (2.1). Then

$$(2.2) \quad S_1^1(u, v)(t, x) = 0, \quad S_1^2(u, v)(t, x) = 0, \quad (t, x) \in [0, \infty) \times \mathbb{R}.$$

We differentiate both equations of (2.2) with respect to t and x and we find

$$u_t(t, x) + u(t, x)u_x(t, x) + v_x(t, x) = 0,$$

$$v_t(t, x) + u_x(t, x) + 2u_x(t, x)v(t, x) + 2u(t, x)v_x(t, x) + u_{xxx}(t, x) = 0,$$

$(t, x) \in [0, \infty) \times \mathbb{R}$. We put $t = 0$ in both equations of (2.2) and we arrive at

$$u(0, x) = u_0(x), \quad v(0, x) = v_0(x), \quad x \in \mathbb{R}.$$

This completes the proof. □

Lemma 2.2. *Suppose (H1) and let $h \in \mathcal{C}([0, \infty) \times \mathbb{R})$ be a positive function almost everywhere on $[0, \infty) \times \mathbb{R}$. If $(u, v) \in X$ satisfies the following integral equations:*

$$\int_0^t \int_0^x (t - t_1)(x - x_1)^3 h(t_1, x_1) S_1^1(u, v)(t_1, x_1) dx_1 dt_1 = 0, \quad (t, x) \in [0, \infty) \times \mathbb{R},$$

and

$$\int_0^t \int_0^x (t - t_1)(x - x_1)^3 h(t_1, x_1) S_1^2(u, v)(t_1, x_1) dx_1 dt_1 = 0, \quad (t, x) \in [0, \infty) \times \mathbb{R},$$

then (u, v) is a solution to the IVP (1.1).

Proof. We differentiate three times with respect to t and three times with respect to x the integral equations of Lemma 2.2 and we find

$$h(t, x)S_1(u, v)(t, x) = 0, \quad (t, x) \in [0, \infty) \times \mathbb{R},$$

whereupon

$$S_1(u, v)(t, x) = 0, \quad (t, x) \in [0, \infty) \times \mathbb{R}.$$

Hence and Lemma 2.1, we conclude that (u, v) is a solution to the IVP (1.1). This completes the proof. \square

2.2. A priori estimates. In the sequel, $X = X^1 \times X^1$ where $X^1 = \mathcal{C}^1([0, \infty), \mathcal{C}^3(\mathbb{R}))$ will be endowed with the norm

$$\|(u, v)\| = \max \{\|u\|_{X^1}, \|v\|_{X^1}\}, \quad (u, v) \in X,$$

with

$$\|u\|_{X^1} = \max \left\{ \sup_{(t,x) \in [0,\infty) \times \mathbb{R}} |u(t, x)|, \sup_{(t,x) \in [0,\infty) \times \mathbb{R}} |u_t(t, x)|, \sup_{(t,x) \in [0,\infty) \times \mathbb{R}} |u_x(t, x)|, \right. \\ \left. \sup_{(t,x) \in [0,\infty) \times \mathbb{R}} |u_{xx}(t, x)|, \sup_{(t,x) \in [0,\infty) \times \mathbb{R}} |u_{xxx}(t, x)| \right\},$$

provided it exists. Let

$$B_1 = 4(B + B^2).$$

Lemma 2.3. *Under hypothesis (H1) and for $(u, v) \in X$ with $\|(u, v)\| \leq B$, the following estimates hold:*

$$|S_1^1(u, v)(t, x)| \leq B_1(1 + t), \quad (t, x) \in [0, \infty) \times \mathbb{R},$$

and

$$|S_1^2(u, v)(t, x)| \leq B_1(1 + t), \quad (t, x) \in [0, \infty) \times \mathbb{R}.$$

Proof. Suppose that (H1) is satisfied and let $(u, v) \in X$ with $\|(u, v)\| \leq B$.

(i) Estimation of $|S_1^1(u, v)(t, x)|$, $(t, x) \in [0, \infty) \times \mathbb{R}$:

$$\begin{aligned} |S_1^1(u, v)(t, x)| &= \left| u(t, x) - u_0(x) + \int_0^t (u(t_1, x)u_x(t_1, x) + v_x(t_1, x)) dt_1 \right| \\ &\leq |u(t, x)| + |u_0(x)| + \int_0^t (|u(t_1, x)||u_x(t_1, x)| + |v_x(t_1, x)|) dt_1 \\ &\leq 2B + (B + B^2)t \\ &\leq (3B + B^2)(1 + t) \\ &\leq B_1(1 + t). \end{aligned}$$

(ii) Estimation of $|S_1^2(u, v)(t, x)|$, $(t, x) \in [0, \infty) \times \mathbb{R}$:

$$\begin{aligned}
|S_1^2(u, v)(t, x)| &= \left| v(t, x) - v(0, x) + \int_0^x \left(u_x(t_1, x) + 2u_x(t_1, x)v(t_1, x) \right. \right. \\
&\quad \left. \left. + 2u(t_1, x)v_x(t_1, x) + u_{xxx}(t_1, x) \right) dx_1 dt_1 \right| \\
&\leq |v(t, x)| + |v(0, x)| + \int_0^x \left(|u_x(t_1, x)| + 2|u_x(t_1, x)||v(t_1, x)| \right. \\
&\quad \left. + 2|u(t_1, x)||v_x(t_1, x)| + |u_{xxx}(t_1, x)| \right) dx_1 dt_1 \\
&\leq 2B + (2B + 4B^2)t \\
&\leq 4(B + B^2)(1 + t) \\
&= B_1(1 + t).
\end{aligned}$$

This completes the proof. □

Suppose

(H2): $g \in \mathcal{C}([0, \infty) \times \mathbb{R})$ is a positive function almost everywhere on $[0, \infty) \times \mathbb{R}$ such that

$$16(1 + t)^2 \left(1 + |x| + x^2 + |x|^3 \right) \int_0^t \left| \int_0^x g(t_1, x_1) dx_1 \right| dt_1 \leq A,$$

$(t, x) \in [0, \infty) \times \mathbb{R}$, for some constant $A > 0$.

In the last section, we will give an example for a function g that satisfies **(H2)**. For $(u, v) \in X$, define the operators

$$\begin{aligned}
S_2^1(u, v)(t, x) &= \int_0^t \int_0^x (t - t_1)(x - x_1)^3 g(t_1, x_1) S_1^1(u, v)(t_1, x_1) dx_1 dt_1, \\
S_2^2(u, v)(t, x) &= \int_0^t \int_0^x (t - t_1)(x - x_1)^3 g(t_1, x_1) S_1^2(u, v)(t_1, x_1) dx_1 dt_1
\end{aligned}$$

and

$$(2.3) \quad S_2(u, v)(t, x) = \left(S_2^1(u, v)(t, x), S_2^2(u, v)(t, x) \right), \quad (t, x) \in [0, \infty) \times \mathbb{R}.$$

Lemma 2.4. *Under hypothesis **(H1)** and **(H2)** and for $(u, v) \in X$, with $\|(u, v)\| \leq B$, the following estimate holds:*

$$\|S_2(u, v)\| \leq AB_1.$$

Proof. Suppose **(H1)** and **(H2)** and let $(u, v) \in X$, with $\|(u, v)\| \leq B$.

(i) Estimation of $|S_2^1(u, v)(t, x)|$, $(t, x) \in [0, \infty) \times \mathbb{R}$:

$$\begin{aligned}
|S_2^1(u, v)(t, x)| &= \left| \int_0^t \int_0^x (t - t_1)(x - x_1)^3 g(t_1, x_1) S_1^1(u, v)(t_1, x_1) dx_1 dt_1 \right| \\
&\leq \int_0^t \left| \int_0^x (t - t_1) |x - x_1|^3 g(t_1, x_1) |S_1^1(u, v)(t_1, x_1)| dx_1 \right| dt_1 \\
&\leq B_1(1 + t) \int_0^t \left| \int_0^x (t - t_1) |x - x_1|^3 g(t_1, x_1) dx_1 \right| dt_1 \\
&\leq 8B_1(1 + t)^2 |x|^3 \int_0^t \left| \int_0^x g(t_1, x_1) dx_1 \right| dt_1 \\
&\leq 8B_1(1 + t)^2 (1 + |x| + x^2 + |x|^3) \int_0^t \left| \int_0^x g(t_1, x_1) dx_1 \right| dt_1 \\
&\leq AB_1.
\end{aligned}$$

(ii) Estimation of $\left| \frac{\partial}{\partial t} S_2^1(u, v)(t, x) \right|$, $(t, x) \in [0, \infty) \times \mathbb{R}$:

$$\begin{aligned}
\left| \frac{\partial}{\partial t} S_2^1(u, v)(t, x) \right| &= \left| \int_0^t \int_0^x (x - x_1)^3 g(t_1, x_1) S_1^1(u, v)(t_1, x_1) dx_1 dt_1 \right| \\
&\leq \int_0^t \left| \int_0^x |x - x_1|^3 g(t_1, x_1) |S_1^1(u, v)(t_1, x_1)| dx_1 \right| dt_1 \\
&\leq B_1(1 + t) \int_0^t \left| \int_0^x |x - x_1|^3 g(t_1, x_1) dx_1 \right| dt_1 \\
&\leq 8B_1(1 + t)^2 |x|^3 \int_0^t \left| \int_0^x g(t_1, x_1) dx_1 \right| dt_1 \\
&\leq 8B_1(1 + t)^2 (1 + |x| + x^2 + |x|^3) \int_0^t \left| \int_0^x g(t_1, x_1) dx_1 \right| dt_1 \\
&\leq AB_1.
\end{aligned}$$

(iii) Estimation of $\left| \frac{\partial}{\partial x} S_2^1(u, v)(t, x) \right|$, $(t, x) \in [0, \infty) \times \mathbb{R}$:

$$\begin{aligned}
\left| \frac{\partial}{\partial x} S_2^1(u, v)(t, x) \right| &= 3 \left| \int_0^t \int_0^x (t - t_1)(x - x_1)^2 g(t_1, x_1) S_1^1(u, v)(t_1, x_1) dx_1 dt_1 \right| \\
&\leq 3B_1(1 + t) \int_0^t \left| \int_0^x (t - t_1)(x - x_1)^2 g(t_1, x_1) dx_1 \right| dt_1 \\
&\leq 12B_1(1 + t)^2 x^2 \int_0^t \left| \int_0^x g(t_1, x_1) dx_1 \right| dt_1 \\
&\leq 12B_1(1 + t)^2 (1 + |x| + x^2 + |x|^3) \int_0^t \left| \int_0^x g(t_1, x_1) dx_1 \right| dt_1 \\
&\leq AB_1.
\end{aligned}$$

(iv) Estimation of $\left| \frac{\partial^2}{\partial x^2} S_2^1(u, v)(t, x) \right|$, $(t, x) \in [0, \infty) \times \mathbb{R}$:

$$\begin{aligned}
\left| \frac{\partial^2}{\partial x^2} S_2^1(u, v)(t, x) \right| &= 6 \left| \int_0^t \int_0^x (t - t_1)(x - x_1) g(t_1, x_1) S_1^1(u, v)(t_1, x_1) dx_1 dt_1 \right| \\
&\leq 6 \int_0^t \left| \int_0^x (t - t_1) |x - x_1| g(t_1, x_1) |S_1^1(u, v)(t_1, x_1)| dx_1 \right| dt_1 \\
&\leq 6B_1(1 + t) \int_0^t \left| \int_0^x (t - t_1) |x - x_1| g(t_1, x_1) dx_1 \right| dt_1 \\
&\leq 12B_1(1 + t)^2 |x| \int_0^t \left| \int_0^x g(t_1, x_1) dx_1 \right| dt_1 \\
&\leq 12B_1(1 + t)^2 (1 + |x| + x^2 + |x|^3) \int_0^t \left| \int_0^x g(t_1, x_1) dx_1 \right| dt_1 \\
&\leq AB_1.
\end{aligned}$$

(v) Estimation of $\left| \frac{\partial^3}{\partial x^3} S_2^1(u, v)(t, x) \right|$, $(t, x) \in [0, \infty) \times \mathbb{R}$:

$$\begin{aligned}
\left| \frac{\partial^3}{\partial x^3} S_2^1(u, v)(t, x) \right| &= 6 \left| \int_0^t \int_0^x (t - t_1) g(t_1, x_1) S_1^1(u, v)(t_1, x_1) dx_1 dt_1 \right| \\
&\leq 6 \int_0^t \left| \int_0^x (t - t_1) g(t_1, x_1) |S_1^1(u, v)(t_1, x_1)| dx_1 \right| dt_1 \\
&\leq 6B_1(1 + t) \int_0^t \left| \int_0^x (t - t_1) g(t_1, x_1) dx_1 \right| dt_1 \\
&\leq 12B_1(1 + t)^2 \int_0^t \left| \int_0^x g(t_1, x_1) dx_1 \right| dt_1 \\
&\leq 12B_1(1 + t)^2 (1 + |x| + x^2 + |x|^3) \int_0^t \left| \int_0^x g(t_1, x_1) dx_1 \right| dt_1 \\
&\leq AB_1.
\end{aligned}$$

Similarly, the same estimates (i)-(v) can be proved for the operator S_2^2 . Finally,

$$\|S_2(u, v)\| \leq AB_1.$$

This completes the proof. □

3. MAIN RESULTS

3.1. Existence of nonnegative solutions. The following theorem (see its proof in [17]) will be used to prove Theorem 3.2.

Theorem 3.1. *Let E be a Banach space and*

$$E_1 = \{x \in E : \|x\| \leq R\},$$

with $R > 0$. Consider two operators T and S , where

$$Tx = -\epsilon x, \quad x \in E_1,$$

with $\epsilon > 0$ and $S : E_1 \rightarrow E$ be continuous and such that

- (i) $(I - S)(E_1)$ resides in a compact subset of E and
- (ii) $\{x \in E : x = \lambda(I - S)x, \|x\| = R\} = \emptyset$ for any $\lambda \in (0, \frac{1}{\epsilon})$.

Then there exists $x^* \in E_1$ such that

$$Tx^* + Sx^* = x^*.$$

In the sequel, suppose that the constants B and A which appear in the conditions **(H1)** and **(H2)**, respectively, satisfy the following inequality:

(H3): $AB_1 < B$, where $B_1 = 4(B + B^2)$.

Our first main result for existence of classical solutions of the IVP (1.1) is as follows.

Theorem 3.2. *Assume that the hypotheses **(H1)**, **(H2)** and **(H3)** are satisfied. Then the IVP (1.1) has at least one nonnegative solution $(u, v) \in \mathcal{C}^1([0, \infty), \mathcal{C}^3(\mathbb{R})) \times \mathcal{C}^1([0, \infty), \mathcal{C}^3(\mathbb{R}))$.*

Proof. Choose $\epsilon \in (0, 1)$, such that $\epsilon B_1(1 + A) < B$.

For $(u, v) \in X = \mathcal{C}^1([0, \infty), \mathcal{C}^3(\mathbb{R})) \times \mathcal{C}^1([0, \infty), \mathcal{C}^3(\mathbb{R}))$, we will write

$$(u, v) \geq 0 \text{ if } u(t, x) \geq 0 \text{ and } v(t, x) \geq 0, \text{ for any } (t, x) \in [0, \infty) \times \mathbb{R}.$$

Let $\tilde{\tilde{Y}}$ denotes the set of all equi-continuous families in X with respect to the norm $\|\cdot\|$, $\tilde{\tilde{Y}} = \overline{\tilde{Y}}$ be the closure of \tilde{Y} , $\tilde{Y} = \tilde{Y} \cup \{(u_0, v_0)\}$ and

$$Y = \{(u, v) \in \tilde{Y} : (u, v) \geq 0, \|(u, v)\| \leq B\}.$$

Note that Y is a compact set in X . For $(u, v) \in X$, define the operators

$$T(u, v)(t, x) = -\epsilon(u, v)(t, x), \quad (t, x) \in [0, \infty) \times \mathbb{R},$$

$$S(u, v)(t, x) = (u, v)(t, x) + \epsilon(u, v)(t, x) + \epsilon S_2(u, v)(t, x), \quad (t, x) \in [0, \infty) \times \mathbb{R}.$$

For $(u, v) \in Y$ and by using Lemma 2.4, it follows that

$$\begin{aligned} \|(I - S)(u, v)\| &= \|\epsilon(u, v) - \epsilon S_2(u, v)\| \\ &\leq \epsilon\|(u, v)\| + \epsilon\|S_2(u, v)\| \\ &\leq \epsilon B_1 + \epsilon A B_1 \\ &= \epsilon B_1(1 + A) \\ &< B. \end{aligned}$$

Thus, $S : Y \rightarrow X$ is continuous and $(I - S)(Y)$ resides in a compact subset of X . Now, suppose that there is a $(u, v) \in X$ so that $\|(u, v)\| = B$ and

$$(u, v) = \lambda(I - S)(u, v)$$

or

$$\frac{1}{\lambda}(u, v) = (I - S)(u, v) = -\epsilon(u, v) - \epsilon S_2(u, v)$$

or

$$\left(\frac{1}{\lambda} + \epsilon\right)(u, v) = -\epsilon S_2(u, v),$$

for some $\lambda \in (0, \frac{1}{\epsilon})$. Hence, $\|S_2(u, v)\| \leq AB_1 < B$,

$$\epsilon B < \left(\frac{1}{\lambda} + \epsilon\right) B = \left(\frac{1}{\lambda} + \epsilon\right) \|(u, v)\| = \epsilon \|S_2(u, v)\| < \epsilon B,$$

which is a contradiction. In virtue of Theorem 3.1, the operator $T + S$ has a fixed point $(u^*, v^*) \in Y$. Therefore,

$$\begin{aligned} (u^*, v^*)(t, x) &= T(u^*, v^*)(t, x) + S(u^*, v^*)(t, x) \\ &= -\epsilon(u^*, v^*)(t, x) + (u^*, v^*)(t, x) + \epsilon(u^*, v^*)(t, x) + \epsilon S_2(u^*, v^*)(t, x), \end{aligned}$$

$(t, x) \in [0, \infty) \times \mathbb{R}$, whereupon

$$0 = S_2(u^*, v^*)(t, x), \quad (t, x) \in [0, \infty) \times \mathbb{R}.$$

Lemma 2.2 yields that (u^*, v^*) is a solution to the IVP (1.1). This completes the proof. \square

3.2. Multiplicity of nonnegative solutions. Let E be a real Banach space.

Definition 3.1. A closed, convex set \mathcal{P} in E is said to be cone if

- (a) $\alpha x \in \mathcal{P}$ for any $\alpha \geq 0$ and for any $x \in \mathcal{P}$;
- (b) $x, -x \in \mathcal{P}$ implies $x = 0$.

Definition 3.2. A mapping $K : E \rightarrow E$ is said to be completely continuous if it is continuous and maps bounded sets into relatively compact sets.

Definition 3.3. Let X and Y be real Banach spaces. A mapping $K : X \rightarrow Y$ is said to be expansive if there exists a constant $h > 1$ such that

$$\|Kx - Ky\|_Y \geq h\|x - y\|_X,$$

for any $x, y \in X$.

The following result (see details of its proof in [5] and [17]) will be used to prove Theorem 3.4.

Theorem 3.3. Let \mathcal{P} be a cone of a Banach space E ; Ω a subset of \mathcal{P} and U_1, U_2 and U_3 three open bounded subsets of \mathcal{P} such that $\overline{U_1} \subset \overline{U_2} \subset U_3$ and $0 \in U_1$. Assume that $T : \Omega \rightarrow \mathcal{P}$ is an expansive mapping, $S : \overline{U_3} \rightarrow E$ is a completely continuous and $S(\overline{U_3}) \subset (I - T)(\Omega)$. Suppose that $(U_2 \setminus \overline{U_1}) \cap \Omega \neq \emptyset$, $(U_3 \setminus \overline{U_2}) \cap \Omega \neq \emptyset$, and there exists $w_0 \in \mathcal{P} \setminus \{0\}$ such that the following conditions hold:

- (i) $Sx \neq (I - T)(x - \lambda w_0)$ for all $\lambda > 0$ and $x \in \partial U_1 \cap (\Omega + \lambda w_0)$;
- (ii) there exists $\varepsilon > 0$ such that $Sx \neq (I - T)(\lambda x)$ for all $\lambda \geq 1 + \varepsilon$, $x \in \partial U_2$ and $\lambda x \in \Omega$;

(iii) $Sx \neq (I - T)(x - \lambda w_0)$ for all $\lambda > 0$ and $x \in \partial U_3 \cap (\Omega + \lambda w_0)$.
Then $T + S$ has at least two non-zero fixed points $x_1, x_2 \in \mathcal{P}$ such that

$$x_1 \in \partial U_2 \cap \Omega \quad \text{and} \quad x_2 \in (\overline{U}_3 \setminus \overline{U}_2) \cap \Omega$$

or

$$x_1 \in (U_2 \setminus U_1) \cap \Omega \quad \text{and} \quad x_2 \in (\overline{U}_3 \setminus \overline{U}_2) \cap \Omega.$$

In the sequel, suppose that the constants B and A which appear in the conditions **(H1)** and **(H2)**, respectively, satisfy the following inequality:

(H4): $AB_1 < \frac{L}{5}$, where $B_1 = 4(B + B^2)$ and L is a positive constant that satisfies the following conditions:

$$r < L < R_1 \leq B, \quad R_1 > \left(\frac{2}{5m} + 1 \right) L,$$

with r and R_1 are positive constants and $m > 0$ is large enough.

Our second main result for existence and multiplicity of classical solutions of the IVP (1.1) is as follows.

Theorem 3.4. *Assume that the hypotheses **(H1)**, **(H2)** and **(H4)** are satisfied. Then the IVP (1.1) has at least two nonnegative solutions*

$$(u_1, v_1), (u_2, v_2) \in \mathcal{C}^1([0, \infty), \mathcal{C}^3(\mathbb{R})) \times \mathcal{C}^1([0, \infty), \mathcal{C}^3(\mathbb{R})).$$

Proof. Set $X = \mathcal{C}^1([0, \infty), \mathcal{C}^3(\mathbb{R})) \times \mathcal{C}^1([0, \infty), \mathcal{C}^3(\mathbb{R}))$ and let

$$\tilde{P} = \{(u, v) \in X : (u, v) \geq 0 \text{ on } [0, \infty) \times \mathbb{R}\}.$$

With \mathcal{P} we will denote the set of all equi-continuous families in \tilde{P} . For $(u, v) \in X$, define the operators

$$T_1(u, v)(t, x) = (1 + m\epsilon)(u, v)(t, x) - \left(\epsilon \frac{L}{10}, \epsilon \frac{L}{10} \right), \quad (t, x) \in [0, \infty) \times \mathbb{R},$$

$$S_3(u, v)(t, x) = -\epsilon S_2(u, v)(t, x) - m\epsilon(u, v)(t, x) - \left(\epsilon \frac{L}{10}, \epsilon \frac{L}{10} \right), \quad (t, x) \in [0, \infty) \times \mathbb{R},$$

where ϵ is a positive constant, $m > 0$ is large enough and the operator S_2 is given by formula (2.3). Note that any fixed point $(u, v) \in X$ of the operator $T_1 + S_3$ is a solution to the IVP (1.1). Now, let us define

$$U_1 = \mathcal{P}_r = \{(u, v) \in \mathcal{P} : \|(u, v)\| < r\},$$

$$U_2 = \mathcal{P}_L = \{(u, v) \in \mathcal{P} : \|(u, v)\| < L\},$$

$$U_3 = \mathcal{P}_{R_1} = \{(u, v) \in \mathcal{P} : \|(u, v)\| < R_1\},$$

$$\Omega = \overline{\mathcal{P}_{R_2}} = \{(u, v) \in \mathcal{P} : \|(u, v)\| \leq R_2\}, \quad \text{with } R_2 = R_1 + \frac{A}{m}B_1 + \frac{L}{5m}.$$

(a) Let $(u_1, v_1), (u_2, v_2) \in \Omega$, then

$$\|T_1(u_1, v_1) - T_1(u_2, v_2)\| = (1 + m\epsilon)\|(u_1, v_1) - (u_2, v_2)\|,$$

whereupon $T_1 : \Omega \rightarrow X$ is an expansive operator with a constant $h = 1 + m\epsilon > 1$.

(b) Let $(u, v) \in \overline{\mathcal{P}_{R_1}}$, then Lemma 2.4 yields

$$\|S_3(u, v)\| \leq \epsilon \|S_2(u, v)\| + m\epsilon \|(u, v)\| + \epsilon \frac{L}{10} \leq \epsilon \left(AB_1 + mR_1 + \frac{L}{10} \right).$$

Therefore, $S_3(\overline{\mathcal{P}_{R_1}})$ is uniformly bounded. Since $S_3 : \overline{\mathcal{P}_{R_1}} \rightarrow X$ is continuous, we have that $S_3(\overline{\mathcal{P}_{R_1}})$ is equi-continuous. Consequently, $S_3 : \overline{\mathcal{P}_{R_1}} \rightarrow X$ is completely continuous.

(c) Let $(u_1, v_1) \in \overline{\mathcal{P}_{R_1}}$ and set

$$(u_2, v_2) = (u_1, v_1) + \frac{1}{m} S_2(u_1, v_1) + \left(\frac{L}{5m}, \frac{L}{5m} \right).$$

Note that $S_2^1(u_1, v_1) + \frac{L}{5} \geq 0$, $S_2^2(u_1, v_1) + \frac{L}{5} \geq 0$ on $[0, \infty) \times \mathbb{R}$. We have $u_2, v_2 \geq 0$ on $[0, \infty) \times \mathbb{R}$ and

$$\|(u_2, v_2)\| \leq \|(u_1, v_1)\| + \frac{1}{m} \|S_2(u_1, v_1)\| + \frac{L}{5m} \leq R_1 + \frac{A}{m} B_1 + \frac{L}{5m} = R_2.$$

Therefore, $(u_2, v_2) \in \Omega$ and

$$-\epsilon m(u_2, v_2) = -\epsilon m(u_1, v_1) - \epsilon S_2(u_1, v_1) - \epsilon \left(\frac{L}{10}, \frac{L}{10} \right) - \epsilon \left(\frac{L}{10}, \frac{L}{10} \right)$$

or

$$(I - T_1)(u_2, v_2) = -\epsilon m(u_2, v_2) + \epsilon \left(\frac{L}{10}, \frac{L}{10} \right) = S_3(u_1, v_1).$$

Consequently, $S_3(\overline{\mathcal{P}_{R_1}}) \subset (I - T_1)(\Omega)$.

(d) Assume that for any $(w_0, z_0) \in \mathcal{P}^* = \mathcal{P} \setminus \{0\}$ there exist $\lambda \geq 0$ and $(u, v) \in \partial\mathcal{P}_r \cap (\Omega + \lambda(w_0, z_0))$ or $(u, v) \in \overline{\mathcal{P}_{R_1}} \cap (\Omega + \lambda(w_0, z_0))$ such that

$$S_3(u, v) = (I - T_1)((u, v) - \lambda(w_0, z_0)).$$

Then

$$-\epsilon S_2(u, v) - m\epsilon(u, v) - \epsilon \left(\frac{L}{10}, \frac{L}{10} \right) = -m\epsilon((u, v) - \lambda(w_0, z_0)) + \epsilon \left(\frac{L}{10}, \frac{L}{10} \right)$$

or

$$-S_2(u, v) = \lambda m(w_0, z_0) + \left(\frac{L}{5}, \frac{L}{5} \right).$$

Hence,

$$\|S_2 v\| = \left\| \lambda m(w_0, z_0) + \left(\frac{L}{5}, \frac{L}{5} \right) \right\| > \frac{L}{5}.$$

This is a contradiction.

(e) Let $\varepsilon_1 = \frac{2}{5m}$. Assume that there exist $(u_1, v_1) \in \partial\mathcal{P}_L$ and $\lambda_1 \geq 1 + \varepsilon_1$ such that $\lambda_1(u_1, v_1) \in \overline{\mathcal{P}_{R_1}}$ and

$$(3.1) \quad S_3(u_1, v_1) = (I - T_1)(\lambda_1(u_1, v_1)).$$

Since $(u_1, v_1) \in \partial\mathcal{P}_L$ and $\lambda_1(u_1, v_1) \in \overline{\mathcal{P}_{R_1}}$, it follows that

$$\left(\frac{2}{5m} + 1 \right) L < \lambda_1 L = \lambda_1 \|(u_1, v_1)\| \leq R_1.$$

Moreover,

$$-\epsilon S_2(u_1, v_1) - m\epsilon(u_1, v_1) - \epsilon\left(\frac{L}{10}, \frac{L}{10}\right) = -\lambda_1 m\epsilon(u_1, v_1) + \epsilon\left(\frac{L}{10}, \frac{L}{10}\right)$$

or

$$S_2(u_1, v_1) + \left(\frac{L}{5}, \frac{L}{5}\right) = (\lambda_1 - 1)m(u_1, v_1).$$

From here,

$$2\frac{L}{5} \geq \left\| S_2(u_1, v_1) + \left(\frac{L}{5}, \frac{L}{5}\right) \right\| = (\lambda_1 - 1)m\|(u_1, v_1)\| = (\lambda_1 - 1)mL$$

and

$$\frac{2}{5m} + 1 \geq \lambda_1,$$

which is a contradiction.

Therefore, all conditions of Theorem 3.4 hold. Hence, the IVP (1.1) has at least two solutions (u_1, v_1) and (u_2, v_2) so that

$$\|(u_1, v_1)\| = L < \|(u_2, v_2)\| \leq R_1$$

or

$$r \leq \|(u_1, v_1)\| < L < \|(u_2, v_2)\| \leq R_1.$$

□

4. AN EXAMPLE

Below, we will illustrate our main results. Let

$$h(s) = \log \frac{1 + s^{11}\sqrt{2} + s^{22}}{1 - s^{11}\sqrt{2} + s^{22}}, \quad l(s) = \arctan \frac{s^{11}\sqrt{2}}{1 - s^{22}}, \quad s \in \mathbb{R}, \quad s \neq \pm 1.$$

Then

$$\begin{aligned} h'(s) &= \frac{22\sqrt{2}s^{10}(1 - s^{22})}{(1 - s^{11}\sqrt{2} + s^{22})(1 + s^{11}\sqrt{2} + s^{22})}, \\ l'(s) &= \frac{11\sqrt{2}s^{10}(1 + s^{20})}{1 + s^{40}}, \quad s \in \mathbb{R}, \quad s \neq \pm 1. \end{aligned}$$

Therefore,

$$\begin{aligned} -\infty &< \lim_{s \rightarrow \pm\infty} (1 + s + s^2)h(s) < \infty, \\ -\infty &< \lim_{s \rightarrow \pm\infty} (1 + s + s^2)l(s) < \infty. \end{aligned}$$

Hence, there exists a positive constant C_1 so that

$$\begin{aligned} (1 + s + s^2 + s^3 + s^4 + s^5 + s^6) \left(\frac{1}{44\sqrt{2}} \log \frac{1 + s^{11}\sqrt{2} + s^{22}}{1 - s^{11}\sqrt{2} + s^{22}} + \frac{1}{22\sqrt{2}} \arctan \frac{s^{11}\sqrt{2}}{1 - s^{22}} \right) &\leq C_1, \\ (1 + s + s^2 + s^3 + s^4 + s^5 + s^6) \left(\frac{1}{44\sqrt{2}} \log \frac{1 + s^{11}\sqrt{2} + s^{22}}{1 - s^{11}\sqrt{2} + s^{22}} + \frac{1}{22\sqrt{2}} \arctan \frac{s^{11}\sqrt{2}}{1 - s^{22}} \right) &\leq C_1, \end{aligned}$$

$s \in \mathbb{R}$. Note that $\lim_{s \rightarrow \pm 1} l(s) = \frac{\pi}{2}$ and by [15, pp. 707, Integral 79], we have

$$\int \frac{dz}{1+z^4} = \frac{1}{4\sqrt{2}} \log \frac{1+z\sqrt{2}+z^2}{1-z\sqrt{2}+z^2} + \frac{1}{2\sqrt{2}} \arctan \frac{z\sqrt{2}}{1-z^2}.$$

Let

$$Q(s) = \frac{s^{10}}{(1+s^{44})(1+s+s^2)^2}, \quad s \in \mathbb{R},$$

and

$$g_1(t, x) = Q(t)Q(x), \quad t \in [0, \infty), \quad x \in \mathbb{R}.$$

Then there exists a constant $C_2 > 0$ such that

$$12(1+t)^2 (1+|x|+x^2+|x|^3) \int_0^t \left| \int_0^x g_1(t_1, x_1) dx_1 \right| dt_1 \leq C_2, \quad (t, x) \in [0, \infty) \times \mathbb{R}.$$

Let

$$g(t, x) = \frac{A}{C_2} g_1(t, x), \quad (t, x) \in [0, \infty) \times \mathbb{R}.$$

Then

$$12(1+t)^2 (1+|x|+x^2+|x|^3) \int_0^t \left| \int_0^x g(t_1, x_1) dx_1 \right| dt_1 \leq A, \quad (t, x) \in [0, \infty) \times \mathbb{R},$$

i.e., **(H2)** holds. Now, consider the initial value problem

$$\begin{aligned} u_t + uu_x + v_x &= 0, \quad t \in (0, \infty), \quad x \in \mathbb{R}, \\ v_t + u_x + 2(uv)_x + u_{xxx} &= 0, \quad t \in (0, \infty), \quad x \in \mathbb{R}, \\ (4.1) \quad u(0, x) &= \frac{1}{1+x^2+x^4}, \quad x \in \mathbb{R}, \\ v(0, x) &= \frac{1}{1+3x^2+x^8}, \quad x \in \mathbb{R}, \end{aligned}$$

so that **(H1)** holds, with $B = 10$, for example. Take

$$B = 10 \quad \text{and} \quad A = \frac{1}{10^4}.$$

Then

$$AB_1 = A \cdot 4(B + B^2) = \frac{1}{10^4} \cdot 4(10 + 10^2) < B.$$

So, condition **(H3)** is fulfilled. Thus, the conditions **(H1)**, **(H2)** and **(H3)** are satisfied. Hence, by Theorem 3.2, it follows that problem (4.1) has at least one solution $(u, v) \in \mathcal{C}^1([0, \infty), \mathcal{C}^3(\mathbb{R})) \times \mathcal{C}^1([0, \infty), \mathcal{C}^3(\mathbb{R}))$.

In the sequel, take

$$R_1 = B = 10, \quad L = 5, \quad r = 4, \quad m = 10^{50}, \quad A = \epsilon = \frac{1}{10^4}.$$

Clearly,

$$r < L < R_1 \leq B, \quad \epsilon > 0, \quad R_1 > \left(\frac{2}{5m} + 1\right)L, \quad AB_1 < \frac{L}{5},$$

i.e., **(H4)** holds. Hence, by Theorem 3.4, it follows that the initial value problem (4.1) has at least two nonnegative solutions $(u_1, v_1), (u_2, v_2) \in \mathcal{C}^1([0, \infty), \mathcal{C}^3(\mathbb{R})) \times \mathcal{C}^1([0, \infty), \mathcal{C}^3(\mathbb{R}))$.

Acknowledgements. The authors D. Boureni, A. Kheloufi and K. Mebarki acknowledge support of "Direction Générale de la Recherche Scientifique et du Développement Technologique (DGRSDT)", MESRS, Algeria.

REFERENCES

- [1] C. L. Chen and S. Y. Lou, *CTE solvability and exact solution to the Broer-Kaup system*, Chin. Phys. Lett. **30**(11) (2013), Article ID 110202, 4 pages. <https://doi.org/10.1088/0256-307X/30/11/110202>
- [2] L. J. F. Broer, *Approximate equations for long water waves*, Appl. Sci. Res. **31** (1975), 377–395. <https://doi.org/10.1007/BF00418048>
- [3] X. Q. Cao, S. C. Hou, Y. N. Guo, C. Z. Zhang and K. C. Peng, *Variational Principle for $(2+1)$ -dimensional Broer-Kaup equations with fractal derivatives*, Fractals **28**(7) (2020), Article ID 2050107, 13 pages. <https://doi.org/10.1142/S0218348X20501078>
- [4] G. Chavchanidze, *Non-Noether symmetries in Hamiltonian dynamical systems*, Mem. Differential Equations Math. Phys. **36** (2005), 81–134.
- [5] S. Djebali and K. Mebarki, *Fixed point index theory for perturbation of expansive mappings by k -set contractions*, Topol. Methods Nonlinear Anal. **54**(2A) (2019), 613–640. <https://doi.org/10.12775/TMNA.2019.055>
- [6] M. F. El-Sabbagh and S. I. El-Ganaini, *The He's variational principle to the Broer-Kaup (BK) and Whitham Broer-Kaup (WBK) systems*, International Mathematical Forum **7**(43) (2012), 2131–2141.
- [7] S. Guo, Y. Zhou and C. Zhao, *The improved $(\frac{G'}{G})$ -expansion method and its applications to the Broer-Kaup equations and approximate long water wave equations*, Appl. Math. Comput. **216**(7) (2010), 1965–1971. <https://doi.org/10.1016/j.amc.2010.03.026>
- [8] B. Jiang and Q.-S. Bi, *Peaked periodic wave solutions to the Broer-Kaup equation*, Commun. Theor. Phys. **67** (2017), 22–26. <https://doi.org/10.1088/0253-6102/67/1/22>
- [9] D. J. Kaup, *A higher order water wave equation and method for solving it*, Progress of Theoretical Physics **54** (1975), 396–408. <https://doi.org/10.1143/PTP.54.396>
- [10] S. Kumar, K. Singh and R. K. Gupta, *Painlevé analysis, Lie symmetries and exact solutions for $(2+1)$ -dimensional variable coefficients Broer-Kaup equations*, Commun. Nonlinear Sci. Numer. Simul. **17** (2012), 1529–1541. <https://doi.org/10.1016/j.cnsns.2011.09.003>
- [11] B. A. Kupershmidt, *Mathematics of dispersive water waves*, Commun. Math. Phys. **99**(1) (1985), 51–73. <https://doi.org/10.1007/BF01466593>
- [12] S. Y. Lou and X. B. Hu, *Broer-Kaup systems from Darboux transformation related symmetry constraints of Kadomtsev-Petviashvili equation*, Commun. Theor. Phys. **29**(1) (1998), 145–148. <https://doi.org/10.1088/0253-6102/29/1/145>
- [13] Q. Meng, W. Li and B. He, *Smooth and peaked solitary wave solutions of the Broer-Kaup system using the approach of dynamical system*, Commun. Theor. Phys. **62**(3) (2014), 308–314. <https://doi.org/10.1088/0253-6102/62/3/03>

- [14] A. Neirameh, *Solitary wave solutions of the BK equation and ALWW system by using the first integral method*, Comput. Methods Differ. Equ. **1**(2) (2013), 146–156.
- [15] A. Polyanin and A. Manzhirov, *Handbook of Integral Equations*, CRC Press, London, 1998.
- [16] J. Satsuma, K. Kajiwara, J. Matsukidaira and J. Hietarinta, *Solutions of the Broer-Kaup system through its trilinear form*, J. Phys. Soc. Jpn. **61**(9) (1992), 3096–3102. <https://doi.org/10.1143/JPSJ.61.3096>
- [17] S. Georgiev, A. Kheloufi and K. Mebarki, *Classical solutions for the Korteweg-De Vries equation*, New Trends Nonlinear Anal. Appl. (2022) (to appear).
- [18] A. K. Svinin, *Differential constraints for the Kaup-Broer system as a reduction of the 1D Toda lattice*, Inverse Probl. **17**(4) (2001), 1061–1066. <https://doi.org/10.1088/0266-5611/17/4/332>
- [19] M. Wang, J. Zhang and X. Li, *Application of the $(\frac{G'}{G})$ -expansion to travelling wave solutions of the Broer-Kaup and the approximate long water wave equations*, Appl. Math. Comput. **206** (2008), 321–326. <https://doi.org/10.1016/j.amc.2008.08.045>
- [20] H. Wang, H. Chen, Z. Ouyang and F. Li, *Lie symmetry analysis and conservation law for the equation arising from higher order Broer-Kaup equation*, Open Math. **17** (2019), 1045–1054. <https://doi.org/10.1515/math-2019-0080>
- [21] Z. Wen, *New exact explicit nonlinear wave solutions for the Broer-Kaup equation*, J. Appl. Math. **2014** (2014), Article ID 984791, 7 pages. <https://doi.org/10.1155/2014/984791>
- [22] S. Xie, X. Hong and J. Lu, *The bifurcation travelling waves of a generalized Broer-Kaup equation*, J. Appl. Anal. Comput. **11**(1) (2021), 210–226. <https://doi.org/10.11948/20190268>
- [23] X.-P. Xin, *Interaction solutions for $(1+1)$ -dimensional higher-order Broer-Kaup system*, Commun. Theor. Phys. **66** (2016), 479–482. <https://doi.org/10.1088/0253-6102/66/5/479>
- [24] H.-M. Yin, B. Tian, J. Chai and X.-Y. Wu, *Stochastic soliton solutions for the $(2+1)$ -dimensional stochastic Broer-Kaup equations in a fluid or plasma*, Appl. Math. Lett. **82** (2018), 126–131. <https://doi.org/10.1016/j.aml.2017.12.005>
- [25] Q. Zeng and Y. Li, *Application of the sub-ode method for the Broer-Kaup equation*, Advanced Materials Research **760–762** (2013), 1655–1660. <https://doi.org/10.4028/www.scientific.net/AMR.760-762.1655>
- [26] C.-L. Zheng and J.-X. Fei, *Complex wave excitations in generalized Broer-Kaup system*, Commun. Theor. Phys. **48**(4) (2007), 657–661. <https://doi.org/10.1088/0253-6102/48/4/018>
- [27] Zh. J. Zhou and Z. B. Li, *A Darboux transformation and new exact solutions for Broer-Kaup system*, Acta Phys. Sin. **52**(2) (2003), 262–266. <https://doi.org/10.7498/aps.52.262>
- [28] S. Zhu and J. Song, *Residual symmetries, n th Bäcklund transformation and interaction solutions for $(2+1)$ -dimensional generalized Broer-Kaup equations*, Appl. Math. Lett. **83** (2018), 33–39. <https://doi.org/10.1016/j.aml.2018.03.021>
- [29] J. Zhu and X. Wang, *Broer-Kaup system revisit: Inelastic interaction and blowup solutions*, J. Math. Anal. Appl. **496** (2021), Article ID 124794, 15 pages. <https://doi.org/10.1016/j.jmaa.2020.124794>

¹LABORATORY OF APPLIED MATHEMATICS,
BEJAIA UNIVERSITY,
06000 BEJAIA, ALGERIA
Email address: dalila.boureni@univ-bejaia.dz

²DEPARTMENT OF DIFFERENTIAL EQUATIONS, FACULTY OF MATHEMATICS AND INFORMATICS,
UNIVERSITY OF SOFIA,
SOFIA, BULGARIA
Email address: svetlinggeorgiev1@gmail.com
ORCID id: <https://orcid.org/0000-0001-8015-4226>

³LABORATORY OF APPLIED MATHEMATICS,
BEJAIA UNIVERSITY,
06000 BEJAIA, ALGERIA
Email address: arezki.kheloufi@univ-bejaia.dz; arezkinet2000@yahoo.fr
ORCID id: <https://orcid.org/0000-0001-5584-1454>

⁴LABORATORY OF APPLIED MATHEMATICS, FACULTY OF EXACT SCIENCES,
BEJAIA UNIVERSITY,
06000 BEJAIA, ALGERIA
Email address: karima.mebarki@univ-bejaia.dz; mebarqi_karima@hotmail.fr