

BERTRAND'S PARADOX: NEW PROBABILISTIC MODELS

ZORAN VIDOVIĆ¹

ABSTRACT. In this paper two new generating procedure of a random chord are obtained and thereby new solutions of Bertrand's paradox are proposed.

1. INTRODUCTION

Paradox, on its own, is a puzzle that confronts some already established principles. Bertrand's paradox was developed as a probability question that raised severe objections on the principle of indifference while dealing with geometrical probability. The question that defines this paradox: "What is the probability that a chord selected "at random" in a circle is larger than a side of the inscribed equilateral triangle?"

In [3], Bertrand obtained probabilities $1/3$, $1/2$ and $1/4$ by different random chord generation procedures: by choosing a chord with one end at a vertex of the inscribed equilateral triangle in a circle; by choosing a chord perpendicular to the diameter which is the right bisector of the equilateral triangle; and selecting a point inside a circle and denoting it as a chord midpoint, respectively. This puzzle has fascinated many since its discovery and a series of papers with outstanding solutions of this problem have been published, see e.g. [1, 2, 4–9]. Here, we provide two new models of random chord construction in a circle and obtain associated probabilities of Bertrand's paradox.

The paper is organized as follows. In Section 2, we propose two new procedures for generating a random chord in a circle and obtain probabilities of Bertrand's paradox for each case. Section 3 concludes this paper.

Key words and phrases. Bertrand's paradox, new solutions, Monte Carlo simulations.
2020 Mathematics Subject Classification. Primary: 60D05.
DOI 10.46793/KgJMat2501.061V
Received: July 09, 2021.
Accepted: November 19, 2021.

2. NEW MODELS AND SOLUTIONS

In [7], an attempt was made to look at classical models of Bertrand's paradox as limits of a continuous family of planar probabilistic models. Such family is seen by fixing a point, say A , at a distance $h > 1$ from a unit circle and constructing lines that intersect the circle and point A . However, this family of chord constructing models undermines the randomness selection of distance h and, so, it yields inappropriate results with respect to Bertrand's paradox. Motivated by this issue, in [10] a chord generating procedure is presented that overcomes this obstacle. Here, we additionally provide two new methods of generating random chords in a circle with the same intention.

For both models, we will denote X as the distance from the center of the circle and the chord and L as the corresponding chord length.

2.1. First model. The first method is obtained as follows.

- Step 1. Let a point A be such that its distance from the center of the circle OA is a random variable $Y \sim U(0, 1)$ and is lying on the x axis.
- Step 2. Using the circle invariance property we can obtain a point on a x axis, say P , so that the relation $OP \cdot OA = 1$ holds.
- Step 3. Angle ϕ is determined by the circle tangent and the x axis, with P as its vertex;
- Step 4. Select a line which is directed by an angle $\theta \in U(0, \phi)$, with P as its starting point. A chord is formed by its intersection with the circle (Figure 1.).

In this case, we have $X = \sqrt{1 - \frac{L^2}{4}}$, $\phi = \arcsin(Y)$ and $\theta = \arcsin\left(Y\sqrt{1 - \frac{L^2}{4}}\right)$.

Using transformation technique, the distribution function of L can be found as

$$\begin{aligned}
 F_I(l) &= \int_0^1 \int_0^l \frac{xy}{4 \arcsin(y) \sqrt{1 - \frac{x^2}{4}} \sqrt{1 - (1 - \frac{x^2}{4})y^2}} dx dy \\
 (2.1) \quad &= \int_0^1 \frac{\arcsin(y) - \arcsin\left(\frac{y\sqrt{4-l^2}}{2}\right)}{\arcsin(y)} dy, \quad 0 < l < 2.
 \end{aligned}$$

Integral (2.1) cannot be obtained explicitly, so we can only provide numerical solutions. For the Bertrand's case $l = \sqrt{3}$ we have

$$(2.2) \quad P\{L_I \geq \sqrt{3}\} = 1 - F_I(\sqrt{3}) = 0.4694.$$

2.2. Second model. The second method is obtained as follows.

- Step 1. Let a point A be determined by a random angle $\phi \sim U(0, \pi/2)$ on a circumference of a circle.
- Step 2. Let a tangent t of a circle be determined by point A .
- Step 3. Angle δ is determined by the circle tangent and the x axis, with P as its vertex.
- Step 4. Select a line which is directed by an angle $\theta \in U(0, \delta)$, with P as its starting point. A chord is formed by its intersection with the circle (Figure 2).




FIGURE 1. Solution I.




FIGURE 2. Solution II.

For this case, we have $X = \sqrt{1 - \frac{L^2}{4}}$ and $\sin \theta = \cos \phi \sqrt{1 - \frac{L^2}{4}}$. Further, the distribution function of L can be obtained as

$$\begin{aligned}
 F_{II}(l) &= \frac{2}{\pi} \int_0^{\pi/2} \int_0^l \frac{x \cos y}{4(\frac{\pi}{2} - y) \sqrt{1 - \frac{x^2}{4}} \sqrt{1 - (1 - \frac{x^2}{4}) \cos^2 y}} dx dy \\
 (2.3) \quad &= \frac{2}{\pi} \int_0^{\pi/2} \frac{\operatorname{arcsec} \left(\frac{2 \sec(y)}{\sqrt{4-l^2}} \right) - y}{\pi - 2y} dy, \quad 0 < l < 2.
 \end{aligned}$$

As above, integral (2.3) cannot be obtained explicitly, so we can obtain numerical solutions. For the Bertrand's case $l = \sqrt{3}$ we have

$$(2.4) \quad P \{L_{II} \geq \sqrt{3}\} = 1 - F_{II}(\sqrt{3}) = 0.4454.$$

3. CONCLUSION

Overall, in this paper we presented two new generating procedures of random chords in a circle. The distribution function (2.3) is also obtained in [10] using a different method of constructing random chords. The results presented in this paper extend those can be found in [4, 9, 10] on Bertrand's paradox.

In [6], procedures of chord construction were classified by disjoint procedures: (i) inside the circle, (ii) on the circle circumference and (iii) outside of the circle. Proposed generating models connect procedures (i), (ii) and (iii), and confronts such classification. This may be a motivation to overlook Bertrand's paradox in a quite different manner.

Acknowledgements. The author would like to thank the referees for their valuable comments and suggestions that significantly improved the quality of the paper.

REFERENCES

- [1] M. K. Arkadani and S. S. Wulff, *An extended problem to Bertrand's paradox*, Discuss. Math. Probab. Stat. **34**(1–2) (2014), 23–34. <http://eudml.org/doc/271059>
- [2] S. Bangu, *On Bertrand's paradox*, Analysis **70**(1) (2009), 30–35.
- [3] J. Bertrand, *Calcul des Probabilités*, Gautier-Villars et Fils, Paris, 1889.
- [4] S. S. Chiu and R. C. Larson, *Bertrand's paradox revisited: More lessons about that ambiguous word, random*, Journal of Industrial and System Engineering **3**(1) (2009), 1–26.
- [5] V. Jevremović and M. Obradović, *Bertrand's paradox: Is there anything else?*, Quality & Quantity **46**(6) (2012), 1709–1709. <https://doi.org/10.1007/s11135-011-9553-7>
- [6] L. Marinof, *A resolution of Bertrand's paradox*, Philosophy of Science **61**(1) (1992), 1–24. <https://www.jstor.org/stable/188286>
- [7] A. Soranzo and A. Volčič, *On the Bertrand's paradox*, Rend. Circ. Mat. Palermo **47**(3) (1998), 503–509.
- [8] G. J. Székely, *Paradoxes in probability theory and mathematical statistics*, Technical Report (1986).
- [9] Z. Vidović, *Limit distribution of maximal random chord length*, Journal of Applied Statistics and Probability **5**(2) (2016), 213–220. <https://doi.org/10.18576/jsap/050202>
- [10] Z. Vidović, *Random chord in a circle and Bertrand's paradox: New generation method, extreme behaviour and length moments*, Bull. Korean Math. Soc. **58**(2) (2021), 433–444. <https://doi.org/10.4134/BKMS.b200345>

¹TEACHER EDUCATION FACULTY,
UNIVERSITY OF BELGRADE,
KRALJICE NATALIJE 43, BELGRADE 11000, SERBIA
Email address: zoran.vidovic@uf.bg.ac.rs