

INVESTIGATIONS ON A RIEMANNIAN MANIFOLD WITH A SEMI-SYMMETRIC NON-METRIC CONNECTION AND GRADIENT SOLITONS

KRISHNENDU DE¹, UDAY CHAND DE², AND AYDIN GEZER³

ABSTRACT. This article carries out the investigation of a three-dimensional Riemannian manifold N^3 endowed with a semi-symmetric type non-metric connection. Firstly, we construct a non-trivial example to prove the existence of a semi-symmetric type non-metric connection on N^3 . It is established that a N^3 with the semi-symmetric type non-metric connection, whose metric is a gradient Ricci soliton, is a manifold of constant sectional curvature with respect to the semi-symmetric type non-metric connection. Moreover, we prove that if the Riemannian metric of N^3 with the semi-symmetric type non-metric connection is a gradient Yamabe soliton, then either N^3 is a manifold of constant scalar curvature or the gradient Yamabe soliton is trivial with respect to the semi-symmetric type non-metric connection. We also characterize the manifold N^3 with a semi-symmetric type non-metric connection whose metrics are Einstein solitons and m -quasi Einstein solitons of gradient type, respectively.

1. INTRODUCTION

In this paper, on a Riemannian manifold N^3 , we carry out an investigation of gradient solitons with a semi-symmetric type non-metric connection (briefly, *SSNMC*). Many years ago, on a differentiable manifold, Friedman and Schouten [11] presented the concept of semi-symmetric linear connection. After that in 1932, on a Riemannian manifold, Hayden [15] introduced the notion of metric connection with torsion. In 1970, a systematic investigation of semi-symmetric metric connection which plays a

Key words and phrases. Riemannian manifolds, gradient Ricci solitons, gradient Yamabe solitons, gradient Einstein solitons, m -quasi Einstein solitons.

2010 *Mathematics Subject Classification.* Primary: 53C05, 53C20, 53C25.

DOI 10.46793/KgJMat2503.387D

Received: March 01, 2022.

Accepted: April 26, 2022.

significant role in the study of Riemannian manifolds, was conducted by Yano [23]. In this connection, we may mention the work of Zengin et al. [24, 25].

On N^3 , a linear connection $\widehat{\nabla}$ is named semi-symmetric if \widehat{T} , the torsion tensor defined by

$$(1.1) \quad \widehat{T}(U, V) = \widehat{\nabla}_U V - \widehat{\nabla}_V U - [U, V]$$

obeys

$$(1.2) \quad \widehat{T}(U, V) = \psi(V)U - \psi(U)V,$$

where ψ is a 1-form defined by $\psi(U) = g(U, \xi)$, for a fixed vector field ξ (the associated vector field of $\widehat{\nabla}$). If in the right side of the equation (1.2) we substitute the independent vector fields U and V , respectively, by ϕU and ϕV , where ϕ is a $(1, 1)$ -tensor field [12], then the connection $\widehat{\nabla}$ transforms into a quarter-symmetric connection.

Again, if a semi-symmetric connection $\widehat{\nabla}$ on N^3 obeys

$$(1.3) \quad (\widehat{\nabla}_U g)(V, Y) = 0,$$

then $\widehat{\nabla}$ is called metric [23]. If $\widehat{\nabla}g \neq 0$, then it is called non-metric [15]. Here, we choose the *SSNMC*, that is, $\widehat{\nabla}g \neq 0$ and the connection $\widehat{\nabla}$ obeys the equation (1.2). The concept of the *SSNMC* on a Riemannian manifold was investigated in [1]. After that, several researchers investigated the properties of *SSNMC* on manifolds with different structures (see [6, 10, 18, 19]).

Hamilton [14] introduced the concept of Ricci flow as a solution to the challenge of obtaining a canonical metric on a smooth manifold. Ricci flow occurs when the metric of a Riemannian manifold N^3 is fulfilled by the evolution equation $\frac{\partial}{\partial t}g_{ij}(t) = -2\mathcal{S}_{ij}$, where \mathcal{S}_{ij} and g_{ij} are the components of the Ricci tensor and the metric tensor, respectively. Ricci solitons were created via self-similar solutions to the Ricci flow.

A metric of N^3 is named a Ricci soliton [13] if it fulfills

$$(1.4) \quad \mathfrak{L}_W g + 2\lambda g + 2\widehat{\mathcal{S}} = 0,$$

for some $\lambda \in \mathbb{R}$, the set of real numbers. Here, \mathfrak{L} being the Lie derivative operator and $\widehat{\mathcal{S}}$ is the Ricci tensor with respect to the non-metric connection $\widehat{\nabla}$. W is a complete vector field known as a potential vector field. The Ricci soliton is considered to be shrinking, expanding or steady depending on whether λ is negative, positive, or zero. If W is Killing or zero, the Ricci soliton is trivial and N^3 is Einstein. Also, if $W = Df$ for some smooth function f , then equation (1.4) turns into

$$(1.5) \quad \widehat{\nabla}^2 f + \widehat{\mathcal{S}} + \lambda g = 0,$$

where $\widehat{\nabla}^2$ and D indicate the Hessian and the gradient operator of g , respectively. The metric obeying the equation (1.5) is called a gradient Ricci soliton. Here, f is said to be the potential function of the gradient Ricci soliton.

On a complete Riemannian manifold N^3 , Hamilton [14] proposed the idea of Yamabe flow, which was inspired by Yamabe's conjecture ("metric of a complete Riemannian manifold is conformally connected to a metric with constant scalar curvature"). A

Riemannian manifold N^3 equipped with a Riemannian metric g is called a Yamabe flow if it obeys:

$$(1.6) \quad \frac{\partial}{\partial t}g(t) + rg(t) = 0, \quad g_0 = g(t),$$

where t indicates the time and r being the scalar curvature of N^3 . A Riemannian manifold N^3 equipped with a Riemannian metric g is named a Yamabe soliton if it fulfills

$$(1.7) \quad \mathfrak{L}_W g - 2(\hat{r} - \lambda)g = 0,$$

for real constant $\lambda : M \rightarrow \mathbb{R}$ and \hat{r} is the scalar curvature with respect to the non-metric connection $\widehat{\nabla}$. Here, W is called the potential vector field. In N^3 , with the condition $W = Df$, the Yamabe soliton reduces to the gradient Yamabe soliton. Thus, (1.7) takes the form

$$(1.8) \quad \widehat{\nabla}^2 f - (\hat{r} - \lambda)g = 0.$$

If f is constant (or, W is Killing) on M , then the soliton becomes trivial. The 3-Kenmotsu manifolds and almost co-Kähler manifolds with Yamabe solitons have been characterized by Wang [21] and Suh and De [20], respectively. Chen and Deshmukh [5, 9] studied the Yamabe solitons on Riemannian manifolds. Some interesting outcomes on this solitons have been investigated in [2, 3, 7, 8, 17] and also by others.

The notion of gradient Einstein soliton was presented in [4] and obeys

$$(1.9) \quad \hat{\mathfrak{S}} - \frac{1}{2}\hat{r}g + \widehat{\nabla}^2 f + \lambda g = 0,$$

where $\lambda \in \mathbb{R}$ is a constant and f indicates a smooth function.

A Riemannian manifold N^3 endowed with the Riemannian metric g is named a gradient m -quasi Einstein metric [4] if there exists a constant λ , a smooth function $f : N^3 \rightarrow \mathbb{R}$ and obeys

$$(1.10) \quad \hat{\mathfrak{S}} - \lambda g + \widehat{\nabla}^2 f - \frac{1}{m}df \otimes df = 0,$$

where \otimes indicate the tensor product and m is an integer. In this case f being the m -quasi Einstein potential function [4]. Here, the gradient m -quasi Einstein soliton is expanding for $\lambda > 0$, steady for $\lambda = 0$ and shrinking when $\lambda < 0$. If $m = \infty$, the foregoing equation represents a gradient Ricci soliton and the metric represents almost gradient Ricci soliton if it obeys the condition $m = \infty$ and λ is a smooth function. Few characterizations of the above metrics were characterized by He et al. [16].

The foregoing investigations motivate us to study the Riemannian manifold N^3 endowed with a $SSNMC$.

The content of the paper is laid out as: In Section 2, we produce the preliminary ideas of $SSNMC$. The existence of a $SSNMC$ on a Riemannian manifold are established in Section 3. The gradient Ricci soliton on N^3 equipped with a $SSNMC$ is investigated in Section 4. Section 5 concerns with gradient Yamabe soliton on N^3 with a $SSNMC$. We study the properties of N^3 with a $SSNMC$ whose metrics are

gradient Einstein solitons and gradient m -quasi Einstein solitons, in Section 6 and Section 7, respectively.

2. SEMI-SYMMETRIC NON-METRIC CONNECTION

A linear connection $\widehat{\nabla}$ on N , defined by

$$(2.1) \quad \widehat{\nabla}_U V = \nabla_U V + \psi(V)U,$$

∇ being the Levi-Civita connection, is a *SSNMC*. It also obeys

$$(2.2) \quad (\widehat{\nabla}_U g)(V, Y) = -\psi(V)g(U, Y) - \psi(Y)g(U, V).$$

Then \widehat{R} , the curvature tensor with respect to the *SSNMC*, $\widehat{\nabla}$, and R , the Riemannian curvature tensor are related by [1]

$$(2.3) \quad \widehat{R}(U, V)Y = R(U, V)Y - \alpha^*(V, Y)U + \alpha^*(U, Y)V,$$

for all U, V, Y on N^3 , where α^* is a $(0, 2)$ -tensor field defined by

$$(2.4) \quad \alpha^*(U, V) = (\nabla_U \psi)(V) - \psi(U)\psi(V).$$

Throughout this article, we choose that the vector field ξ is a unit parallel vector field with respect to the Levi-Civita connection ∇ . Then $\nabla_U \xi = 0$, which immediately implies

$$(2.5) \quad R(U, V)\xi = 0$$

and

$$(2.6) \quad \mathcal{S}(U, \xi) = 0.$$

Also, using $\nabla_U \xi = 0$, we obtain

$$(2.7) \quad (\nabla_U \psi)V = 0.$$

Hence, by the preceding equation, we get from (2.3)

$$(2.8) \quad \widehat{R}(U, V)Y = R(U, V)Y + \psi(Y)[\psi(V)U - \psi(U)V].$$

From the foregoing equation, we can easily have

$$(2.9) \quad \widehat{\mathcal{S}}(U, V) = \mathcal{S}(U, V) + 2\psi(U)\psi(V).$$

Contracting the above equation, we lead

$$(2.10) \quad \widehat{r} = r - 2,$$

since $\psi(\xi) = g(\xi, \xi) = 1$. Making use of (2.5), we infer from (2.8)

$$(2.11) \quad \widehat{R}(U, V)\xi = \psi(V)U - \psi(U)V.$$

Therefore, we obtain the subsequent relations

$$(2.12) \quad \psi(\widehat{R}(U, V)Y) = 0,$$

$$(2.13) \quad \widehat{\mathcal{S}}(U, \xi) = 2\psi(U), \quad \widehat{Q}\xi = 2\xi.$$

We first establish the subsequent lemma.

Lemma 2.1. *Let N^3 be a Riemannian manifold with a SSNMC, $\widehat{\nabla}$. Then we have*

$$(2.14) \quad \xi \widehat{r} = 0.$$

Proof. In N^3 , the Riemannian curvature tensor is expressed by

$$(2.15) \quad R(U, V)Y = g(V, Y)QU - g(U, Y)QV + \mathcal{S}(V, Y)U - \mathcal{S}(U, Y)V \\ - \frac{r}{2}[g(V, Y)U - g(U, Y)V].$$

Making use of (2.8) and (2.9), we acquire

$$(2.16) \quad \widehat{R}(U, V)Y - \psi(Y)[\psi(V)U - \psi(U)V] \\ = g(V, Y)[\widehat{Q}U - 2\xi\psi(U)] - g(U, Y)[\widehat{Q}V - 2\xi\psi(V)] + [\widehat{\mathcal{S}}(V, Y) - 2\psi(V)\psi(Y)]U \\ - [\widehat{\mathcal{S}}(U, Y) - 2\psi(U)\psi(Y)]V - \frac{r}{2}[g(V, Y)U - g(U, Y)V].$$

Putting $V = Y = \xi$, the foregoing equation yields

$$(2.17) \quad \widehat{Q}U = \left(\frac{\widehat{r}}{2} + 1\right)U - \left(\frac{\widehat{r}}{2} - 1\right)\psi(U)\xi.$$

Taking covariant derivative along V , we write

$$(2.18) \quad (\nabla_V \widehat{Q})U = \frac{(V\widehat{r})}{2}[U - \psi(U)\xi].$$

Contracting the foregoing equation we acquire the desired result. \square

The projective curvature tensor \widehat{P} of N^3 with respect to $\widehat{\nabla}$ is defined by

$$(2.19) \quad \widehat{P}(U, V)Y = \widehat{R}(U, V)Y - \frac{1}{2}[\widehat{\mathcal{S}}(V, Y)U - \widehat{\mathcal{S}}(U, Y)V].$$

Making use of (2.8) and (2.9), (2.19) reduces to

$$(2.20) \quad \widehat{P}(U, V)Y = P(U, V)Y,$$

where P represents the projective curvature tensor with respect to the Levi-Civita connection ∇ defined by

$$(2.21) \quad P(U, V)Y = R(U, V)Y - \frac{1}{2}[\mathcal{S}(V, Y)U - \mathcal{S}(U, Y)V].$$

Theorem 2.1. *If N^3 is endowed with a SSNMC $\widehat{\nabla}$, then the projective curvature tensor with respect to $\widehat{\nabla}$ and ∇ , respectively, coincide on N^3 .*

In differential geometry, the investigation of conformal curvature tensor performs a significant role. Also, it has various applications in applied physics and the other branches of modern sciences. Motivated by the above facts we investigate the properties of the conformal curvature tensor C . With respect to $\widehat{\nabla}$, the conformal curvature

tensor \widehat{C} is defined by

$$(2.22) \quad \begin{aligned} \widehat{C}(U, V)Y = & \widehat{R}(U, V)Y - [\widehat{S}(V, Y)U - \widehat{S}(U, Y)V + g(V, Y)\widehat{Q}U \\ & - g(U, Y)\widehat{Q}V] + \frac{\widehat{r}}{2}[g(V, Y)U - g(U, Y)V], \end{aligned}$$

for all U, V and Y on N^3 [22]. Utilizing (2.8) and (2.9) in (2.22), we obtain

$$(2.23) \quad \begin{aligned} \widehat{C}(U, V)Y = & C(U, V)Y - \widehat{\psi}(V)\widehat{\psi}(Y)U + \widehat{\psi}(U)\widehat{\psi}(Y)V \\ & + 2\xi g(V, Y)\widehat{\psi}(U) - 2\xi g(U, Y)\widehat{\psi}(V) + g(V, Y)U - g(U, Y)V, \end{aligned}$$

where C represents the conformal curvature tensor with respect to the Levi-Civita connection ∇ defined by

$$(2.24) \quad \begin{aligned} C(U, V)Y = & R(U, V)Y - [S(V, Y)U - S(U, Y)V + g(V, Y)QU - g(U, Y)QV] \\ & + \frac{r}{2}[g(V, Y)U - g(U, Y)V]. \end{aligned}$$

Putting $Y = \xi$ in (2.23), we get

$$(2.25) \quad \widehat{C}(U, V)\xi = C(U, V)\xi.$$

Hence, we have the subsequent theorem.

Theorem 2.2. *If N^3 is equipped with a SSNMC $\widehat{\nabla}$, then the the conformal curvature tensor with respect to $\widehat{\nabla}$ and ∇ , satisfy the relation (2.25).*

3. EXISTENCE OF A SEMI-SYMMETRIC TYPE NON-METRIC CONNECTION

Here we construct a non-trivial example of semi-symmetric type non-metric connection on a Riemannian manifold.

Example 3.1. Let us consider a three-dimensional differentiable manifold $N^3 = \{(u, v, w) \in \mathbb{R}^3, w \neq 0\}$, where (u, v, w) indicates the standard coordinate of \mathbb{R}^3 . Let us choose

$$(3.1) \quad k_1 = e^w \frac{\partial}{\partial u}, \quad k_2 = e^w \frac{\partial}{\partial v}, \quad k_3 = \frac{\partial}{\partial w}.$$

At each point of N^3 the preceding vector fields are linearly independent. Here we define the Riemannian metric g as

$$\begin{aligned} g(k_1, k_3) &= g(k_1, k_2) = g(k_2, k_3) = 0, \\ g(k_1, k_1) &= g(k_2, k_2) = g(k_3, k_3) = 1, \end{aligned}$$

ψ indicates a 1-form defined by $\psi(U) = g(U, \xi)$, where $\xi = k_3$. Hence, (N^3, g) is a three-dimensional Riemannian manifold. The Lie brackets are calculated as

$$\begin{aligned}
 [k_1, k_3] &= k_1 k_3 - k_3 k_1 \\
 &= e^w \frac{\partial}{\partial u} \left(\frac{\partial}{\partial w} \right) - \left(\frac{\partial}{\partial w} \right) \left(e^w \frac{\partial}{\partial u} \right) \\
 &= e^w \frac{\partial^2}{\partial u \partial w} - e^w \frac{\partial^2}{\partial w \partial u} - e^w \frac{\partial}{\partial u} \\
 &= -k_1.
 \end{aligned}
 \tag{3.2}$$

Similarly,

$$[k_1, k_2] = 0 \quad \text{and} \quad [k_2, k_3] = -k_2. \tag{3.3}$$

∇ , the Levi-Civita connection with respect to g , is obtained by

$$\begin{aligned}
 2g(\nabla_U V, Y) &= U g(V, Y) + V g(Y, U) - Y g(U, V) \\
 &\quad - g(U, [V, Y]) - g(V, [U, Y]) + g(Y, [U, V]),
 \end{aligned}
 \tag{3.4}$$

which is termed as Koszul's formula.

Making use of (3.4) we have

$$2g(\nabla_{k_1} k_3, k_1) = -2g(k_1, k_1). \tag{3.5}$$

Again by (3.4)

$$2g(\nabla_{k_1} k_3, k_2) = 0 = -2g(k_1, k_2) \tag{3.6}$$

and

$$2g(\nabla_{k_1} k_3, k_3) = 0 = -2g(k_1, k_3). \tag{3.7}$$

From (3.5), (3.6) and (3.7) we get

$$2g(\nabla_{k_1} k_3, U) = -2g(k_1, U),$$

for all $U \in \mathfrak{X}(N)$.

Thus, $\nabla_{k_1} k_3 = -k_1$. Therefore, (3.4) further gives

$$\begin{aligned}
 \nabla_{k_1} k_2 &= 0, \quad \nabla_{k_1} k_1 = k_3, \\
 \nabla_{k_2} k_3 &= -k_2, \quad \nabla_{k_2} k_2 = k_3, \quad \nabla_{k_2} k_1 = 0, \\
 \nabla_{k_3} k_3 &= 0, \quad \nabla_{k_3} k_2 = 0, \quad \nabla_{k_3} k_1 = 0.
 \end{aligned}
 \tag{3.8}$$

We know that

$$R(U, V)Y = \nabla_U \nabla_V Y - \nabla_V \nabla_U Y - \nabla_{[U, V]} Y, \tag{3.9}$$

where R is the Riemann curvature tensor. Utilizing the foregoing results and with the help of (3.9), we acquire

$$\begin{aligned}
 R(k_1, k_2)k_3 &= 0, \quad R(k_1, k_3)k_3 = -k_1, \\
 R(k_1, k_2)k_2 &= -k_1, \quad R(k_2, k_3)k_2 = k_3, \quad R(k_1, k_3)k_2 = 0,
 \end{aligned}$$

$$R(k_1, k_2)k_1 = k_2, \quad R(k_2, k_3)k_1 = 0, \quad R(k_1, k_3)k_1 = k_3.$$

Using the above expressions, the Ricci tensor can be obtained as

$$(3.10) \quad \mathcal{S}(k_1, k_1) = g(R(k_1, k_2)k_2, k_1) + g(R(k_1, k_3)k_3, k_1) = -2.$$

Similarly, we get

$$(3.11) \quad \mathcal{S}(k_2, k_2) = \mathcal{S}(k_3, k_3) = -2.$$

Therefore, the scalar curvature r is calculated as

$$(3.12) \quad r = \mathcal{S}(k_1, k_1) + \mathcal{S}(k_2, k_2) + \mathcal{S}(k_3, k_3) = -6.$$

Making use of the above expressions and using the equation (2.1), we have

$$(3.13) \quad \begin{aligned} \widehat{\nabla}_{k_1}k_3 &= 0, & \widehat{\nabla}_{k_1}k_2 &= 0, & \widehat{\nabla}_{k_1}k_1 &= k_3, \\ \widehat{\nabla}_{k_2}k_3 &= 0, & \widehat{\nabla}_{k_2}k_2 &= k_3, & \widehat{\nabla}_{k_2}k_1 &= 0, \\ \widehat{\nabla}_{k_3}k_3 &= k_3, & \widehat{\nabla}_{k_3}k_2 &= 0, & \widehat{\nabla}_{k_3}k_1 &= 0. \end{aligned}$$

From the last equation and using (1.2), we obtain $\widehat{T}(k_1, k_3) = k_1$ and $\psi(k_3)k_1 - \psi(k_1)k_3 = k_1$. Similarly, other components can be verified. Therefore, the linear connection $\widehat{\nabla}$ defined on (N^3, g) as (2.1), is a semi-symmetric connection. Also, we have

$$(3.14) \quad (\widehat{\nabla}_{k_1}g)(k_1, k_3) = -1 \neq 0.$$

Thus, the linear connection $\widehat{\nabla}$ is non-metric on (N^3, g) .

4. GRADIENT RICCI SOLITONS ON N^3 WITH A *SSNMC*

This section carries out the study of gradient Ricci solitons in N^3 with a *SSNMC*.

Let us choose that the soliton vector W of the Ricci soliton (g, W, λ) in N^3 with a *SSNMC* is a gradient of some smooth function f . Then using (1.5), we infer

$$(4.1) \quad \widehat{\nabla}_U Df = -\widehat{Q}U - \lambda U,$$

for all $U \in \mathfrak{X}(N)$. Making use of the above equation, the subsequent relation

$$(4.2) \quad \widehat{R}(U, V)Df = \widehat{\nabla}_U \widehat{\nabla}_V Df - \widehat{\nabla}_V \widehat{\nabla}_U Df - \widehat{\nabla}_{[U, V]} Df$$

yields

$$(4.3) \quad \widehat{R}(U, V)Df = (\widehat{\nabla}_U \widehat{Q})(V) - (\widehat{\nabla}_V \widehat{Q})(U).$$

The contraction of the preceding equation gives

$$(4.4) \quad \widehat{S}(U, Df) = -\frac{1}{2}(U\widehat{r}).$$

Again, from (2.17) we obtain

$$(4.5) \quad \widehat{S}(U, Df) = \left(\frac{\widehat{r}}{2} + 1\right)(Uf) - \left(\frac{\widehat{r}}{2} - 1\right)\psi(U)(\xi f).$$

Comparing the equations (4.4) and (4.5)

$$(4.6) \quad -\frac{1}{2}(U\hat{r}) = \left(\frac{\hat{r}}{2} + 1\right)(Uf) - \left(\frac{\hat{r}}{2} - 1\right)\psi(U)(\xi f).$$

Now, putting $U = \xi$ in (4.6), we find

$$(4.7) \quad \xi f = 0,$$

since $\xi\hat{r} = 0$.

Equation (4.3) gives

$$(4.8) \quad g(\hat{R}(U, V)\xi, Df) = 0.$$

Again, from equation (2.11) we infer that

$$(4.9) \quad g(\hat{R}(U, V)\xi, Df) = \psi(V)(Uf) - \psi(U)(Vf).$$

Comparing last two equations and putting $V = \xi$ and using $\xi f = 0$, we lead

$$(4.10) \quad Uf = 0,$$

which shows that $f = \text{constant}$. Making use of the fact that f is constant, equation (4.1) infers that the manifold is an Einstein manifold. Hence, the Riemannian manifold N^3 is of constant sectional curvature.

Theorem 4.1. *Let the soliton vector field W of the Ricci soliton (g, W, λ) in N^3 with a SSNMC be a gradient Ricci soliton. Then N^3 is a manifold of constant sectional curvature with respect to the SSNMC.*

5. GRADIENT YAMABE SOLITONS ON N^3 WITH A SSNMC

From equation (1.8), we find

$$(5.1) \quad \widehat{\nabla}_V Df = (\hat{r} - \lambda)V.$$

Differentiating (5.1) covariantly along the vector field U , we obtain

$$(5.2) \quad \widehat{\nabla}_U \widehat{\nabla}_V Df = (U\hat{r})V + (\hat{r} - \lambda)\widehat{\nabla}_U V.$$

Interchanging U and V in the above equation and then utilizing the preceding equation in $\hat{R}(U, V)Df = \widehat{\nabla}_U \widehat{\nabla}_V Df - \widehat{\nabla}_V \widehat{\nabla}_U Df - \widehat{\nabla}_{[U, V]} Df$, we lead

$$(5.3) \quad \hat{R}(U, V)Df = (U\hat{r})V - (V\hat{r})U.$$

Contracting the previous equation over U , we get

$$(5.4) \quad \hat{S}(V, Df) = -2(V\hat{r}).$$

Combining the last equation and (4.5), we infer

$$(5.5) \quad -2(U\hat{r}) = \left(\frac{\hat{r}}{2} + 1\right)(Uf) - \left(\frac{\hat{r}}{2} - 1\right)\psi(U)(\xi f).$$

Putting $U = \xi$ in the foregoing equation, we have

$$(5.6) \quad \xi f = 0,$$

since $\xi\hat{r} = 0$. Thus, from (5.5), we obtain

$$(5.7) \quad -2(U\hat{r}) = \left(\frac{\hat{r}}{2} + 1\right)(Uf).$$

Now, from equation (5.3) we find that

$$(5.8) \quad g(\hat{R}(U, V)\xi, Df) = \psi(U)(V\hat{r}) - \psi(V)(U\hat{r}).$$

Combining equation (4.9) and (5.8), we have

$$(5.9) \quad \psi(V)(Uf) - \psi(U)(Vf) = \psi(U)(V\hat{r}) - \psi(V)(U\hat{r}).$$

Setting $V = \xi$ in the previous equation gives

$$(5.10) \quad (U\hat{r}) = -(Uf).$$

Utilizing (5.10) in (5.7) we infer that

$$(5.11) \quad \left(\frac{\hat{r}}{2} - 1\right)(Uf) = 0,$$

which entails that either $\hat{r} = 2$ or $\hat{r} \neq 2$.

If $\hat{r} = 2$, then from (2.10) we infer that $r = 4$. Therefore, N^3 is of constant scalar curvature.

Next, we suppose that $\hat{r} \neq 2$, that is, $(Uf) = 0$, which implies f is a constant. Therefore, the gradient Yamabe soliton is trivial.

Hence, we state the result.

Theorem 5.1. *Let the Riemannian metric of N^3 with a SSNMC be the gradient Yamabe soliton. Then, either N^3 is a manifold of constant scalar curvature or the gradient Yamabe soliton is trivial with respect to the SSNMC.*

Also, if $\hat{r} = 2$, then using the equation (2.17) we acquire that the manifold is an Einstein manifold. Hence, the Riemannian manifold N^3 is of constant sectional curvature.

Corollary 5.1. *Let the Riemannian metric of N^3 with a SSNMC be the gradient Yamabe soliton. Then, either N^3 is a manifold of constant sectional curvature or the gradient Yamabe soliton is trivial with respect to the SSNMC.*

6. GRADIENT EINSTEIN SOLITONS ON N^3 WITH A SSNMC

Making use of (1.9), we have

$$(6.1) \quad \widehat{\nabla}_V Df = -\widehat{Q}V + \frac{\hat{r}}{2}V - \lambda V.$$

Differentiating (6.1) covariantly along U , we find

$$(6.2) \quad \widehat{\nabla}_U \widehat{\nabla}_V Df = -\widehat{\nabla}_U \widehat{Q}V + \frac{1}{2}(U\hat{r})V + \left(\frac{\hat{r}}{2} - \lambda\right)\widehat{\nabla}_U V.$$

Interchanging U and V and then making use of the above equation in $\widehat{R}(U, V)Df = \widehat{\nabla}_U \widehat{\nabla}_V Df - \widehat{\nabla}_V \widehat{\nabla}_U Df - \widehat{\nabla}_{[U, V]} Df$, we infer

$$(6.3) \quad \widehat{R}(U, V)Df = \frac{1}{2}[(U\widehat{r})V - (V\widehat{r})U] - (\widehat{\nabla}_U \widehat{Q})(V) + (\widehat{\nabla}_V \widehat{Q})(U).$$

Contracting the foregoing equation over U , we obtain

$$(6.4) \quad \widehat{S}(V, Df) = -\frac{1}{2}(V\widehat{r}).$$

Combining the last equation and (4.5), we get

$$(6.5) \quad -\frac{1}{2}(U\widehat{r}) = \left(\frac{\widehat{r}}{2} + 1\right)(Uf) - \left(\frac{\widehat{r}}{2} - 1\right)\psi(U)(\xi f).$$

Setting $U = \xi$ in (6.5), we have

$$(6.6) \quad (\xi f) = 0,$$

since $\xi\widehat{r} = 0$. Thus, from (6.5), we acquire

$$(6.7) \quad -\frac{1}{2}(U\widehat{r}) = \left(\frac{\widehat{r}}{2} + 1\right)(Uf).$$

Now, from equation (6.3) we obtain that

$$(6.8) \quad g(\widehat{R}(U, V)\xi, Df) = -\frac{1}{2}[\psi(U)(V\widehat{r}) - \psi(V)(U\widehat{r})].$$

Combining equation (4.9) and (6.8), we lead

$$(6.9) \quad \psi(V)(Uf) - \psi(U)(Vf) = -\frac{1}{2}[\psi(U)(V\widehat{r}) - \psi(V)(U\widehat{r})].$$

Putting $V = \xi$ in the last equation yields

$$(6.10) \quad (Uf) = -\frac{1}{2}(U\widehat{r}).$$

Using (6.10) in (6.7) we find that

$$(6.11) \quad \frac{\widehat{r}}{2}(Uf) = 0.$$

Hence, either $\widehat{r} = 0$ or $\widehat{r} \neq 0$.

If $\widehat{r} = 0$, then from (2.10) we acquire that $r = 2$. Therefore, N^3 is of constant scalar curvature.

Next, we suppose that $\widehat{r} \neq 0$, that is, $(Uf) = 0$, which implies f is a constant. Then, equation (6.1) reveals that N^3 is an Einstein manifold. Hence, N^3 is of constant sectional curvature, since the manifold is of dimension 3.

Thus, we state the subsequent.

Theorem 6.1. *If the Riemannian metric of N^3 with a SSNMC is a gradient Einstein soliton, then N^3 is either a manifold of constant scalar curvature or a manifold of constant sectional curvature with respect to the SSNMC.*

7. GRADIENT m -QUASI EINSTEIN SOLITONS ON N^3 WITH A $SSNMC$

Here, we investigate the m -quasi Einstein metric on N^3 with a $SSNMC$. Initially, we prove the following lemma.

Lemma 7.1. *In N^3 , we have the following:*

$$(7.1) \quad \begin{aligned} \hat{R}(U, V)Df = & (\widehat{\nabla}_V \hat{Q})U - (\widehat{\nabla}_U \hat{Q})V + \frac{\lambda}{m} \{(Vf)U - (Uf)V\} \\ & + \frac{1}{m} \{(Uf)\hat{Q}V - (Vf)\hat{Q}U\}, \end{aligned}$$

for all $U, V \in \mathfrak{X}(M)$.

Proof. Let the Riemannian metric of N^3 with a $SSNMC$ be a m -quasi Einstein metric. Therefore, the equation (1.10) can be represented as

$$(7.2) \quad \widehat{\nabla}_U Df = -\hat{Q}U + \frac{1}{m}g(U, Df)Df + \lambda U.$$

Covariant derivative of (7.2) along V yields

$$(7.3) \quad \widehat{\nabla}_V \widehat{\nabla}_U Df = -\widehat{\nabla}_V \hat{Q}U + \frac{1}{m}\widehat{\nabla}_V g(U, Df)Df + \frac{1}{m}g(U, Df)\widehat{\nabla}_V Df + \lambda \widehat{\nabla}_V U.$$

Exchanging U and V in (7.3), we obtain

$$(7.4) \quad \widehat{\nabla}_U \widehat{\nabla}_V Df = -\widehat{\nabla}_U \hat{Q}V + \frac{1}{m}\widehat{\nabla}_U g(V, Df)Df + \frac{1}{m}g(V, Df)\widehat{\nabla}_U Df + \lambda \widehat{\nabla}_U V$$

and

$$(7.5) \quad \widehat{\nabla}_{[U, V]} Df = -\hat{Q}[U, V] + \frac{1}{m}g([U, V], Df)Df + \lambda[U, V].$$

Utilizing (7.2)–(7.5) and the relation $\hat{R}(U, V)Df = \widehat{\nabla}_U \widehat{\nabla}_V Df - \widehat{\nabla}_V \widehat{\nabla}_U Df - \widehat{\nabla}_{[U, V]} Df$, we have

$$\begin{aligned} \hat{R}(U, V)Df = & (\widehat{\nabla}_V \hat{Q})U - (\widehat{\nabla}_U \hat{Q})V + \frac{\lambda}{m} \{(Vf)U - (Uf)V\} \\ & + \frac{1}{m} \{(Uf)\hat{Q}V - (Vf)\hat{Q}U\}. \end{aligned} \quad \square$$

Now contracting the equation (7.1) over U , we obtain

$$(7.6) \quad \hat{S}(V, Df) = \frac{1}{2}(V\hat{r}) + \frac{2\lambda}{m}(Vf) - \frac{1}{m} \left\{ \left(\frac{\hat{r}}{2} + 3 \right) (Vf) + \left(\frac{\hat{r}}{2} - 1 \right) (\xi f)\psi(V) \right\}.$$

Combining (7.6) and (4.5), we have

$$(7.7) \quad \begin{aligned} & \frac{1}{2}(V\hat{r}) + \frac{2\lambda}{m}(Vf) - \frac{1}{m} \left\{ \left(\frac{\hat{r}}{2} + 3 \right) (Vf) + \left(\frac{\hat{r}}{2} - 1 \right) (\xi f)\psi(V) \right\} \\ & = \left(\frac{\hat{r}}{2} + 1 \right) (Vf) - \left(\frac{\hat{r}}{2} - 1 \right) \psi(V)(\xi f). \end{aligned}$$

Setting $V = \xi$ in (7.7), we obtain

$$(7.8) \quad (2m + \hat{r} - 2\lambda + 2)(\xi f) = 0,$$

since $\xi\hat{r} = 0$.

Now, from equation (7.1) we have

$$(7.9) \quad g(\hat{R}(U, V)\xi, Df) = \left(\frac{\lambda}{m} - \frac{2}{m}\right) [\psi(V)(Uf) - \psi(U)(Vf)].$$

Combining equations (4.9) and (7.9), we find that

$$(7.10) \quad \psi(V)(Uf) - \psi(U)(Vf) = \left(\frac{\lambda}{m} - \frac{2}{m}\right) [\psi(V)(Uf) - \psi(U)(Vf)].$$

Putting $V = \xi$ in the foregoing equation yields

$$(7.11) \quad (\lambda - m - 2)(Uf) = 0,$$

where we have used $\xi f = 0$.

Hence, either $(\lambda - m - 2) = 0$ or $(\lambda - m - 2) \neq 0$.

If $(\lambda - m - 2) = 0$, then we get $\lambda = m + 2 =$ positive integer. Hence, the gradient m -quasi Einstein soliton is expanding.

If we suppose that $(\lambda - m - 2) \neq 0$, then $(Uf) = 0$, which implies f is a constant. Then, equation (7.1) reveals that N^3 is an Einstein manifold. Hence, N^3 is of constant sectional curvature, since the manifold is of dimension 3.

Hence, we state the following.

Theorem 7.1. *If the Riemannian metric of N^3 with a SSNMC is a gradient m -quasi Einstein soliton, then either the soliton is expanding or it is a manifold of constant sectional curvature with respect to the SSNMC, provided $(2m + \hat{r} - 2\lambda + 2) \neq 0$.*

8. ACKNOWLEDGEMENT

We would like to thank the referee for reviewing the paper carefully and his or her valuable comments to improve the quality of the paper.

REFERENCES

- [1] S. Agashe and M. R. Chafle, *A semi-symmetric non-metric connection on a Riemannian manifold*, Indian J. Pure Appl. Math. **23** (1992), 399–409.
- [2] A. M. Blaga, *Solitons and geometrical structures in a perfect fluid spacetime*, Rocky Mountain J. Math. **50** (2020), 41–53.
- [3] A. M. Blaga and C. Özgür, *Almost η -Ricci and almost η -Yamabe solitons with torse forming potential vector field*, Quaest. Math. (2020), 1–21.
- [4] A. Barros and J. N. Gomes, *A compact gradient generalized quasi-Einstein metric with constant scalar curvature*, J. Math. Anal. Appl. **401** (2013), 702–705.
- [5] B. Y. Chen and S. Deshmukh, *Yamabe and quasi-Yamabe solitons on Euclidean submanifolds*, Mediterr. J. Math. **15**(194) (2018), 8 pages.
- [6] A. De, *On almost pseudo symmetric manifolds admitting a type of semi-symmetric non-metric connection*, Acta Math. Hungar. **125**(1–2) (2009), 183–190.
- [7] K. De and U. C. De, *A note on gradient solitons on para-Kenmotsu manifolds*, Int. J. Geom. Methods Mod. Phys. **18**(1) (2021), Article ID 2150007, 11 pages.
- [8] K. De and U. C. De, *Almost quasi-Yamabe solitons and gradient almost quasi-Yamabe solitons in paracontact geometry*, Quaest. Math. **44**(11) (2021), 1429–1440.

- [9] S. Deshmukh and B. Y. Chen, *A note on Yamabe solitons*, Balkan J. Geom. Appl. **23**(1) (2018), 37–43.
- [10] Y. Dogru, C. Özgür and C. Murathan, *Riemannian manifolds with a semi-symmetric nonmetric connection satisfying some semisymmetry conditions*, Bull. Math. Anal. Appl. **3**(2) (2011), 206–212.
- [11] A. Friedmann and J. A. Schouten, *Über die geometrie der holbsymmetrischen ubertragungen*, Math. Z. **21** (1924), 211–233.
- [12] S. Golab, *On semi-symmetric and quarter-symmetric linear connections*, Tensor (N.S.) **29** (1975), 249–254.
- [13] R. S. Hamilton, *Three-manifolds with positive Ricci curvature*, J. Differential Geom. **17** (1982), 255–306.
- [14] R. S. Hamilton, *The Ricci flow on surfaces*, Contemp. Math. **71** (1988), 237–261.
- [15] H. A. Hayden, *Subspace of space with torsion*, Proc. Lon. Math. Soc. **34** (1932), 27–50.
- [16] C. He, P. Petersen and W. Wylie, *On the classification of warped product Einstein metrics*, Comm. Anal. Geom. **20** (2012), 271–312.
- [17] S. Y. Hsu, *A note on compact gradient Yamabe solitons*, J. Math. Anal. Appl. **388** (2012), 725–726.
- [18] C. Özgür and S. Sular, *Warped products with a semi-symmetric non-metric connection*, Arab J. Math. Sci. **36** (2011), 461–473.
- [19] J. Sengupta, U. C. De and T. Q. Binh, *On a type of semi-symmetric non-metric connection on a Riemannian manifold*, Indian J. Pure Appl. Math. **31**(12) (2000), 1659–1670.
- [20] Y. J. Suh and U. C. De, *Yamabe solitons and Ricci solitons on almost co-Kähler manifolds*, Canad. Math. Bull. **62** (2019), 653–661.
- [21] Y. Wang, *Yamabe solitons on three-dimensional Kenmotsu manifolds*, Bull. Belg. Math. Soc. Simon Stevin **23** (2016), 345–355.
- [22] H. Weyl, *Reine Infinitesimal geometrie*, Math. Z. **2**(1918), 384–411.
- [23] K. Yano, *On semi-symmetric metric connection*, Rev. Roumaine Math. Pures Appl. **15** (1970), 1579–1586.
- [24] F. O. Zengin, S. A. Demirbag, S. A. Uysal and H. B. Yilmaz, *Some vector fields on a Riemannian manifold with semi-symmetric metric connection*, Bull. Iranian Math. Soc. **38** (2012), 479–490.
- [25] F. O. Zengin, S. A. Uysal and S. A. Demirbag, *On sectional curvature of a Riemannian manifold with semi-symmetric metric connection*, Ann. Polon. Math. **101** (2011), 131–138.

¹DEPARTMENT OF MATHEMATICS,
KABI SUKANTA MAHAVIDYALAYA,
BHADRESWAR, P.O.-ANGUS, HOOGHLY,
PIN 712221, WEST BENGAL, INDIA
Email address: krishnendu.de@outlook.in

²DEPARTMENT OF PURE MATHEMATICS,
UNIVERSITY OF CALCUTTA
35, BALLYGUNGE CIRCULAR ROAD
KOL- 700019, WEST BENGAL, INDIA
Email address: uc_de@yahoo.com

³DEPARTMENT OF MATHEMATICS,
ATATURK UNIVERSITY,
ERZURUM-TURKEY
Email address: aydingzr@gmail.com