

QUANTITATIVE UNCERTAINTY PRINCIPLES FOR THE CANONICAL FOURIER BESSEL TRANSFORM

KHALED HLEILI¹ AND MANEL HLEILI²

ABSTRACT. The aim of this paper is to prove new uncertainty principles for the Canonical Fourier Bessel transform. To do so we prove a quantitative uncertainty inequality about the essential supports of a nonzero function for this transformation.

1. INTRODUCTION

The classical linear canonical transform (LCT) is considered as a generalization of the Fourier transform, and was first proposed in the 1970s by Collins [5] and Moshinsky and Quesne [26]. Very recently, many works have been devoted the LCT under many different names and in different contexts. Namely, in [22] the LCT is known as the generalized Fresnel transform, in [4] is called ABCD transform and in [1] is also called the special affine Fourier transform. Also, the LCT has been studied by many authors for various Fourier transforms, for examples [11, 23, 34]. In [11], the authors introduced the Dunkl linear canonical transform (DLCT) which is a generalization of the LCT in the framework of Dunkl transform [7]. DLCT includes many well-known transforms such as the Dunkl transform [7, 10] and the canonical Fourier Bessel transform [8, 11]. The LCT plays an important role in many fields of optics, radar system analysis, GRIN medium system analysis, filter design, phase retrieval, pattern recognition and many others [3, 28, 29]. In [8] the authors established some important properties of the Canonical Fourier Bessel transform (QFBT) such as Riemann-Lebesgue lemma, inversion formula, Plancherel theorem and some uncertainty principles.

Key words and phrases. Canonical Fourier Bessel transform, Donoho-Stark's uncertainty principle, Matolcsi-Szücs-type inequality.

2020 *Mathematics Subject Classification.* Primary: 42A38.

DOI 10.46793/KgJMat2504.567H

Received: March 16, 2021.

Accepted: July 30, 2022.

On the other hand, the uncertainty principle plays one important role in signal processing. It describes a function and its Fourier transform, which cannot both be simultaneously sharply localized. If we try to limit the behaviour of one we lose control of the other. Many of these uncertainty principles have already been studied from several points of view for the Fourier transform, such as Heisenberg-Pauli-Weyl inequality [6] and local uncertainty inequality [30]. Uncertainty principles have implications in two main areas: quantum physics and signal analysis. In quantum physics, they tell us that a particle’s speed and position cannot both be measured with arbitrary precision. In signal analysis, they tell us that if we observe a signal only for a finite period of time, we will lose information about the frequencies the signal consists of. Timelimited functions and bandlimited functions are basic tools of signal and image processing. Unfortunately, the simplest form of the uncertainty principle tells us that a signal cannot be simultaneously time and bandlimited. This leads to the investigation of the set of almost time and almost bandlimited functions, which has been initially carried through Landau, Pollak [24, 25] and then by Donoho, Stark [9]. In recent past, many works have been devoted to establish some uncertainty principles in different setting and for various transforms (see for example [2, 12–21, 31]) and others.

The purpose of this paper is to obtain uncertainty principle similar to Donoho-Stark’s principle for the QFBT.

In order to describe our results, we first need to introduce some facts about harmonic analysis related to Canonical Fourier Bessel transform. For more details, see [8].

Throughout this paper, α denotes a real number such that $\alpha \geq -\frac{1}{2}$. We use the following notation.

- $C_{e,0}(\mathbb{R})$ denotes the space of even continuous functions on \mathbb{R} and vanishing at infinity. We provide $C_{e,0}(\mathbb{R})$ with the topology of uniform convergence.
- $L^{p,\alpha}$ denotes the Lebesgue space of measurable functions f on \mathbb{R}_+ , such that

$$\|f\|_{p,\alpha} = \left(\int_0^{+\infty} |f(y)|^p y^{2\alpha+1} dy \right)^{\frac{1}{p}} < +\infty, \quad \text{if } 1 \leq p < +\infty,$$

$$\|f\|_{\infty,\alpha} = \text{ess sup}_{y \in \mathbb{R}_+} |f(y)| < +\infty, \quad \text{if } p = +\infty.$$

We provide $L^{p,\alpha}$ with the topology defined by the norm $\|\cdot\|_{p,\alpha}$.

- $L^{2,\alpha}$ denotes the Hilbert space equipped with the inner product $\langle \cdot, \cdot \rangle$ given by

$$\langle f, g \rangle = \int_0^{+\infty} f(y) \overline{g(y)} y^{2\alpha+1} dy.$$

- $m = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ is an arbitrary matrix in $SL(2, \mathbb{R})$, such that $b \neq 0$.

Definition 1.1. The canonical Fourier Bessel transform of a function $f \in L^{1,\alpha}$ is defined by

$$\mathcal{F}_\alpha^m(f)(x) = \frac{c_\alpha}{(ib)^{\alpha+1}} \int_0^{+\infty} K_\alpha^m(x, y) f(y) y^{2\alpha+1} dy,$$

where

$$(1.1) \quad c_\alpha = \frac{1}{2^\alpha \Gamma(\alpha + 1)}$$

and

$$K_\alpha^m(x, y) = e^{\frac{i}{2}\left(\frac{dx^2}{b} + \frac{ay^2}{b}\right)} j_\alpha\left(\frac{xy}{b}\right).$$

Here j_α denotes the normalized Bessel function of order $\alpha \geq -\frac{1}{2}$ and defined by [33]

$$j_\alpha(z) = 2^\alpha \Gamma(\alpha + 1) \frac{J_\alpha(z)}{z^\alpha} = \Gamma(\alpha + 1) \sum_{k=0}^{+\infty} \frac{(-1)^k}{k! \Gamma(\alpha + 1 + k)} \left(\frac{z}{2}\right)^{2k}, \quad z \in \mathbb{C}.$$

Proposition 1.1 ([8]). *We denote by Δ_α^m the differential operator*

$$\Delta_\alpha^m = \frac{d^2}{dx^2} + \left(\frac{2\alpha + 1}{x} - 2i\frac{d}{b}x\right) \frac{d}{dx} - \left(\frac{d^2}{b^2}x^2 + 2i(\alpha + 1)\frac{d}{b}\right).$$

(1) *For each $y \in \mathbb{R}$, the kernel $K_\alpha^m(\cdot, y)$ of the canonical Fourier Bessel transform \mathcal{F}_α^m is the unique solution of*

$$\begin{cases} \Delta_\alpha^m K_\alpha^m(\cdot, y) = \frac{-y^2}{b^2} K_\alpha^m(\cdot, y), \\ K_\alpha^m(0, y) = e^{\frac{ia y^2}{2b}}, \\ \frac{d}{dx} K_\alpha^m(0, y) = 0. \end{cases}$$

(2) *For each $x, y \in \mathbb{R}$ the kernel K_α^m has the following integral representation*

$$K_\alpha^m(x, y) = \begin{cases} \frac{2\Gamma(\alpha+1)}{\sqrt{\pi}\Gamma(\alpha+\frac{1}{2})} e^{\frac{i}{2}\left(\frac{dx^2}{b} + \frac{ay^2}{b}\right)} \int_0^1 (1-t^2)^{\alpha-\frac{1}{2}} \cos\left(\frac{xyt}{b}\right) dt, & \text{if } \alpha > -\frac{1}{2}, \\ e^{\frac{i}{2}\left(\frac{dx^2}{b} + \frac{ay^2}{b}\right)} \cos\left(\frac{xy}{b}\right), & \text{if } \alpha = -\frac{1}{2}. \end{cases}$$

In particular, we have

$$(1.2) \quad |K_\alpha^m(x, y)| \leq 1 \quad \text{for all } x, y \in \mathbb{R}.$$

Theorem 1.1 ([8]). (1) *(Plancherel theorem) If $f \in L^{1,\alpha} \cap L^{2,\alpha}$, then $\mathcal{F}_\alpha^m(f) \in L^{2,\alpha}$ and*

$$(1.3) \quad \|\mathcal{F}_\alpha^m(f)\|_{2,\alpha} = \|f\|_{2,\alpha}.$$

(2) *(Orthogonality relation) For every $f, g \in L^{2,\alpha}$, we have*

$$(1.4) \quad \langle f, g \rangle = \langle \mathcal{F}_\alpha^m(f), \mathcal{F}_\alpha^m(g) \rangle.$$

(3) *(The reversibility property) For all $f \in L^{1,\alpha}$, with $\mathcal{F}_\alpha^m \in L^{1,\alpha}$, we have*

$$(1.5) \quad (\mathcal{F}_\alpha^m \circ \mathcal{F}_\alpha^{m^{-1}})(f) = (\mathcal{F}_\alpha^{m^{-1}} \circ \mathcal{F}_\alpha^m)(f) = f, \quad \text{a.e.}$$

Babenko-Beckner inequality. Let $m = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ an arbitrary matrix in $SL(2, \mathbb{R})$, such that $b \neq 0$. Let p and q be real numbers such that $1 < p \leq 2$

and $\frac{1}{p} + \frac{1}{q} = 1$. Then, \mathcal{F}_α^m extends to a bounded linear operator on $L^{p,\alpha}$, $\alpha \geq -\frac{1}{2}$ and we have

$$(1.6) \quad \|\mathcal{F}_\alpha^m(f)\|_{q,\alpha} \leq |b|^{(\alpha+1)(\frac{2}{q}-1)} \left(\frac{(c_\alpha p)^{\frac{1}{p}}}{(c_\alpha q)^{\frac{1}{q}}} \right)^{\alpha+1} \|f\|_{p,\alpha},$$

where c_α is the constant given by (1.1).

Riemann-Lebesgue lemma. For all $f \in L^{1,\alpha}$, the canonical Fourier Bessel transform $\mathcal{F}_\alpha^m(f)$ belongs to $C_{e,0}(\mathbb{R})$ and verifies

$$(1.7) \quad \|\mathcal{F}_\alpha^m(f)\|_{\infty,\alpha} \leq c_\alpha |b|^{-(\alpha+1)} \|f\|_{1,\alpha}.$$

2. DONOHO-STARK'S UNCERTAINTY PRINCIPLE FOR THE CANONICAL FOURIER BESSEL TRANSFORM

In this section, based on the techniques of Donoho-Stark [9], we will show uncertainty principle of concentration-type the canonical Fourier Bessel transform.

In the following, we consider a pair of orthogonal projections on $L^{2,\alpha}$. The first is the time-limiting operator defined

$$(2.1) \quad P_S f = \chi_S f,$$

and the second is the frequency-limiting operator defined by

$$(2.2) \quad \mathcal{F}_\alpha^m(Q_\Sigma f) = \chi_\Sigma \mathcal{F}_\alpha^m(f),$$

where S and Σ are two measurable subsets of \mathbb{R}_+ and χ_S and χ_Σ denote the characteristic functions of S and Σ .

Definition 2.1. Let $0 < \varepsilon_S, \varepsilon_\Sigma < 1$ and let $f \in L^{2,\alpha}$ be a nonzero function.

(1) We say that f is ε_S -concentrated on S if

$$(2.3) \quad \|P_{S^c} f\|_{2,\alpha} \leq \varepsilon_S \|f\|_{2,\alpha}.$$

(2) We say that f is ε_Σ -concentrated on Σ for the canonical Fourier Bessel transform if

$$(2.4) \quad \|Q_{\Sigma^c} f\|_{2,\alpha} \leq \varepsilon_\Sigma \|f\|_{2,\alpha}.$$

P_S and Q_Σ are projections. Indeed, let $f, g \in L^{2,\alpha}$. By relation (1.4), we have

$$\begin{aligned} \langle P_S^2 f, g \rangle &= \langle P_S f, P_S g \rangle = \langle \mathcal{F}_\alpha^m(P_S f), \mathcal{F}_\alpha^m(P_S g) \rangle \\ &= \int_0^{+\infty} \mathcal{F}_\alpha^m(P_S f)(y) \overline{\mathcal{F}_\alpha^m(P_S g)(y)} y^{2\alpha+1} dy \\ &= \int_S \mathcal{F}_\alpha^m(f)(y) \overline{\mathcal{F}_\alpha^m(g)(y)} y^{2\alpha+1} dy \\ &= \int_0^{+\infty} \mathcal{F}_\alpha^m(P_S f)(y) \overline{\mathcal{F}_\alpha^m(g)(y)} y^{2\alpha+1} dy \\ &= \langle P_S f, g \rangle. \end{aligned}$$

Thus, $P_S^2 = P_S$ and hence P_S is a projection.

By the same way,

$$\begin{aligned} \langle Q_\Sigma^2 f, g \rangle &= \langle Q_\Sigma f, Q_\Sigma g \rangle = \langle \mathcal{F}_\alpha^m(Q_\Sigma f), \mathcal{F}_\alpha^m(Q_\Sigma g) \rangle \\ &= \int_0^{+\infty} \mathcal{F}_\alpha^m(Q_\Sigma f)(y) \overline{\mathcal{F}_\alpha^m(Q_\Sigma g)(y)} y^{2\alpha+1} dy \\ &= \int_\Sigma \mathcal{F}_\alpha^m(f)(y) \overline{\mathcal{F}_\alpha^m(g)(y)} y^{2\alpha+1} dy \\ &= \int_0^{+\infty} \mathcal{F}_\alpha^m(Q_\Sigma f)(y) \overline{\mathcal{F}_\alpha^m(g)(y)} y^{2\alpha+1} dy \\ &= \langle Q_\Sigma f, g \rangle. \end{aligned}$$

Thus, $Q_\Sigma^2 = Q_\Sigma f$ and hence $Q_\Sigma f$ is a projection.

For all $f \in L^{2,\alpha}$, given the kernel N which satisfies the following two conditions: $f(\cdot)N(\cdot, y) \in L^{1,\alpha}$ for almost every $y \in \mathbb{R}_+$ and if

$$\mathcal{M}f(x) = \int_0^{+\infty} f(y)N(x, y)y^{2\alpha+1}dy,$$

then $\mathcal{M}f \in L^{2,\alpha}$. Then we define the norm of \mathcal{M} to be

$$\|\mathcal{M}\| = \sup_{f \in L^{2,\alpha}} \frac{\|\mathcal{M}f\|_{2,\alpha}}{\|f\|_{2,\alpha}}, \quad f \neq 0,$$

and the Hilbert-Schmidt norm of \mathcal{M} is given by

$$\|\mathcal{M}\|_{HS} = \left(\int_0^{+\infty} \int_0^{+\infty} |N(x, y)|^2 x^{2\alpha+1} y^{2\alpha+1} dx dy \right)^{\frac{1}{2}}.$$

It is clear that $\|P_S\| = \|Q_\Sigma\| = 1$ (see [9]). If $|\Sigma| < +\infty$, where Σ is a set of finite measure of \mathbb{R}_+ , we have by [27]

$$|\Sigma| = \int_\Sigma x^{2\alpha+1} dx.$$

Lemma 2.1. *If S and Σ are two measurable sets of \mathbb{R}_+ such that $|S| < +\infty$ and $|\Sigma| < +\infty$, then*

$$\|P_S Q_\Sigma\|_{HS} = \|Q_\Sigma P_S\|_{HS}.$$

Proof. From relations (1.5), (2.1) and (2.2), we have

$$\begin{aligned} Q_\Sigma P_S(f)(x) &= \frac{c_\alpha}{(-ib)^{\alpha+1}} \int_\Sigma \overline{K_\alpha^m(y, x)} \mathcal{F}_\alpha^m(\chi_S f)(y) y^{2\alpha+1} dy \\ &= \frac{c_\alpha}{(-ib)^{\alpha+1}} \int_\Sigma \overline{K_\alpha^m(y, x)} \left(\frac{c_\alpha}{(ib)^{\alpha+1}} \int_S K_\alpha^m(y, z) f(z) z^{2\alpha+1} dz \right) y^{2\alpha+1} dy \\ &= \frac{c_\alpha^2}{b^{2\alpha+2}} \int_S f(z) \left(\int_\Sigma \overline{K_\alpha^m(y, x)} K_\alpha^m(y, z) y^{2\alpha+1} dy \right) z^{2\alpha+1} dz \\ &= \int_S f(z) k(x, z) z^{2\alpha+1} dz, \end{aligned}$$

where

$$k(x, z) = \frac{c_\alpha^2}{b^{2\alpha+2}} \int_\Sigma \overline{K_\alpha^m(y, x)} K_\alpha^m(y, z) y^{2\alpha+1} dy, \quad z \in S, x \in \mathbb{R}_+.$$

In the same way, we get

$$\begin{aligned} P_S Q_\Sigma(f)(x) &= \chi_S(x) Q_\Sigma(f)(x) \\ &= \chi_S(x) \frac{c_\alpha}{(-ib)^{\alpha+1}} \int_\Sigma \overline{K_\alpha^m(y, x)} \mathcal{F}_\alpha^m(f)(y) y^{2\alpha+1} dy \\ &= \chi_S(x) \frac{c_\alpha^2}{b^{2\alpha+2}} \int_\Sigma \overline{K_\alpha^m(y, x)} \left(\int_0^{+\infty} K_\alpha^m(y, z) f(z) z^{2\alpha+1} dz \right) y^{2\alpha+1} dy \\ &= \chi_S(x) \frac{c_\alpha^2}{b^{2\alpha+2}} \int_0^{+\infty} f(z) \left(\int_\Sigma \overline{K_\alpha^m(y, x)} K_\alpha^m(y, z) y^{2\alpha+1} dy \right) z^{2\alpha+1} dz \\ &= \chi_S(x) \int_0^{+\infty} f(z) k(x, z) z^{2\alpha+1} dz. \end{aligned}$$

Then, from the above results we can easily obtain that

$$\|Q_\Sigma P_S\|_{HS} = \left(\int_S \int_0^{+\infty} |k(x, z)|^2 x^{2\alpha+1} z^{2\alpha+1} dx dz \right)^{\frac{1}{2}}$$

and

$$\|P_S Q_\Sigma\|_{HS} = \left(\int_0^{+\infty} \int_S |k(x, z)|^2 x^{2\alpha+1} z^{2\alpha+1} dx dz \right)^{\frac{1}{2}},$$

which yields the desired result. □

Using Cauchy-Schwarz inequality, we can easily obtain that

$$(2.5) \quad \|P_S Q_\Sigma\| \leq \|P_S Q_\Sigma\|_{HS}.$$

Lemma 2.2. *If S and Σ are two measurable subsets of \mathbb{R}_+ such that $|S| < +\infty$ and $|\Sigma| < +\infty$, then*

$$\|P_S Q_\Sigma\| \leq \frac{c_\alpha}{|b|^{\alpha+1}} \sqrt{|S||\Sigma|},$$

where c_α is the constant given by relation (1.1).

Proof. For $x \in S$, let $g_x(t) = k(x, t)$. Note that

$$\mathcal{F}_\alpha^m(g_x)(y) = \frac{c_\alpha}{(ib)^{\alpha+1}} \chi_\Sigma(y) K_\alpha^m(x, y).$$

By relations (1.3) and (1.2), we have

$$\begin{aligned} \int_0^{+\infty} |g_x(t)|^2 t^{2\alpha+1} dt &= \int_0^{+\infty} |\mathcal{F}_\alpha^m(g_x)(y)|^2 y^{2\alpha+1} dy \\ &= \frac{c_\alpha^2}{|b|^{2\alpha+2}} \int_\Sigma |K_\alpha^m(x, y)|^2 y^{2\alpha+1} dy \\ &\leq \frac{c_\alpha^2}{|b|^{2\alpha+2}} |\Sigma|. \end{aligned}$$

Hence,

$$\int_0^{+\infty} \int_0^{+\infty} |k(x, t)|^2 x^{2\alpha+1} t^{2\alpha+1} dx dt \leq \frac{c_\alpha^2}{|b|^{2\alpha+2}} |\Sigma| \int_S x^{2\alpha+1} dx = \frac{c_\alpha^2}{|b|^{2\alpha+2}} |\Sigma| |S|.$$

Therefore,

$$\|P_S Q_\Sigma\|_{HS}^2 \leq \frac{c_\alpha^2}{|b|^{2\alpha+2}} |\Sigma| |S|.$$

And the proof is complete by (2.5). \square

Proposition 2.1. *Let S and Σ be two measurable subsets of \mathbb{R}_+ and assume that $\varepsilon_S + \varepsilon_\Sigma < 1$, f is ε_S -concentrated on S and \mathcal{F}_α^m is ε_Σ -concentrated on Σ , with $\|f\|_{2,\alpha} = 1$. Then*

$$\frac{c_\alpha^2}{|b|^{2\alpha+2}} |\Sigma| |S| \geq (1 - \varepsilon_S - \varepsilon_\Sigma)^2.$$

Proof. Assume that $0 < |S|, |\Sigma| < +\infty$. As $\|Q_\Sigma\| = 1$, it follows that

$$\begin{aligned} \|f - Q_\Sigma P_S(f)\|_{2,\alpha} &\leq \|f - Q_\Sigma(f)\|_{2,\alpha} + \|Q_\Sigma(f) - Q_\Sigma P_S(f)\|_{2,\alpha} \\ &\leq \varepsilon_\Sigma + \|Q_\Sigma\| \|f - P_S(f)\|_{2,\alpha} \\ &\leq \varepsilon_\Sigma + \varepsilon_S. \end{aligned}$$

The triangle inequality gives

$$\|Q_\Sigma P_S(f)\|_{2,\alpha} \geq \|f\|_{2,\alpha} - \|f - Q_\Sigma P_S(f)\|_{2,\alpha} \geq 1 - \varepsilon_\Sigma - \varepsilon_S.$$

Hence,

$$\|Q_\Sigma P_S\| \geq 1 - \varepsilon_\Sigma - \varepsilon_S.$$

Then from lemmas 2.1 and 2.2, we get the desired result. \square

Theorem 2.1 (Donoho-Stark uncertainty principle-type). *Let $f \in L^{2,\alpha}$ and S, Σ be two measurable subsets of \mathbb{R}_+ such that $|S| |\Sigma| < \frac{|b|^{2\alpha+2}}{c_\alpha^2}$ and let $\varepsilon_S, \varepsilon_\Sigma > 0$ such that $\varepsilon_S^2 + \varepsilon_\Sigma^2 < 1$. If f is ε_S -concentrated on S and ε_Σ -concentrated on Σ for the canonical Fourier Bessel transform, then*

$$\frac{c_\alpha^2}{|b|^{2\alpha+2}} |S| |\Sigma| \geq \left(1 - \sqrt{\varepsilon_S^2 + \varepsilon_\Sigma^2}\right)^2.$$

Proof. Since $I = P_S + P_{S^c} = P_S Q_\Sigma + P_S Q_{\Sigma^c} + P_{S^c}$, then, using the orthogonality of P_S and P_{S^c} , we have

$$\begin{aligned} \|f - P_S Q_\Sigma(f)\|_{2,\alpha}^2 &= \|P_S Q_{\Sigma^c}(f) + P_{S^c}(f)\|_{2,\alpha}^2 \\ &= \|P_S Q_{\Sigma^c}(f)\|_{2,\alpha}^2 + \|P_{S^c}(f)\|_{2,\alpha}^2 \\ &\leq \|P_S\|^2 \|Q_{\Sigma^c}(f)\|_{2,\alpha}^2 + \|P_{S^c}(f)\|_{2,\alpha}^2. \end{aligned}$$

From (2.1), we have

$$(2.6) \quad \|P_S\| \leq 1.$$

Since P_S is a projection on $L^{2,\alpha}$, then

$$(2.7) \quad \|P_S\| = \|P_S \circ P_S\| \leq \|P_S\|^2.$$

By (2.6) and (2.7), we deduce that $\|P_S\| = 1$. Thus,

$$(2.8) \quad \|f - P_S Q_\Sigma(f)\|_{2,\alpha} \leq \sqrt{\|Q_{\Sigma^c}(f)\|_{2,\alpha}^2 + \|P_{S^c}(f)\|_{2,\alpha}^2}.$$

On the other hand,

$$\|f - P_S Q_\Sigma(f)\|_{2,\alpha} \geq \|f\|_{2,\alpha} - \|P_S Q_\Sigma(f)\|_{2,\alpha} \geq (1 - \|P_S Q_\Sigma\|)\|f\|_{2,\alpha}.$$

Then, by (2.8), we have

$$(1 - \|P_S Q_\Sigma\|)\|f\|_{2,\alpha} \leq \sqrt{\|Q_{\Sigma^c}(f)\|_{2,\alpha}^2 + \|P_{S^c}(f)\|_{2,\alpha}^2}.$$

Since $\frac{c_\alpha}{|b|^{\alpha+1}} \sqrt{|S||\Sigma|} < 1$, it follows from Lemma 2.2 that

$$(2.9) \quad \|f\|_{2,\alpha}^2 \leq \left(1 - \frac{c_\alpha}{|b|^{\alpha+1}} \sqrt{|S||\Sigma|}\right)^{-2} (\|Q_{\Sigma^c}(f)\|_{2,\alpha}^2 + \|P_{S^c}(f)\|_{2,\alpha}^2).$$

Now, by relations (2.3) and (2.4), we get

$$(2.10) \quad \|Q_{\Sigma^c}(f)\|_{2,\alpha}^2 + \|P_{S^c}(f)\|_{2,\alpha}^2 \leq (\varepsilon_S^2 + \varepsilon_\Sigma^2)\|f\|_{2,\alpha}^2.$$

By combining relations (2.9) and (2.10), we obtain the desired result. □

3. $L^{p,\alpha}$ -UNCERTAINTY PRINCIPLES FOR THE CANONICAL FOURIER BESSEL TRANSFORM

In this section, building on the techniques of Donoho and Stark [9] and Soltani [32], we show a quantitative uncertainty inequality about the essential supports of a nonzero function $f \in L^{p,\alpha}$, $1 \leq p \leq 2$ and its canonical Fourier Bessel transform.

Proposition 3.1. *Let $f \in L^{1,\alpha} \cap L^{p,\alpha}$, $1 < p \leq 2$. Then*

$$\|\mathcal{F}_\alpha^m(f)\|_{q,\alpha} \leq \frac{c_\alpha}{|b|^{\alpha+1}} |\text{supp } \mathcal{F}_\alpha^m(f)|^{\frac{1}{q}} |\text{supp } f|^{\frac{1}{q}} \|f\|_{p,\alpha},$$

with $q = \frac{p}{p-1}$.

Proof. Let $f \in L^{1,\alpha} \cap L^{p,\alpha}$, $1 \leq p \leq 2$. Then by Hölder's inequality and (1.7), we get

$$\begin{aligned} \|\mathcal{F}_\alpha^m(f)\|_{q,\alpha} &\leq |\text{supp } \mathcal{F}_\alpha^m(f)|^{\frac{1}{q}} \|\mathcal{F}_\alpha^m(f)\|_{\infty,\alpha} \\ &\leq \frac{c_\alpha}{|b|^{\alpha+1}} |\text{supp } \mathcal{F}_\alpha^m(f)|^{\frac{1}{q}} \|f\|_{1,\alpha} \\ &\leq \frac{c_\alpha}{|b|^{\alpha+1}} |\text{supp } \mathcal{F}_\alpha^m(f)|^{\frac{1}{q}} |\text{supp } f|^{\frac{1}{q}} \|f\|_{p,\alpha}, \end{aligned}$$

which gives the desired result. □

Proposition 3.2. *Let $f \in L^{2,\alpha} \cap L^{p,\alpha}$, $1 < p < 2$. Then*

$$1 \leq |b|^{(\alpha+1)(\frac{2}{q}-1)} \left(\frac{(c_\alpha p)^{\frac{1}{p}}}{(c_\alpha q)^{\frac{1}{q}}} \right)^{\alpha+1} |\text{supp } \mathcal{F}_\alpha^m(f)|^{\frac{q-2}{2q}} |\text{supp } f|^{\frac{2-p}{2p}},$$

with $q = \frac{p}{p-1}$.

Proof. Let $f \in L^{2,\alpha} \cap L^{p,\alpha}$, $1 \leq p \leq 2$. Then by Hölder's inequality and (1.6), we get

$$\begin{aligned} \|\mathcal{F}_\alpha^m(f)\|_{q,\alpha} &\leq |\text{supp } \mathcal{F}_\alpha^m(f)|^{\frac{q-2}{2q}} \|\mathcal{F}_\alpha^m(f)\|_{q,\alpha} \\ &\leq |b|^{(\alpha+1)(\frac{2}{q}-1)} \left(\frac{(c_\alpha p)^{\frac{1}{p}}}{(c_\alpha q)^{\frac{1}{q}}} \right)^{\alpha+1} |\text{supp } \mathcal{F}_\alpha^m(f)|^{\frac{q-2}{2q}} \|f\|_{p,\alpha} \\ &\leq |b|^{(\alpha+1)(\frac{2}{q}-1)} \left(\frac{(c_\alpha p)^{\frac{1}{p}}}{(c_\alpha q)^{\frac{1}{q}}} \right)^{\alpha+1} |\text{supp } \mathcal{F}_\alpha^m(f)|^{\frac{q-2}{2q}} |\text{supp } f|^{\frac{2-p}{2p}} \|f\|_{2,\alpha}. \end{aligned}$$

Relation (1.3) completes the proof. \square

Definition 3.1. Let $0 < \varepsilon_S, \varepsilon_\Sigma < 1$.

(1) We say that a function $f \in L^{p,\alpha}$, $1 \leq p \leq 2$ is ε_S -concentrated to S in $L^{p,\alpha}$ -norm if and only if

$$(3.1) \quad \|f - P_S f\|_{p,\alpha} \leq \varepsilon_S \|f\|_{p,\alpha}.$$

(2) Let $f \in L^{p,\alpha}$, $1 \leq p \leq 2$. We say that $\mathcal{F}_\alpha^m(f)$ is ε_Σ -concentrated on Σ in $L^{q,\alpha}$ -norm, $q = \frac{p}{p-1}$ if and only if

$$(3.2) \quad \|\mathcal{F}_\alpha^m(f) - \mathcal{F}_\alpha^m(Q_\Sigma f)\|_{q,\alpha} \leq \varepsilon_\Sigma \|\mathcal{F}_\alpha^m(f)\|_{q,\alpha}.$$

Lemma 3.1. *Let $f \in L^{p,\alpha}$, $1 < p \leq 2$. Then*

$$\|\mathcal{F}_\alpha^m(Q_\Sigma f)\|_{q,\alpha} \leq |b|^{(\alpha+1)(\frac{2}{q}-1)} \left(\frac{(c_\alpha p)^{\frac{1}{p}}}{(c_\alpha q)^{\frac{1}{q}}} \right)^{\alpha+1} \|f\|_{p,\alpha},$$

with $q = \frac{p}{p-1}$.

Proof. Let $f \in L^{p,\alpha}$, $1 < p \leq 2$ and $q = \frac{p}{p-1}$. From relations (1.6) and (2.2), we get

$$\begin{aligned} \|\mathcal{F}_\alpha^m(Q_\Sigma f)\|_{q,\alpha} &= \left(\int_\Sigma |\mathcal{F}_\alpha^m(f)(x)|^q x^{2\alpha+1} dx \right)^{\frac{1}{q}} \\ &\leq \|\mathcal{F}_\alpha^m(f)\|_{q,\alpha} \\ &\leq |b|^{(\alpha+1)(\frac{2}{q}-1)} \left(\frac{(c_\alpha p)^{\frac{1}{p}}}{(c_\alpha q)^{\frac{1}{q}}} \right)^{\alpha+1} \|f\|_{p,\alpha}, \end{aligned}$$

which yields the desired result. \square

Lemma 3.2. *Let S and Σ be two measurable subsets of \mathbb{R}_+ and $f \in L^{p,\alpha}$, $1 < p \leq 2$, $q = \frac{p}{p-1}$. Then*

$$\|\mathcal{F}_\alpha^m(Q_\Sigma P_S f)\|_{q,\alpha} \leq \frac{c_\alpha}{|b|^{\alpha+1}} |S|^{\frac{1}{q}} |\Sigma|^{\frac{1}{q}} \|f\|_{p,\alpha}.$$

Proof. Assume that $|S| < +\infty$ and $|\Sigma| < +\infty$. From relation (2.2), we have

$$(3.3) \quad \|\mathcal{F}_\alpha^m(Q_\Sigma P_S f)\|_{q,\alpha} = \left(\int_\Sigma |\mathcal{F}_\alpha^m(\chi_S f)(x)|^q x^{2\alpha+1} dx \right)^{\frac{1}{q}}.$$

By (1.2) and Hölder’s inequality it follows that

$$\begin{aligned} |\mathcal{F}_\alpha^m(\chi_S f)(x)| &\leq \frac{c_\alpha}{|b|^{\alpha+1}} \left(\int_S |f(y)|^p y^{2\alpha+1} dy \right)^{\frac{1}{p}} \left(\int_S |K_\alpha^m(x, y)|^q y^{2\alpha+1} dy \right)^{\frac{1}{q}} \\ &\leq \frac{c_\alpha}{|b|^{\alpha+1}} |S|^{\frac{1}{q}} \|f\|_{p,\alpha}. \end{aligned}$$

Then from (3.3), we obtain the desired result. □

Theorem 3.1. *Let S and Σ be two measurable subsets of \mathbb{R}_+ and $f \in L^{p,\alpha}$, $1 < p \leq 2$, $q = \frac{p}{p-1}$. If f is ε_S -concentration to S in $L^{p,\alpha}$ -norm and $\mathcal{F}_\alpha^m(f)$ is ε_Σ -concentration to Σ in $L^{q,\alpha}$ -norm, then*

$$\|\mathcal{F}_\alpha^m(f)\|_{q,\alpha} \leq \frac{1}{1 - \varepsilon_\Sigma} \left(\frac{c_\alpha}{|b|^{\alpha+1}} |S|^{\frac{1}{q}} |\Sigma|^{\frac{1}{q}} + \varepsilon_S |b|^{(\alpha+1)(\frac{2}{q}-1)} \left(\frac{(c_\alpha p)^{\frac{1}{p}}}{(c_\alpha q)^{\frac{1}{q}}} \right)^{\alpha+1} \right) \|f\|_{p,\alpha}.$$

Proof. Assume that $|S| < +\infty$ and $|\Sigma| < +\infty$. From the triangle inequality, relations (1.6), (3.1), (3.2) and Lemma 3.2, we get

$$\begin{aligned} \|\mathcal{F}_\alpha^m(f)\|_{q,\alpha} &\leq \|\mathcal{F}_\alpha^m(Q_\Sigma P_S f)\|_{q,\alpha} + \|\mathcal{F}_\alpha^m(f) - \mathcal{F}_\alpha^m(Q_\Sigma P_S f)\|_{q,\alpha} \\ &\leq \|\mathcal{F}_\alpha^m(Q_\Sigma P_S f)\|_{q,\alpha} + \|\mathcal{F}_\alpha^m(f) - \mathcal{F}_\alpha^m(Q_\Sigma f)\|_{q,\alpha} \\ &\quad + \|\mathcal{F}_\alpha^m(Q_\Sigma f) - \mathcal{F}_\alpha^m(Q_\Sigma P_S f)\|_{q,\alpha} \\ &\leq \frac{c_\alpha}{|b|^{\alpha+1}} |S|^{\frac{1}{q}} |\Sigma|^{\frac{1}{q}} \|f\|_{p,\alpha} + \varepsilon_\Sigma \|\mathcal{F}_\alpha^m(f)\|_{q,\alpha} \\ &\quad + |b|^{(\alpha+1)(\frac{2}{q}-1)} \left(\frac{(c_\alpha p)^{\frac{1}{p}}}{(c_\alpha q)^{\frac{1}{q}}} \right)^{\alpha+1} \|f - P_S f\|_{p,\alpha} \\ &\leq \left(\frac{c_\alpha}{|b|^{\alpha+1}} |S|^{\frac{1}{q}} |\Sigma|^{\frac{1}{q}} + \varepsilon_S |b|^{(\alpha+1)(\frac{2}{q}-1)} \left(\frac{(c_\alpha p)^{\frac{1}{p}}}{(c_\alpha q)^{\frac{1}{q}}} \right)^{\alpha+1} \right) \|f\|_{p,\alpha} \\ &\quad + \varepsilon_\Sigma \|\mathcal{F}_\alpha^m(f)\|_{q,\alpha}, \end{aligned}$$

which gives the desired result. □

Theorem 3.2 (Donoho-Stark’s uncertainty principle-type). *Let S and Σ be two measurable subsets of \mathbb{R}_+ and $f \in L^{p_1,\alpha} \cap L^{p_2,\alpha}$, $1 < p_1 < p_2 \leq 2$. If f is ε_S -concentration to S in $L^{p_1,\alpha}$ -norm and $\mathcal{F}_\alpha^m(f)$ is ε_Σ -concentration to Σ in $L^{q_2,\alpha}$ -norm,*

$q_2 = \frac{p_2}{p_2-1}$, then

$$\|\mathcal{F}_\alpha^m(f)\|_{q_2,\alpha} \leq \frac{|S|^{\frac{p_2-p_1}{p_1 p_2}} |\Sigma|^{\frac{q_1-q_2}{q_1 q_2}}}{(1-\varepsilon_\Sigma)(1-\varepsilon_S)} |b|^{(\alpha+1)(\frac{2}{q_1}-1)} \left(\frac{(c_\alpha p_1)^{\frac{1}{p_1}}}{(c_\alpha q_1)^{\frac{1}{q_1}}} \right)^{\alpha+1} \|f\|_{p_2,\alpha},$$

where $q_1 = \frac{p_1}{p_1-1}$.

Proof. Assume that $|S| < +\infty$ and $|\Sigma| < +\infty$. Let $f \in L^{p_1,\alpha} \cap L^{p_2,\alpha}$, $1 < p_1 < p_2 \leq 2$. Since $\mathcal{F}_\alpha^m(f)$ is ε_Σ -concentration to Σ in $L^{q_2,\alpha}$ -norm, then, by Hölder's inequality, we obtain

$$\begin{aligned} \|\mathcal{F}_\alpha^m(f)\|_{q_2,\alpha} &\leq \varepsilon_\Sigma \|\mathcal{F}_\alpha^m(f)\|_{q_2,\alpha} + \|\chi_\Sigma \mathcal{F}_\alpha^m(f)\|_{q_2,\alpha} \\ &\leq \varepsilon_\Sigma \|\mathcal{F}_\alpha^m(f)\|_{q_2,\alpha} + |\Sigma|^{\frac{q_1-q_2}{q_1 q_2}} \|\mathcal{F}_\alpha^m(f)\|_{q_1,\alpha}. \end{aligned}$$

Thus, by (1.6),

$$(3.4) \quad \|\mathcal{F}_\alpha^m(f)\|_{q_2,\alpha} \leq \frac{|\Sigma|^{\frac{q_1-q_2}{q_1 q_2}}}{1-\varepsilon_\Sigma} |b|^{(\alpha+1)(\frac{2}{q_1}-1)} \left(\frac{(c_\alpha p_1)^{\frac{1}{p_1}}}{(c_\alpha q_1)^{\frac{1}{q_1}}} \right)^{\alpha+1} \|f\|_{p_1,\alpha}.$$

On the other hand, since f is ε_S -concentration to S in $L^{p_1,\alpha}$ -norm, then by Hölder's inequality, we deduce that

$$\|f\|_{p_1,\alpha} \leq \varepsilon_S \|f\|_{p_1,\alpha} + \|\chi_S f\|_{p_1,\alpha} \leq \varepsilon_S \|f\|_{p_1,\alpha} + |S|^{\frac{p_2-p_1}{p_1 p_2}} \|f\|_{p_2,\alpha}.$$

Thus,

$$(3.5) \quad \|f\|_{p_1,\alpha} \leq \frac{|S|^{\frac{p_2-p_1}{p_1 p_2}}}{1-\varepsilon_S} \|f\|_{p_2,\alpha}.$$

Combining (3.4) and (3.5), we obtain the result of this theorem. \square

Corollary 3.1. *Let S and Σ be two measurable subsets of \mathbb{R}_+ and $f \in L^{2,\alpha} \cap L^{p,\alpha}$, $1 < p < 2$. If f is ε_S -concentration to S in $L^{p,\alpha}$ -norm and $\mathcal{F}_\alpha^m(f)$ is ε_Σ -concentration to Σ in $L^{2,\alpha}$ -norm, then*

$$(1-\varepsilon_\Sigma)(1-\varepsilon_S) \leq |S|^{\frac{2-p}{2p}} |\Sigma|^{\frac{q-2}{2q}} |b|^{(\alpha+1)(\frac{2}{q}-1)} \left(\frac{(c_\alpha p)^{\frac{1}{p}}}{(c_\alpha q)^{\frac{1}{q}}} \right)^{\alpha+1},$$

where $q = \frac{p}{p-1}$.

Let $B^p(\Sigma)$, $1 \leq p \leq 2$, be the set of functions $g \in L^{p,\alpha}$ that are bandlimited to Σ , i.e., ($g \in B^p(\Sigma)$ implies $Q_\Sigma g = g$).

We say that f is ε_Σ -bandlimited to Σ in $L^{p,\alpha}$ -norm if there is a $g \in B^p(\Sigma)$ with

$$\|f - g\|_{p,\alpha} \leq \varepsilon_\Sigma \|f\|_{p,\alpha}.$$

In the following, we state an $L^{p_1,\alpha} \cap L^{p_2,\alpha}$ bandlimited uncertainty principle of concentration-type.

Theorem 3.3 (Bandlimited principle-type). *Let S and Σ be two measurable subsets of \mathbb{R}_+ and $f \in L^{p_1,\alpha} \cap L^{p_2,\alpha}$, $1 \leq p_1 < p_2 \leq 2$. If f is ε_S -concentration to S in $L^{p_1,\alpha}$ -norm and ε_Σ -bandlimited to Σ in $L^{q_2,\alpha}$ -norm, $q_2 = \frac{p_2}{p_2-1}$, then*

$$\begin{aligned} & \|f\|_{p_1,\alpha} \\ & \leq \frac{|S|^{\frac{p_2-p_1}{p_1 p_2}}}{1 - \varepsilon_S} \left[(1 + \varepsilon_\Sigma) c_\alpha |\Sigma|^{\frac{1}{p_2}} |S|^{\frac{1}{p_2}} |b|^{(\alpha+1)(\frac{2}{q_2}-2)} \left(\frac{(c_\alpha p_2)^{\frac{1}{p_2}}}{(c_\alpha q_2)^{\frac{1}{q_2}}} \right)^{\alpha+1} + \varepsilon_\Sigma \right] \|f\|_{p_2,\alpha}. \end{aligned}$$

Proof. Assume that $|S| < +\infty$ and $|\Sigma| < +\infty$. Let $f \in L^{p_1,\alpha} \cap L^{p_2,\alpha}$, $1 \leq p_1 < p_2 \leq 2$. Since f is ε_S -concentration to S in $L^{p_1,\alpha}$ -norm, then by Hölder’s inequality, we deduce that

$$\|f\|_{p_1,\alpha} \leq \varepsilon_S \|f\|_{p_1,\alpha} + \|P_S f\|_{p_1,\alpha} \leq \varepsilon_S \|f\|_{p_1,\alpha} + |S|^{\frac{p_2-p_1}{p_1 p_2}} \|P_S f\|_{p_2,\alpha}.$$

Thus,

$$(3.6) \quad \|f\|_{p_1,\alpha} \leq \frac{|S|^{\frac{p_2-p_1}{p_1 p_2}}}{1 - \varepsilon_S} \|P_S f\|_{p_2,\alpha}.$$

As f is ε_Σ -bandlimited to Σ in $L^{q_2,\alpha}$ -norm, there is a $g \in B^{p_2}(\Sigma)$ with

$$\|f - g\|_{p_2,\alpha} \leq \varepsilon_\Sigma \|f\|_{p_2,\alpha}.$$

On the other hand, we have

$$\|P_S f\|_{p_2,\alpha} \leq \|P_S g\|_{p_2,\alpha} + \|P_S(f - g)\|_{p_2,\alpha} \leq \|P_S g\|_{p_2,\alpha} + \varepsilon_\Sigma \|f\|_{p_2,\alpha}.$$

But $g \in B^{p_2}(\Sigma)$, from (2.2), $g(x) = \mathcal{F}_\alpha^{m-1}(\chi_\Sigma \mathcal{F}_\alpha^m(g))(x)$ and by (1.6) and Hölder’s inequality, we deduce that

$$\begin{aligned} |g(x)| & \leq \frac{c_\alpha}{|b|^{\alpha+1}} |\Sigma|^{\frac{1}{p_2}} \|\mathcal{F}_\alpha^m(g)\|_{q_2,\alpha} \\ & \leq c_\alpha |\Sigma|^{\frac{1}{p_2}} |b|^{(\alpha+1)(\frac{2}{q_2}-2)} \left(\frac{(c_\alpha p_2)^{\frac{1}{p_2}}}{(c_\alpha q_2)^{\frac{1}{q_2}}} \right)^{\alpha+1} \|g\|_{p_2,\alpha}. \end{aligned}$$

Hence,

$$\begin{aligned} \|P_S g\|_{p_2,\alpha} & = \left(\int_S |g(x)|^{p_2} x^{2\alpha+1} dx \right)^{\frac{1}{p_2}} \\ & \leq c_\alpha |\Sigma|^{\frac{1}{p_2}} |S|^{\frac{1}{p_2}} |b|^{(\alpha+1)(\frac{2}{q_2}-2)} \left(\frac{(c_\alpha p_2)^{\frac{1}{p_2}}}{(c_\alpha q_2)^{\frac{1}{q_2}}} \right)^{\alpha+1} \|g\|_{p_2,\alpha}. \end{aligned}$$

Then by (3.6) and the fact that $\|g\|_{p_2,\alpha} \leq (1 + \varepsilon_\Sigma) \|f\|_{p_2,\alpha}$, we get

$$\|f\|_{p_1,\alpha} \leq \frac{|S|^{\frac{p_2-p_1}{p_1 p_2}}}{1 - \varepsilon_S} \left[(1 + \varepsilon_\Sigma) c_\alpha |\Sigma|^{\frac{1}{p_2}} |S|^{\frac{1}{p_2}} |b|^{(\alpha+1)(\frac{2}{q_2}-2)} \left(\frac{(c_\alpha p_2)^{\frac{1}{p_2}}}{(c_\alpha q_2)^{\frac{1}{q_2}}} \right)^{\alpha+1} + \varepsilon_\Sigma \right] \|f\|_{p_2,\alpha}.$$

This completes the desired result. □

Corollary 3.2. *Let S and Σ be two measurable subsets of \mathbb{R}_+ and $f \in L^{p,\alpha}$, $1 < p \leq 2$. If f is ε_S -concentration to S and ε_Σ -bandlimited to Σ in $L^{p,\alpha}$ -norm, then*

$$\frac{1 - \varepsilon_S - \varepsilon_\Sigma}{1 + \varepsilon_\Sigma} \leq c_\alpha |\Sigma|^{\frac{1}{p}} |S|^{\frac{1}{p}} |b|^{(\alpha+1)(\frac{2}{q}-2)} \left(\frac{(c_\alpha p)^{\frac{1}{p}}}{(c_\alpha q)^{\frac{1}{q}}} \right)^{\alpha+1}.$$

Theorem 3.4 (Matolcsi-Szűcs-type inequality). *Let $f \in L^{p_1,\alpha} \cap L^{p_2,\alpha}$, $1 < p_1 \leq p_2 \leq 2$. Then*

$$\|\mathcal{F}_\alpha^m(f)\|_{q_2,\alpha} \leq |b|^{(\alpha+1)(\frac{2}{q_1}-1)} \left(\frac{(c_\alpha p_1)^{\frac{1}{p_1}}}{(c_\alpha q_1)^{\frac{1}{q_1}}} \right)^{\alpha+1} |\text{supp } \mathcal{F}_\alpha^m(f)|^{\frac{q_1-q_2}{q_1 q_2}} |\text{supp } f|^{\frac{p_2-p_1}{p_1 p_2}} \|f\|_{p_2,\alpha},$$

where $q_1 = \frac{p_1}{p_1-1}$ and $q_2 = \frac{p_2}{p_2-1}$.

Proof. Let $f \in L^{p_1,\alpha} \cap L^{p_2,\alpha}$, $1 < p_1 \leq p_2 \leq 2$, $q_1 = \frac{p_1}{p_1-1}$ and $q_2 = \frac{p_2}{p_2-1}$. Then, by relation (1.6) and Hölder’s inequality, we obtain

$$\begin{aligned} & \|\mathcal{F}_\alpha^m(f)\|_{q_2,\alpha} \\ & \leq |\text{supp } \mathcal{F}_\alpha^m(f)|^{\frac{q_1-q_2}{q_1 q_2}} \|\mathcal{F}_\alpha^m(f)\|_{q_1,\alpha} \\ & \leq |b|^{(\alpha+1)(\frac{2}{q_1}-1)} \left(\frac{(c_\alpha p_1)^{\frac{1}{p_1}}}{(c_\alpha q_1)^{\frac{1}{q_1}}} \right)^{\alpha+1} |\text{supp } \mathcal{F}_\alpha^m(f)|^{\frac{q_1-q_2}{q_1 q_2}} \|f\|_{p_1,\alpha} \\ & \leq |b|^{(\alpha+1)(\frac{2}{q_1}-1)} \left(\frac{(c_\alpha p_1)^{\frac{1}{p_1}}}{(c_\alpha q_1)^{\frac{1}{q_1}}} \right)^{\alpha+1} |\text{supp } \mathcal{F}_\alpha^m(f)|^{\frac{q_1-q_2}{q_1 q_2}} |\text{supp } f|^{\frac{p_2-p_1}{p_1 p_2}} \|f\|_{p_2,\alpha}, \end{aligned}$$

which yields the desired result. □

Corollary 3.3. *Let $f \in L^{2,\alpha} \cap L^{p,\alpha}$, $1 < p < 2$ and $q = \frac{p}{p-1}$. Then*

$$\left(\frac{(c_\alpha p)^{\frac{1}{p}}}{(c_\alpha q)^{\frac{1}{q}}} \right)^{-(\alpha+1)} |b|^{(\alpha+1)(1-\frac{2}{q})} \leq |\text{supp } f|^{\frac{2-p}{2p}} |\text{supp } \mathcal{F}_\alpha^m(f)|^{\frac{q-2}{2q}}.$$

REFERENCES

- [1] S. Abe and J. T. Sheridan, *Optical operations on wave functions as the Abelian subgroups of the special affine Fourier transformation*, Optics Letters **19** (1994), 1801–1803.
- [2] M. Bahri and R. Ashino, *A variation on uncertainty principle and logarithmic uncertainty principle for continuous quaternion wavelet transforms*, Abstr. Appl. Anal. **11** (2017), Article ID 3795120. <https://doi.org/10.1155/2017/3795120>
- [3] B. Barshan, M. Kutay and H. Ozaktas, *Optimal filtering with linear canonical transformations*, Optics Communications **135** (1997), 32–36.
- [4] L. M. Bernardo, *ABCD matrix formalism of fractional Fourier optics*, Optical Engineering (Bellingham) **35** (1996), 732–740.
- [5] S. A. J. Collins, *Lens-system diffraction integral written in terms of matrix optics*, Journal of the Optical Society of America **60**(9) (1970), 1168–1177.
- [6] M. Cowling and J. F. Price, *Bandwidth versus time concentration: the Heisenberg-Pauli-Weyl inequality*, SIAM J. Math. Anal. **15** (1984), 151–165.

- [7] M. F. E. De Jeu, *The Dunkl transform*, Inventory Math. **113** (1993), 147–162.
- [8] L. Dhaouadi, J. Sahbani and A. Fitouhi, *Harmonic analysis associated to the canonical Fourier Bessel transform*, Integral Transforms Spec. Funct. (2019), 290–315. <https://doi.org/10.1080/10652469.2020.1823977>
- [9] D. L. Donoho and P. B. Strak, *Uncertainty principles and signal recovery*, SIAM J. Appl. Math. **49** (3) (1989), 906–931.
- [10] C. F. Dunkl, *Hankel transforms associated to finite reflection groups*, Contemp. Math. **138**(1) (1992), 128–138.
- [11] S. Ghazouani, E. A. Soltani and A. Fitouhi, *A unified class of integral transforms related to the Dunkl transform*, J. Math. Anal. Appl. **449**(2) (2017), 1797–1849.
- [12] S. Ghobber, *Variations on uncertainty principles for integral operators*, Appl. Anal. **93**(5) (2014), 1057–1072.
- [13] K. Hleili and S. Omri, *An $L^p - L^q$ version of Miyachi's theorem for the Riemann-Liouville operator*, Indian Journal of Pure and Applied Mathematics **46**(2) (2015), 121–138.
- [14] K. Hleili, *Uncertainty principles for spherical mean L^2 -multiplier operators*, J. Pseudo-Differ. Oper. Appl. **9** (2018), 573–587.
- [15] K. Hleili, *Continuous wavelet transform and uncertainty principle related to the Weinstein operator*, Integral Transforms Spec. Funct. **29**(4) (2018), 252–268.
- [16] K. Hleili, *A dispersion inequality and accumulated Spectrograms in the Weinstein Setting*, Bull. Math. Anal. Appl. **12**(1) (2020), 51–70.
- [17] K. Hleili, *Some Results for the windowed Fourier transform related to the spherical mean operator*, Acta Math. Vietnam. **46**(1) (2021), 179–201.
- [18] K. Hleili, *A variety of uncertainty principles for the Hankel-Stockwell transform*, Open Journal of Mathematical Analysis **5**(1) (2021), 22–34.
- [19] K. Hleili, *A variation on uncertainty principles for quaternion linear canonical transform*, Advances in Applied Clifford Algebras **31**(3) (2021), 1–13.
- [20] K. Hleili, *Windowed linear canonical transform and its applications to the time-frequency analysis*, J. Pseudo-Differ. Oper. Appl. **13**(2) (2022), 1–26.
- [21] K. Hleili, *L^p uncertainty principles for the Windowed Spherical mean transform*, Mem. Differ. Equ. Math. Phys. **85** (2022), 75–90.
- [22] D. F. V. James and G. S. Agarwal, *The generalized Fresnel transform and its application to optics*, Optics Communications **126** (1996), 207–212.
- [23] F. H. Kerr, *A fractional power theory for Hankel transforms in $L^2(\mathbb{R}_+)$* , J. Math. Anal. Appl. **158**(1) (1991), 114–123.
- [24] H. J. Landau, *On Szegő's eigenvalue distribution theorem and non-Hermitian kernels*, J. Anal. Math. **28** (1975), 335–357.
- [25] H. J. Landau and H. O. Pollak, *Prolate spheroidal wave functions, Fourier analysis and uncertainty- III: the dimension of the space of essentially time- and band-limited signals*, The Bell System Technical Journal **41** (1962), 1295–1336.
- [26] M. Moshinsky and C. Quesne, *Linear canonical transformations and their unitary representations*, J. Math. Phys. **12**(8) (1971), 1772–1780.
- [27] S. Omri, *Local uncertainty principle for the Hankel transform*, Integral Transforms Spec. Funct. **21**(9) (2010), 703–712.
- [28] H. Ozaktas, Z. Zalevsky and M. Kutay, *The Fractional Fourier Transform with Applications in Optics and Signal Processing*, Wiley, New York, 2001.
- [29] S. Pei and J. Ding, *Eigenfunctions of linear canonical transform*, IEEE Trans. Signal Process. **50** (2002), 11–26.
- [30] J. F. Price, *Inequalities and local uncertainty principles*, J. Math. Phys. **24** (1983), 1711–1714.
- [31] F. Soltani, *An L^p Heisenberg-Pauli-Weyl uncertainty principle for the Dunkl transform*, Konuralp J. Math. **2**(1) (2014), 1–6.

- [32] F. Soltani and J. Ghazwani, *A variation of the L^p uncertainty principles for the Fourier transform*, Proc. Inst. Math. Mech. Natl. Acad. Sci. Azerb. **42**(1) (2016), 10–24.
- [33] G. N. Watson, *A Treatise on the Theory of Bessel Functions*, Cambridge University Press, Cambridge, 1944.
- [34] K. B. Wolf, *Canonical transforms. II, Complex radial transforms*, J. Math. Phys. **15** (1974), 1295–1301.

¹DEPARTMENT OF MATHEMATICS,
NORTHERN BORDERS UNIVERSITY,
KAIROUAN UNIVERSITY,
FACULTY OF SCIENCES OF ARAR, ARAR, SAUDI ARABIA
PREPARATORY INSTITUTE FOR ENGINEERING STUDIES OF KAIROUAN, KAIROUAN, TUNISIA
Email address: khaled.hleili@gmail.com

²DEPARTMENT OF MATHEMATICS,
UNIVERSITY OF TUNIS EL MANAR,
UNIVERSITY OF TABUK,
FACULTY OF SCIENCES OF TUNIS, TUNIS, TUNISIA
FACULTY OF SCIENCES OF TABUK, TABUK, SAUDI ARABIA
Email address: Hleili.manel@outlook.com