

BI-PERIODIC HYPER-FIBONACCI NUMBERS

NASSIMA BELAGGOUN^{1,2} AND HACÈNE BELBACHIR^{1,2}

ABSTRACT. In the present paper, we introduce and study a new generalization of hyper-Fibonacci numbers, called the bi-periodic hyper-Fibonacci numbers. Furthermore, we give a combinatorial interpretation using the weighted tilings approach and prove several identities relating these numbers. Moreover, we derive their generating function and new identities for the classical hyper-Fibonacci numbers.

1. INTRODUCTION

The Fibonacci numbers F_n are defined, as usual, by the recurrence relation

$$F_0 = 0, F_1 = 1 \quad \text{and} \quad F_n = F_{n-1} + F_{n-2}, \quad \text{for } n \geq 2.$$

The hyper-Fibonacci numbers denoted $F_n^{(r)}$, are introduced by Dil and Mezö [10], for $n, r \in \mathbb{N} \cup \{0\}$, as entries of an infinite matrix arranged such that $F_n^{(r)}$ is the entry of the r th row and n th column, satisfying

$$(1.1) \quad F_n^{(0)} = F_n, F_0^{(r)} = 0 \quad \text{and} \quad F_n^{(r)} = F_{n-1}^{(r)} + F_n^{(r-1)}, \quad \text{for } n, r \geq 1.$$

The sum of the first $n + 1$ elements of row $r - 1$ is expressed by $F_n^{(r)}$, i.e.,

$$(1.2) \quad F_n^{(r)} = \sum_{k=0}^n F_k^{(r-1)}.$$

They satisfy many interesting number theoretical and combinatorial properties, see [9]. Belbachir and Belkhir [3] provided a combinatorial interpretation of the hyper-Fibonacci numbers in terms of linear tilings and gave some combinatorial identities.

Key words and phrases. Hyper-Fibonacci numbers, bi-periodic Fibonacci numbers, bi-periodic hyper-Fibonacci numbers, generating function.

2020 *Mathematics Subject Classification.* Primary: 11B39, 05A15. Secondary: 05A19.

DOI 10.46793/KgJMat2504.603B

Received: June 12, 2022.

Accepted: October 08, 2022.

They also defined bivariate hyper-Fibonacci polynomials in [4], as

$$(1.3) \quad F_n^{(r)}(x, y) = xF_{n-1}^{(r)}(x, y) + yF_n^{(r-1)}(x, y), \quad \text{for } n, r \geq 1,$$

with initial conditions $F_n^{(0)}(x, y) = F_n(x, y)$, $F_0^{(r)}(x, y) = 0$, where x, y are real parameters and $F_n(x, y)$ is the n th bivariate Fibonacci polynomial, defined by (see [1, 5])

$$F_0(x, y) = 0, \quad F_1(x, y) = 1 \quad \text{and} \quad F_n(x, y) = xF_{n-1}(x, y) + yF_{n-2}(x, y).$$

The bivariate hyper-Fibonacci polynomials are given by the following explicit formula

$$(1.4) \quad F_{n+1}^{(r)}(x, y) = \sum_{k=r}^{\lfloor n/2 \rfloor + r} \binom{n + 2r - k}{k} x^{n+2r-2k} y^k.$$

The associated generating function is given as follows

$$(1.5) \quad \sum_{n \geq 0} F_n^{(r)}(x, y) z^n = \frac{y^r z}{(1 - xz - yz^2)(1 - xz)^r}.$$

For $y = 1$, we denote $F_n(x, y)$ by $F_n(x)$.

Edson and Yayenie [12] introduced a new generalization for the Fibonacci sequence, called as bi-periodic Fibonacci sequence, that depends on two real parameters a and b , defined for $n \geq 2$, as follows

$$(1.6) \quad q_n = \begin{cases} aq_{n-1} + q_{n-2}, & \text{if } n \text{ is even,} \\ bq_{n-1} + q_{n-2}, & \text{if } n \text{ is odd,} \end{cases}$$

with initial values $q_0 = 0$ and $q_1 = 1$. These sequences are found in the study of continued fraction expansion of the quadratic irrational numbers and combinatorics on words or dynamical system theory [18]. Some well-known sequences, such as the Fibonacci sequence, the Pell sequence and the k -Fibonacci sequence for some positive integer k , are special cases of this sequence. For more results related to this sequence, see [8, 11–18]

The generating function of q_n is given by

$$(1.7) \quad \sum_{n \geq 0} q_n z^n = \frac{z(1 + az - z^2)}{1 - (ab + 2)z^2 + z^4}.$$

Yayenie [18] gave an explicit formula of bi-periodic Fibonacci numbers, as

$$(1.8) \quad q_{n+1} = a^{\xi(n)} \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n - k}{k} (ab)^{\lfloor n/2 \rfloor - k},$$

where $\xi(n) = n - 2\lfloor n/2 \rfloor$, i.e., $\xi(n) = 0$ when n is even and $\xi(n) = 1$ when n is odd.

In this paper, we define a new generalization of hyper-Fibonacci numbers, which we will also call bi-periodic hyper-Fibonacci numbers. We give a combinatorial interpretation of these numbers using a weighted tilings approach and provide several combinatorial proofs of some identities. We also obtain new identities for the classical hyper-Fibonacci numbers. Moreover, by using the generating function of the bivariate

hyper-Fibonacci polynomials, we establish the generating function of the bi-periodic hyper-Fibonacci sequence.

Definition 1.1. For any integers $n, r \geq 1$ and nonzero real numbers a and b , the bi-periodic hyper-Fibonacci numbers, denoted by $q_n^{(r)}$, are defined by

$$(1.9) \quad q_n^{(r)} = \sum_{k=0}^n a^{\xi(k)\xi(n+1)} b^{\xi(k+1)\xi(n)} (ab)^{\lfloor (n-k)/2 \rfloor} q_k^{(r-1)},$$

with initial values $q_0^{(r)} = 0$ and $q_n^{(0)} = q_n$, where q_n is the n th bi-periodic Fibonacci number.

The first few generations are as follows in Table 1.

TABLE 1. Sequence of bi-periodic hyper-Fibonacci numbers in the first few generations

n	0	1	2	3	4	5	6
$q_n^{(0)}$	0	1	a	$ab + 1$	$a^2b + 2a$	$a^2b^2 + 3ab + 1$	$a^3b^2 + 4a^2b + 3a$
$q_n^{(1)}$	0	1	$2a$	$3ab + 1$	$4a^2b + 3a$	$5a^2b^2 + 6ab + 1$	$6a^3b^2 + 10a^2b + 4a$
$q_n^{(2)}$	0	1	$3a$	$6ab + 1$	$10a^2b + 4a$	$15a^2b^2 + 10ab + 1$	$21a^3b^2 + 20a^2b + 5a$
$q_n^{(3)}$	0	1	$4a$	$10ab + 1$	$20a^2b + 5a$	$35a^2b^2 + 15ab + 1$	$56a^3b^2 + 35a^2b + 6a$
$q_n^{(4)}$	0	1	$5a$	$15ab + 1$	$35a^2b + 6a$	$70a^2b^2 + 21ab + 1$	$126a^3b^2 + 56a^2b + 7a$

From the definition, we have the following recurrence relation:

$$(1.10) \quad q_n^{(r)} = \begin{cases} aq_{n-1}^{(r)} + q_n^{(r-1)}, & \text{if } n \text{ is even,} \\ bq_{n-1}^{(r)} + q_n^{(r-1)}, & \text{if } n \text{ is odd.} \end{cases}$$

Note that, for $a = b = 1$, we obtain the classical hyper-Fibonacci sequence (1.1).

2. COMBINATORIAL IDENTITIES

The Fibonacci numbers can be interpreted as the number of ways to tile a board of length n (i.e., an n -board) with cells numbered 1 to n from left to right using only squares and dominoes; see [6, 7]. We expand the results to bi-periodic Fibonacci numbers using weighted tilings. We assign a weight to each square in a tiling based on its position. It is assigned a weight a if it is in an odd position and a weight b if it is in an even position. The weight of a tiling of an n -board is defined as the product of the weights of its individual tiles. The sum of all possible weighted tilings is given by q_{n+1} . Furthermore, the total of all possible weighted tilings of an $(n + 2r)$ -board with at least r dominoes is given by the bi-periodic hyper-Fibonacci numbers $q_{n+1}^{(r)}$, as shown in Theorem 2.1.

For example, Figure 1 shows the tilings and the sum of their weights of a 5-board. We have $q_6^{(0)} = q_6 = a^3b^2 + 4a^2b + 3a$.




FIGURE 1. Tilings of a 5-board

Figure 2 shows the tilings and the sum of their weights of a 6-board with at least 2 dominoes, there are $q_3^{(2)} = 6ab + 1$ dispositions.




FIGURE 2. Tilings of a 6-board with at least 2 dominos

Therefore, we have the following results.

Theorem 2.1. For $n, r \geq 0$, $q_{n+1}^{(r)}$ gives the weight of all tilings of an $(n + 2r)$ -board having at least r dominoes.

Proof. Given $(n + 2r)$ -board. If it ends with a square, then there are $bq_n^{(r)}$ ways to tile the $(n + 2r - 1)$ -board for n even and $aq_n^{(r)}$ for n odd. If it ends with a domino, then there are $q_{n+1}^{(r-1)}$ ways to tile the $(n + 2(r - 1))$ -board. When $n = 0$, there is one way to tile a $2r$ -board with at least r dominoes and there are q_{n+1} ways to tile a n -board with at least 0 dominoes. There is no way to tile an $(n + 2r)$ -board with at least r dominoes for $n < 0$. \square

Let $f(n, k)$ be the number of weighted tilings having n tiles and exactly k dominoes. Then

$$f(n, k) = a^{\xi(n+k)}b^{\xi(n+k+1)}f(n - 1, k) + f(n - 1, k - 1).$$

In fact, if the $(n + k)$ -board ends in a square there are $a^{\xi(n+k)}b^{\xi(n+k+1)}f(n - 1, k)$ ways to tile the board. If it ends with a domino, then there are $f(n - 1, k - 1)$ ways.

Lemma 2.1. The number of weighted tilings having n tiles and exactly k dominoes is

$$a^{\xi(n+k)}\binom{n}{k}(ab)^{\lfloor (n-k)/2 \rfloor}.$$

Proof. Let $g(n, k) = a^{\xi(n+k)} \binom{n}{k} (ab)^{\lfloor (n-k)/2 \rfloor}$. Then

$$a^{\xi(n+k)} \binom{n}{k} (ab)^{\lfloor (n-k)/2 \rfloor} = a^{\xi(n+k)} \left(\binom{n-1}{k} + \binom{n-1}{k-1} \right) (ab)^{\lfloor (n-k)/2 \rfloor}.$$

Using $\lfloor (n-k)/2 \rfloor = \lfloor (n-k-1)/2 \rfloor + \xi(n+k+1)$, we get

$$\begin{aligned} a^{\xi(n+k)} \binom{n}{k} (ab)^{\lfloor (n-k)/2 \rfloor} &= a^{\xi(n+k)} (ab)^{\xi(n+k+1)} \binom{n-1}{k} (ab)^{\lfloor (n-k-1)/2 \rfloor} \\ &\quad + a^{\xi(n+k)} \binom{n-1}{k-1} (ab)^{\lfloor (n-k)/2 \rfloor} \\ &= a^{\xi(n+k)} b^{\xi(n+k+1)} g(n-1, k) + g(n-1, k-1). \end{aligned}$$

Since $g(n, k)$ satisfies the same recurrence of $f(n, k)$ and the same initial conditions, we get result. □

In the following theorems, we establish an explicit formula for the bi-periodic hyper-Fibonacci sequence.

Theorem 2.2. *For $n, r \geq 0$, we have*

$$(2.1) \quad q_{n+1}^{(r)} = a^{\xi(n)} \sum_{k=r}^{\lfloor n/2 \rfloor + r} \binom{n+2r-k}{k} (ab)^{\lfloor n/2 \rfloor + r - k}.$$

Proof. From Theorem 2.1, $q_{n+1}^{(r)}$ counts the number of ways to tile an $(n+2r)$ -board with at least r dominoes. On the other hand, using Lemma 2.1, the possible tilings with exactly k dominoes contains $n+2r-2k$ squares and $n+2r-k$ tiles, have cardinality $a^{\xi(n)} \binom{n+2r-k}{k} (ab)^{\lfloor n/2 \rfloor + r - k}$. Since it contains at least r dominoes, the sum over $k \geq r$ gives the identity. □

Now, we establish a double-summation formula for even-numbered bi-periodic hyper-Fibonacci numbers $q_{2n+2}^{(r)}$.

Theorem 2.3. *For $n, r \geq 0$, we have*

$$(2.2) \quad q_{2n+2}^{(r)} = a \sum_{k=r}^{n+r} \sum_{j=0}^k (ab)^{\xi(n+r-k)} \binom{n+r-j}{k-j} \binom{n+r-k+j}{j} (ab)^{2\lfloor (n+r-k)/2 \rfloor}.$$

Proof. Consider an $(n+2r+1)$ -board. Since the length of the board is odd, there are an odd number of squares such that we have at least one in each tiling. Suppose there are i dominoes to the left of its median square and j dominoes to its right, whose total is at least r dominoes, i.e., $i+j \geq r$. The median square contributes an $a^{\xi(n+r-i-j+1)} b^{\xi(n+r-i-j)}$ to the weight (according to the position of the median square). Such tiling contains $2n+2r-2i-2j+1$ squares, so there are $n+r-i-j$ squares on each side of the median square. The left side gives $n+r-j$ tiles with i dominos. Hence, there are $a^{\xi(n+r-i-j)} \binom{n+r-j}{i} (ab)^{\lfloor (n+r-i-j)/2 \rfloor}$ different ways. Similarly,

we have $a^{\xi(n+r-i-j)} \binom{n+r-i}{j} (ab)^{\lfloor (n+r-i-j)/2 \rfloor}$ different ways to tile the right side. Thus, the possible tilings have cardinality $a(ab)^{\xi(n+r-i-j)} \binom{n+r-i}{j} (ab)^{2\lfloor (n+r-i-j)/2 \rfloor}$. Summing over $i + j \geq r$, we get

$$\begin{aligned} & a \sum_{r \leq i+j \leq n+r} (ab)^{\xi(n+r-i-j)} \binom{n+r-j}{i} \binom{n+r-i}{j} (ab)^{2\lfloor (n+r-i-j)/2 \rfloor} \\ &= a \sum_{k=r}^{n+r} \sum_{i+j=k} (ab)^{\xi(n+r-k)} \binom{n+r-j}{i} \binom{n+r-i}{j} (ab)^{2\lfloor (n+r-k)/2 \rfloor} \\ &= a \sum_{k=r}^{n+r} \sum_{j=0}^k (ab)^{\xi(n+r-k)} \binom{n+r-j}{k-j} \binom{n+r-k+j}{j} (ab)^{2\lfloor (n+r-k)/2 \rfloor}. \quad \square \end{aligned}$$

For $a = b = 1$, we get the following identity.

Corollary 2.1. *For $n, r \geq 0$, the following identity holds*

$$(2.3) \quad F_{2n+2}^{(r)} = \sum_{k=r}^{n+r} \sum_{j=0}^k \binom{n+r-j}{k-j} \binom{n+r-k+j}{j}.$$

From the explicit formulas (1.8) and (2.1), we state the bi-periodic hyper-Fibonacci sequence in terms of the bi-periodic Fibonacci sequence and binomial sum.

Theorem 2.4. *Let $n \geq 0$ and $r \geq 1$ be integers, then we have*

$$(2.4) \quad q_{n+1}^{(r)} = q_{n+1+2r} - a^{\xi(n)} \sum_{k=0}^{r-1} \binom{n+2r-k}{k} (ab)^{\lfloor n/2 \rfloor + r - k}.$$

Note that, if we take $a = b = 1$, we get the following identity, see [3],

$$F_{n+1}^{(r)} = F_{n+1+2r} - \sum_{k=0}^{r-1} \binom{n+2r-k}{k}.$$

Theorem 2.5. *For $n, r \geq 1$, we have*

$$(2.5) \quad q_{n+1}^{(r)} = q_{n-1} + \sum_{k=0}^r a^{\xi(n)} b^{\xi(n+1)} q_n^{(k)}.$$

Proof. There exists $q_{n+1}^{(r)}$ ways to tile a board of length $n + 2r$ containing at least r dominoes. Consider the number of dominoes at the end of each tiling. If tiling ends in at least r dominoes, then the final r dominoes cover cells $n + 1$ through $n + 2r$, while the remaining tilings can be done in q_{n+1} ways. On the other hand, if tilings ends in exactly $r - k$ dominoes for some $1 \leq k \leq r$, preceded by a square at position $n + 2k$ and contribute $a^{\xi(n)} b^{\xi(n+1)}$ to the weight, then the remaining $(n - 1 + 2k)$ -board can be tiled with at least k dominoes in $q_n^{(k)}$ ways. The result follows from the sum of over k , i.e.,

$$q_{n+1}^{(r)} = q_{n+1} + \sum_{k=1}^r a^{\xi(n)} b^{\xi(n+1)} q_n^{(k)} = q_{n-1} + \sum_{k=0}^r a^{\xi(n)} b^{\xi(n+1)} q_n^{(k)}.$$

□

Note that, if we take $a = b = x$, we get the following hyper-Fibonacci identity.

Corollary 2.2. *For $n, r \geq 1$, we have*

$$(2.6) \quad F_{n+1}^{(r)}(x) = F_{n-1}(x) + \sum_{k=0}^r xF_n^{(k)}(x).$$

For $a = b = 1$, we obtain the following identity, see [2],

$$F_{n+1}^{(r)} = F_{n-1} + \sum_{k=0}^r F_n^{(k)}.$$

In the following theorem, we give the recurrence relation of the bi-periodic hyper-Fibonacci sequence.

Theorem 2.6. *For $n \geq 0$ and $r \geq 2$, we have*

$$(2.7) \quad q_{n+2}^{(r)} = abq_n^{(r)} + 2q_{n+2}^{(r-1)} - q_{n+2}^{(r-2)}.$$

Proof. We will construct a 3-to-1 correspondence between the following two sets.

- The set of all tiled $(n + 2r - 1)$ -boards with at least r dominoes. There are $q_n^{(r)}$ ways.
- The set of all tiled $(n + 2r + 1)$ -boards with at least r dominoes and $(n + 2r - 3)$ -boards with at least $r - 1$ dominoes. There are $q_{n+2}^{(r)} + q_n^{(r-1)}$ ways.

Consider an arbitrary tiling T of length $n + 2r - 1$, we can do the following.

1. Add two squares at the end of T to get an $(n + 2r + 1)$ -board ending in a square. Then there are $abq_n^{(r)}$ ways.
2. Add a domino at the end of T to get an $(n + 2r + 1)$ -board ending in a domino. Then there are $q_{n+2}^{(r-1)}$ ways.
3. Condition on whether T ends in a square or a domino.
 - i. Suppose T ends in a square, then insert a domino immediately to the left of the square to creates $(n + 2r + 1)$ -board ending in a square. Then there are $a^{\xi(n+1)}b^{\xi(n)}q_{n+1}^{(r-1)}$ ways to do it.
 - ii. Suppose T ends in a domino, we remove the domino to get an $(n + 2r - 2)$ -board. Then there are $q_n^{(r-1)}$ ways.

So, we conclude that

$$\begin{aligned} q_{n+2}^{(r)} + q_n^{(r-1)} &= abq_n^{(r)} + q_{n+2}^{(r-1)} + a^{\xi(n+1)}b^{\xi(n)}q_{n+1}^{(r-1)} + q_n^{(r-1)} \\ &= abq_n^{(r)} + 2q_{n+2}^{(r-1)} + q_n^{(r-1)} - q_{n+2}^{(r-2)}. \end{aligned}$$

Therefore

$$q_{n+2}^{(r)} = abq_n^{(r)} + 2q_{n+2}^{(r-1)} - q_{n+2}^{(r-2)}.$$

□

Note that, if we take $a = b = 1$, we get the following hyper-Fibonacci identity.

Corollary 2.3. *For $n \geq 0$ and $r \geq 2$, we have*

$$(2.8) \quad F_{n+2}^{(r)} = F_n^{(r)} + 2F_{n+2}^{(r-1)} - F_{n+2}^{(r-2)}.$$

The following theorem gives the nonhomogeneous recurrence relation for the bi-periodic hyper-Fibonacci sequence.

Theorem 2.7. *For $n, r \geq 1$, we have*

$$(2.9) \quad q_{n+1}^{(r)} = a^{\xi(n)} b^{\xi(n+1)} q_n^{(r)} + q_{n-1}^{(r)} + a^{\xi(n)} (ab)^{\lfloor n/2 \rfloor} \binom{n+r-1}{r-1}.$$

Proof. There are $q_{n+1}^{(r)}$ ways to tile a $(n + 2r)$ -board with at least r dominoes. We consider the last tile in a tiling, which can be either a square or a domino. If the board ends in a square, then there are $bq_n^{(r)}$ ways to tile $(n + 2r - 1)$ -boards with at least r dominoes for n even and $aq_n^{(r)}$ ways to do it for n odd. If the board ends in a domino, we separate the tilings into two disjoint sets A and B . The set A with exactly r dominoes and the set B whose contain tilings with at least $r + 1$ dominoes. Having in mind that one domino is fixed, the tilings in the set A has $n + r - 1$ tiles with exactly $r - 1$ dominoes, then by Lemma 2.1, we have $|A| = a^{\xi(n)} (ab)^{\lfloor n/2 \rfloor} \binom{n+r-1}{r-1}$. The tilings in the set B are equivalent to the tilings of an $(n + 2r - 2)$ -boards with at least r dominoes, i.e., $|B| = q_{n-1}^{(r)}$. Therefore,

$$q_{n+1}^{(r)} = a^{\xi(n)} b^{\xi(n+1)} q_n^{(r)} + |A| + |B|. \quad \square$$

Note that, if we take $a = b = x$, we get the following hyper-Fibonacci identity, see [4],

$$F_{n+1}^{(r)}(x) = xF_n^{(r)}(x) + F_{n-1}^{(r)}(x) + x^n \binom{n+r-1}{r-1}.$$

Theorem 2.8. *For $m, n \in \mathbb{N} \cup \{0\}$ with $m \leq r$, we have*

$$(2.10) \quad q_{n+m}^{(r)} = \sum_{k=0}^m a^{\xi(n+m+1)\xi(n+k)} b^{\xi(n+m)\xi(n+k+1)} \binom{m}{k} (ab)^{\lfloor (m-k)/2 \rfloor} q_{n+k}^{(r-k)}.$$

Proof. There exists $q_{n+m}^{(r)}$ ways to tile a board of length $(n + m + 2r - 1)$ containing at least r dominoes. Consider the number of dominoes among the first m tiles. The k dominoes can be placed among the first m tiles in $\binom{m}{k}$ ways and the remaining tiles which consisting of squares, contribute $a^{\xi(n+m+1)\xi(n+k)} b^{\xi(n+m)\xi(n+k+1)} (ab)^{\lfloor (m-k)/2 \rfloor}$ to the weight. The remaining right board has a length of $n - 1 + 2r - k$, with at least $r - k$ dominos that can be tiled in $q_{n+k}^{(r-k)}$ ways. Summing over all possible k completes the proof. \square

Note that, if we take $a = b = x$ and $m = r$, we get the following hyper-Fibonacci identity, see [4],

$$F_{n+r}^{(r)} = \sum_{k=0}^r \binom{r}{k} x^{r-k} F_{n+k}^{(r-k)}.$$

The bi-periodic hyper-Fibonacci sequence can be expressed in terms of the combinatorial sum of bi-periodic Fibonacci sequence.

Theorem 2.9. For $n, r \geq 1$, we have

$$(2.11) \quad q_n^{(r)} = \sum_{k=1}^n a^{\xi(n+1)\xi(k)} b^{\xi(n)\xi(k+1)} \binom{n+r-k-1}{r-1} (ab)^{\lfloor (n-k)/2 \rfloor} q_k.$$

Proof. The left-hand side of this equality counts the number of ways to tile a board of length $n + 2r - 1$ containing at least r dominoes.

The right-hand side is obtained by conditioning on the location of the r th domino. Suppose that the r th domino occupies cell k and $k + 1$ ($1 \leq k \leq n$) (from the right). The left part is a tiling of some section of length $k - 1$ which can be done in q_k ways. The right part is a tiling of the remaining portion of length $n + 2r - 2 - k$ (i.e., cells $k + 2$ through $n + 2r - 1$) with exactly $r - 1$ dominos, which can be done in $a^{\xi(n+1)\xi(k)} b^{\xi(n)\xi(k+1)} \binom{n+r-k-1}{r-1} (ab)^{\lfloor (n-k)/2 \rfloor}$ ways (according to the parity of the numbers n and k). The result follows from considering the sum of all possible locations of the r th domino. \square

Note that, if we take $a = b = x$, we get the following hyper-Fibonacci identity, see [4],

$$F_n^{(r)}(x) = \sum_{k=1}^n x^{n-k} \binom{n+r-k-1}{r-1} F_k(x).$$

In the following theorem, we give the alternating binomial sum of the bi-periodic hyper-Fibonacci numbers.

Theorem 2.10. For $r, m, n \in \mathbb{N} \cup \{0\}$ with $m \leq r$, we have

$$(2.12) \quad \sum_{j=0}^m (-1)^j \binom{m}{j} q_{n+m}^{(r-j)} = a^{\xi(n)\xi(m)} b^{\xi(n+1)\xi(m)} (ab)^{\lfloor m/2 \rfloor} q_n^{(r)}.$$

Proof. We proceed by induction on $m \leq r$. For $m = 1$ and $m = 2$, we get (1.10) and Theorem 2.6, respectively. Suppose that the result holds for all $i \leq m$. Then we can prove it for $m + 1$

$$\begin{aligned} \sum_{j=0}^{m+1} (-1)^j \binom{m+1}{j} q_{n+m+1}^{(r-j)} &= \sum_{j=0}^{m+1} (-1)^j \left(\binom{m}{j} + \binom{m}{j-1} \right) q_{n+m+1}^{(r-j)} \\ &= \sum_{j=0}^m (-1)^j \binom{m}{j} q_{n+m+1}^{(r-j)} - \sum_{j \geq 0} (-1)^j \binom{m}{j} q_{n+m+1}^{(r-j-1)}. \end{aligned}$$

From (1.10), we obtain

$$\begin{aligned} \sum_{j=0}^{m+1} (-1)^j \binom{m+1}{j} q_{n+m+1}^{(r-j)} &= \sum_{j \geq 0} (-1)^j \binom{m}{j} a^{\xi(n+m)} b^{\xi(n+m+1)} q_{n+m}^{(r-j)} \\ &= a^{\xi(n+m)} b^{\xi(n+m+1)} a^{\xi(n)\xi(m)} b^{\xi(n+1)\xi(m)} (ab)^{\lfloor m/2 \rfloor} q_n^{(r)}. \end{aligned}$$

Using $\xi(n+m) = \xi(n) + \xi(m) - 2\xi(n)\xi(m)$ and $\lfloor m/2 \rfloor = \lfloor (m+1)/2 \rfloor - \xi(m)$, we get

$$\sum_{j=0}^{m+1} (-1)^j \binom{m+1}{j} q_{n+m+1}^{(r-j)} = a^{\xi(n)\xi(m+1)} b^{\xi(n+1)\xi(m+1)} (ab)^{\lfloor (m+1)/2 \rfloor} q_n^{(r)}.$$

Therefore, the identity is valid for all $m \leq r$. \square

Note that, for $a = b = x$, we get the following result.

Corollary 2.4. *The following equality holds for any nonnegative integers $r \geq m$*

$$(2.13) \quad \sum_{j=0}^m (-1)^j \binom{m}{j} F_{n+m}^{(r-j)} = x^m F_n^{(r)}.$$

The bi-periodic Fibonacci sequence can be expressed in terms of the bi-periodic hyper-Fibonacci sequence.

Theorem 2.11. *For $r, m \in \mathbb{N} \cup \{0\}$, we have*

$$(2.14) \quad q_{m+1} = \sum_{k=0}^m \binom{r}{k} (-1)^k a^{\xi(k)\xi(m)} b^{\xi(k)\xi(m+1)} (ab)^{\lfloor k/2 \rfloor} q_{m+1-k}^{(r)}.$$

Proof. We proceed by induction on m . This is true for $m = 0$. Suppose that the result holds for all $i \leq m$. Then we can prove it for $m + 1$. From (1.10), we get

$$\begin{aligned} q_{m+2} &= a^{\xi(m+1)} b^{\xi(m)} q_{m+1} + q_m \\ &= a^{\xi(m+1)} b^{\xi(m)} \sum_{k=0}^m \binom{r}{k} (-1)^k a^{\xi(k)\xi(m)} b^{\xi(k)\xi(m+1)} (ab)^{\lfloor k/2 \rfloor} q_{m+1-k}^{(r)} \\ &\quad + \sum_{k=0}^{m-1} \binom{r}{k} (-1)^k a^{\xi(k)\xi(m+1)} b^{\xi(k)\xi(m)} (ab)^{\lfloor k/2 \rfloor} q_{m-k}^{(r)}. \end{aligned}$$

Using $\xi(m+1) = \xi(m-k+1) + \xi(k)\xi(m+1) - \xi(k)\xi(m)$ and $\xi(m) = \xi(m-k) + \xi(k)\xi(m) - \xi(k)\xi(m+1)$ we get $\xi(k)\xi(m) + \xi(m+1) = \xi(k)\xi(m+1) + \xi(m-k+1)$ and $\xi(k)\xi(m+1) + \xi(m) = \xi(k)\xi(m) + \xi(m-k)$. Therefore, we have

$$\begin{aligned} q_{m+2} &= \sum_{k=0}^m \binom{r}{k} (-1)^k a^{\xi(k)\xi(m+1) + \xi(m-k+1)} b^{\xi(k)\xi(m) + \xi(m-k)} (ab)^{\lfloor k/2 \rfloor} q_{m+1-k}^{(r)} \\ &\quad + \sum_{k=0}^{m-1} \binom{r}{k} (-1)^k a^{\xi(k)\xi(m+1)} b^{\xi(k)\xi(m)} (ab)^{\lfloor k/2 \rfloor} q_{m-k}^{(r)} \\ &= \sum_{k \geq 0} \binom{r}{k} (-1)^k a^{\xi(k)\xi(m+1)} b^{\xi(k)\xi(m)} (ab)^{\lfloor k/2 \rfloor} \left(a^{\xi(m-k+1)} b^{\xi(m-k)} q_{m+1-k}^{(r)} + q_{m-k}^{(r)} \right) \\ &= \sum_{k=0}^{m+1} \binom{r}{k} (-1)^k a^{\xi(k)\xi(m+1)} b^{\xi(k)\xi(m)} (ab)^{\lfloor k/2 \rfloor} q_{m+2-k}^{(r)}. \end{aligned} \quad \square$$

Note that, for $a = b = x$, we get the following result.

Corollary 2.5. *The following equality holds for any integers $r, m \geq 0$*

$$(2.15) \quad F_{m+1}(x) = \sum_{k=0}^m \binom{r}{k} (-1)^k x^k F_{m+1-k}^{(r)}(x).$$

3. GENERATING FUNCTION

We start by establishing the relationship between the bi-periodic hyper-Fibonacci sequence and the hyper-Fibonacci polynomials.

Lemma 3.1. *For $n, r \geq 0$, we have*

$$(3.1) \quad q_n^{(r)} = \frac{1}{2} \left(\left(1 + \sqrt{\frac{a}{b}} \right) - (-1)^n \left(1 - \sqrt{\frac{a}{b}} \right) \right) F_n^{(r)}(\sqrt{ab}).$$

Proof. Using (1.4), (2.1) and $\lfloor n/2 \rfloor = (n - \xi(n))/2$, we have

$$\begin{aligned} q_n^{(r)} &= a^{\xi(n-1)} \sum_{k=r}^{\lfloor (n-1)/2 \rfloor + r} \binom{n-1+2r-k}{k} (ab)^{(n-1-\xi(n-1))/2+r-k} \\ &= \left(\frac{a}{\sqrt{ab}} \right)^{\xi(n-1)} \sum_{k=r}^{\lfloor (n-1)/2 \rfloor + r} \binom{n-1+2r-k}{k} (\sqrt{ab})^{n-1+2r-2k} \\ &= \left(\sqrt{\frac{a}{b}} \right)^{\xi(n-1)} \sum_{k=r}^{\lfloor (n-1)/2 \rfloor + r} \binom{n-1+2r-k}{k} (\sqrt{ab})^{n-1+2r-2k} \\ &= \frac{\left(1 + \sqrt{\frac{a}{b}} \right) - (-1)^n \left(1 - \sqrt{\frac{a}{b}} \right)}{2} \sum_{k=r}^{\lfloor (n-1)/2 \rfloor + r} \binom{n-1+2r-k}{k} (\sqrt{ab})^{n-1+2r-2k}. \quad \square \end{aligned}$$

Theorem 3.1. *The generating function of the bi-periodic hyper-Fibonacci sequence is given by*

$$\begin{aligned} \sum_{n \geq 0} q_n^{(r)} z^n &= \\ z \frac{\left(1 + \sqrt{\frac{a}{b}} \right) \left(1 + \sqrt{ab}z - z^2 \right) \left(1 + \sqrt{ab}z \right)^r + \left(1 - \sqrt{\frac{a}{b}} \right) \left(1 - \sqrt{ab}z - z^2 \right) \left(1 - \sqrt{ab}z \right)^r}{2 \left(1 - (ab+2)z^2 + z^4 \right) \left(1 - abz^2 \right)^r}. \end{aligned}$$

Proof. Using Lemma 3.1 and (1.5), we get

$$\begin{aligned} \sum_{n \geq 0} q_n^{(r)} z^n &= \frac{1}{2} \left(1 + \sqrt{\frac{a}{b}} \right) \sum_{n \geq 0} F_n^{(r)}(\sqrt{ab}) z^n - \frac{1}{2} \left(1 - \sqrt{\frac{a}{b}} \right) \sum_{n \geq 0} F_n^{(r)}(\sqrt{ab}) (-z)^n \\ &= \frac{1}{2} \left(1 + \sqrt{\frac{a}{b}} \right) \frac{z}{\left(1 - \sqrt{ab}z - z^2 \right) \left(1 - \sqrt{ab}z \right)^r} \\ &\quad - \frac{1}{2} \left(1 - \sqrt{\frac{a}{b}} \right) \frac{-z}{\left(1 + \sqrt{ab}z - z^2 \right) \left(1 + \sqrt{ab}z \right)^r}, \end{aligned}$$

which gives the desired result. □

Note that, if we take $r = 0$, we obtain the generating function of the bi-periodic Fibonacci sequence (1.7). If we take $a = b = x$, we obtain the generating function of hyper-Fibonacci polynomials (1.5) with $y = 1$.

REFERENCES

- [1] T. Amdeberhan, X. Chen, V. H. Moll and B. E. Sagan, *Generalized Fibonacci polynomials and Fibonacci coefficients*, Ann. Comb. **18**(4) (2014), 541–562. <https://doi.org/10.1007/s00026-014-0242-9>
- [2] M. Bahşi, I. Mezö and S. Solak, *A symmetric algorithm for hyper-Fibonacci and hyper-Lucas numbers*, Ann. Math. Inform. **43** (2014), 19–27.
- [3] H. Belbachir and A. Belkhir, *Combinatorial expressions involving Fibonacci, hyper-Fibonacci, and incomplete Fibonacci numbers*, J. Integer Seq. **17**(4) (2014), Article ID 14.4.3.
- [4] H. Belbachir and A. Belkhir, *On generalized hyper-Fibonacci and incomplete Fibonacci polynomials in arithmetic progresions*, Šiauliai Math. Semin. **11**(19) (2016), 1–12.
- [5] H. Belbachir and F. Bencherif, *On some properties of bivariate Fibonacci and Lucas polynomials*, J. Integer Seq. **11**(2) (2008), Article ID 08.2.6.
- [6] A. T. Benjamin and J. J. Quinn, *Proofs that Really Count: The Art of Combinatorial Proof*, Volume 27, The Mathematical Association of America, Washington, DC, 2003.
- [7] A. T. Benjamin, J. J. Quinn and F. E. Su, *Phased tilings and generalized Fibonacci identities*, Fibonacci Quart. **38**(3) (2000), 282–288.
- [8] G. Bilgici, *Two generalizations of Lucas sequence*, Appl. Math. Comput. **245** (2014), 526–538. <https://doi.org/10.1016/j.amc.2014.07.111>
- [9] N. N. Cao and F. Z. Zhao, *Some properties of hyperfibonacci and hyperlucas numbers*, J. Integer Seq. **13**(8) (2010), Article ID 10.8.8, 1–11.
- [10] A. Dil and I. Mezö, *A symmetric algorithm for hyperharmonic and Fibonacci numbers*, Appl. Math. Comput. **206**(2) (2008), 942–951. <https://doi.org/10.1016/j.amc.2008.10.013>
- [11] M. Edson, S. Lewis and O. Yayenie, *The k-periodic Fibonacci sequence and extended Binet's formula*, Integers **11**(A32) (2011), 739–751. <https://doi.org/10.1515/INTEG.2011.056>
- [12] M. Edson and O. Yayenie, *A new generalization of Fibonacci sequences and extended Binet's Formula*, Integers **9**(A48) (2009), 639–654. <https://doi.org/10.1515/INTEG.2009.051>
- [13] D. Panario, M. Sahin, and Q. Wan, *A family of Fibonacci-like conditional sequences*, Integers **13**(A78) (2013). <https://doi.org/10.1515/9783110298161.1042>
- [14] D. Panario, M. Sahin, Q. Wang and W. Webb, *General conditional recurrences*, Applied. Math. Comp. **243** (2014), 220–231. <https://doi.org/10.1016/j.amc.2014.05.108>
- [15] J. L. Ramirez, *Bi-periodic incomplete Fibonacci sequences*, Ann. Math. Inform. **42** (2013), 83–92.
- [16] E. Tan, *Some properties of the bi-periodic Horadam sequences*, Notes Number Theory Discrete Math. **23**(4) (2017), 56–65.
- [17] S. Uygun and E. Owusu, *A new generalization of Jacobsthal numbers (Bi-Periodic Jacobsthal Sequences)*, J. Math. Anal. **7**(5) (2016), 28–39. <https://doi.org/10.9734/jamcs/2019/v34i530226>
- [18] O. Yayenie, *A note on generalized Fibonacci sequence*, Appl. Math. Comput. **217**(12) (2011), 5603–5611. <https://doi.org/10.1016/j.amc.2010.12.038>

¹USTHB, FACULTY OF MATHEMATICS, RECITS LABORATORY,
Po. Box 32, EL ALIA 16111, BAB EZZOUAR, ALGIERS, ALGERIA

²CERIST: RESEARCH CENTER ON SCIENTIFIC AND TECHNICAL INFORMATION 05,
RUE DES 3 FRÈRES AISSOU, BEN AKNOUN, ALGIERS, ALGERIA
Email address: belaggounmanassil@gmail.com
Email address: hacenebelbachir@gmail.com