

DISCRETE LOCAL FRACTIONAL HILBERT-TYPE INEQUALITIES

PREDRAG VUKOVIĆ¹ AND WENGUI YANG²

ABSTRACT. The main objective of this paper is a study of some new discrete local fractional Hilbert-type inequalities. We apply our general results to homogeneous kernels. Also, the obtained results have the best possible constants.

1. INTRODUCTION

If $f(x), g(x) \geq 0$, such that $0 < \int_0^{+\infty} f^2(x)dx < +\infty$ and $0 < \int_0^{+\infty} g^2(x)dx < +\infty$, then we have (see [1]):

$$(1.1) \quad \int_0^{+\infty} \int_0^{+\infty} \frac{f(x)g(y)}{x+y} dx dy \leq \pi \left(\int_0^{+\infty} f^2(x)dx \int_0^{+\infty} g^2(y)dy \right)^{\frac{1}{2}},$$

where the constant π is the best possible. The inequality (1.1) is well known as Hilbert's integral inequality, which is important in mathematical analysis and its applications.

Over the last ten years, by using the kinds of generalized fractional integral operators, a great deal of fractional integral inequalities have been presented [2–5]. Recently, local fractional calculus has caused widespread attention from many scholars, we give basic definitions and results of the local fractional calculus (see [6–13]). Based on the local fractal identity and the generalized p -convexity, some novel Newton's type variants for the local differentiable functions were obtained in the paper [14]. Sarikaya et al. [15] established the generalized Grüss type inequality and some generalized Čebyšev type inequalities for local fractional integrals on fractal sets. According to the identity

Key words and phrases. Hilbert inequality, conjugate parameters, homogeneous function, local fractional calculus.

2020 *Mathematics Subject Classification.* Primary: 26D15. Secondary: 26A33.

DOI 10.46793/KgJMat2506.899V

Received: October 27, 2022.

Accepted: April 06, 2023.

involving local fractional integrals, Iftikhar et al. [16] presented some new Newton-type inequalities for functions with the local fractional derivatives. By employing the local fractional integrals, Akkurt et al. [17] investigated the generalized Ostrowski type integral inequalities involving moments of continuous random variables. Sarikaya and Budak [18] gave a generalized Ostrowski inequality and some new inequalities using the generalized convex function for local fractional integrals on fractal sets. Based on two local fractional integral operators with Mittag-Leffler kernel, Sun [19] obtained some Hermite-Hadamard and Hermite-Hadamard-Fejér-type local fractional integral inequalities for generalized preinvex functions on Yang's fractal sets.

For the sake of convenience, we recall Yang's fractal set Ω^α , where the set Ω is called base set of fractional set, and α denotes the dimension of cantor set, $0 < \alpha \leq 1$. The α -type set of integers \mathbb{Z}^α is defined by (see [6–8])

$$\mathbb{Z}^\alpha := \{0^\alpha\} \cup \{\pm m^\alpha : m \in \mathbb{N}\}.$$

The α -type set of rational numbers \mathbb{Q}^α is defined by

$$\mathbb{Q}^\alpha := \{q^\alpha : q \in \mathbb{Q}\} = \left\{ \left(\frac{m}{n} \right)^\alpha : m \in \mathbb{Z}, n \in \mathbb{N} \right\}.$$

The α -type set of irrational numbers \mathbb{J}^α is defined by

$$\mathbb{J}^\alpha := \{r^\alpha : r \in \mathbb{J}\} = \left\{ r^\alpha \neq \left(\frac{m}{n} \right)^\alpha : m \in \mathbb{Z}, n \in \mathbb{N} \right\}.$$

The α -type set of real line numbers \mathbb{R}^α is defined by

$$\mathbb{R}^\alpha = \mathbb{Q}^\alpha \cup \mathbb{J}^\alpha.$$

Some basic operation rules on \mathbb{R}^α are presented as follows: If $a^\alpha, b^\alpha, c^\alpha \in \mathbb{R}^\alpha$, then

- (a1) $a^\alpha + b^\alpha \in \mathbb{R}^\alpha, a^\alpha b^\alpha \in \mathbb{R}^\alpha$;
- (a2) $a^\alpha + b^\alpha = b^\alpha + a^\alpha = (a + b)^\alpha = (b + a)^\alpha$;
- (a3) $a^\alpha + (b^\alpha + c^\alpha) = (a + b)^\alpha + c^\alpha$;
- (a4) $a^\alpha b^\alpha = b^\alpha a^\alpha = (ab)^\alpha = (ba)^\alpha$;
- (a5) $a^\alpha (b^\alpha c^\alpha) = (a^\alpha b^\alpha) c^\alpha$;
- (a6) $a^\alpha (b^\alpha + c^\alpha) = a^\alpha b^\alpha + a^\alpha c^\alpha$;
- (a7) $a^\alpha + 0^\alpha = 0^\alpha + a^\alpha = a^\alpha$ and $a^\alpha 1^\alpha = 1^\alpha a^\alpha = a^\alpha$;
- (a8) for each $a^\alpha \in \mathbb{R}^\alpha$, its inverse element $(-a^\alpha)$ may be written as $-a^\alpha$; for each $b^\alpha \in \mathbb{R}^\alpha \setminus \{0^\alpha\}$, its inverse element $(1/b)^\alpha$ may be written as $1^\alpha/b^\alpha$ but not as $1/b^\alpha$;
- (a9) $a^\alpha < b^\alpha$ if and only if $a < b$;
- (a10) $a^\alpha = b^\alpha$ if and only if $a = b$.

Further, we define the local fractional derivative and integral.

Definition 1.1. A non-differentiable function $f(x)$ is said to be local fractional continuous at $x = x_0$ if for each $\varepsilon > 0$, there exists for $\delta > 0$ such that

$$|f(x) - f(x_0)| < \varepsilon^\alpha,$$

holds for $0 < |x - x_0| < \delta$. If a function f is local continuous on the interval (a, b) , we denote $f \in C_\alpha(a, b)$.

Definition 1.2. Let $f(x) \in C_\alpha[a, b]$. Local fractional derivative of the function $f(x)$ at $x = x_0$ is given by

$$f^{(\alpha)}(x_0) = \left. \frac{d^\alpha f(x)}{dx^\alpha} \right|_{x=x_0} = \lim_{x \rightarrow x_0} \frac{\Gamma(1+\alpha)(f(x) - f(x_0))}{(x - x_0)^\alpha}.$$

Definition 1.3. Let $f(x) \in C_\alpha[a, b]$ and let $P = \{t_0, t_1, \dots, t_N\}$, $N \in \mathbb{N}$, be a partition of interval $[a, b]$ such that $a = t_0 < t_1 < \dots < t_{N-1} < t_N = b$. Further, for this partition P , let $\Delta t_j = t_{j+1} - t_j$, $j = 0, \dots, N-1$, and $\Delta t = \max\{\Delta t_1, \Delta t_2, \dots, \Delta t_{N-1}\}$. Then the local fractional integral of f on the interval $[a, b]$ of order α (denoted by ${}_a I_b^\alpha f(x)$) is defined by

$${}_a I_b^\alpha f(x) = \frac{1}{\Gamma(1+\alpha)} \int_a^b f(t)(dt)^\alpha = \frac{1}{\Gamma(1+\alpha)} \lim_{\Delta t \rightarrow 0} \sum_{j=0}^{N-1} f(t_j)(\Delta t_j)^\alpha.$$

The above definition implies that ${}_a I_b^{(\alpha)} f(x) = 0$ if $a = b$, and ${}_a I_b^{(\alpha)} f(x) = -{}_b I_a^{(\alpha)} f(x)$ if $a < b$. If for any $x \in [a, b]$, there exists ${}_a I_x^{(\alpha)} f(x)$, then we denote by $f(x) \in I_x^{(\alpha)}[a, b]$.

At the end of this summary, we give some useful formulas:

- (b1) $\frac{d^\alpha x^{k\alpha}}{dx^\alpha} = \frac{\Gamma(1+k\alpha)}{\Gamma(1+(k-1)\alpha)} x^{(k-1)\alpha}$, $k > 0$;
- (b2) $\frac{d^\alpha E_\alpha((cx)^\alpha)}{dx^\alpha} = c^\alpha E_\alpha((cx)^\alpha)$, where $E_\alpha(\cdot)$ denotes the Mittag-Leffler function given by $E_\alpha(x^\alpha) = \sum_{k=0}^{+\infty} \frac{x^{k\alpha}}{\Gamma(1+k\alpha)}$;
- (b3) If $y(x) = (f \circ g)(x)$, then $\frac{d^\alpha y(x)}{dx^\alpha} = f^{(\alpha)}(g(x))(g'(x))^\alpha$;
- (b4) $\frac{1}{\Gamma(1+\alpha)} \int_a^b E_\alpha(x^\alpha)(dx)^\alpha = E_\alpha(b^\alpha) - E_\alpha(a^\alpha)$;
- (b5) $\frac{1}{\Gamma(1+\alpha)} \int_a^b x^{k\alpha}(dx)^\alpha = \frac{\Gamma(1+k\alpha)}{\Gamma(1+(k+1)\alpha)} (b^{(k+1)\alpha} - a^{(k+1)\alpha})$, $k > 0$;
- (b6) $B_\alpha(a, b) = \frac{1}{\Gamma(1+\alpha)} \int_0^{+\infty} \frac{x^{\alpha(b-1)}}{(1+x^\alpha)^{a+b}} (dx)^\alpha$, where $B_\alpha(a, b)$ denotes local fractional Beta function.

In this paper, by using the way of weight functions and the technique of local fractional calculus, a new Hilbert-type discrete inequality with homogeneous kernel and a best constant is built. As applications, the equivalent form and some particular cases are obtained.

2. MAIN RESULTS

The starting point in the researching Hilbert-type inequalities is the well-known Hölder's inequality. A fractal version of Hölder's inequality is presented in the following lemma.

Lemma 2.1 ([8]). *Let $1/p + 1/q = 1$, $p > 1$, and let $(a_m)_{m \in \mathbb{N}}$ and $(b_n)_{n \in \mathbb{N}}$ be non-negative real sequences. Then*

$$\sum_{i=1}^n a_i^\alpha b_i^\alpha \leq \left(\sum_{i=1}^n a_i^{\alpha p} \right)^{\frac{1}{p}} \left(\sum_{i=1}^n b_i^{\alpha q} \right)^{\frac{1}{q}}.$$

If $\sum_{i=1}^{+\infty} a_i^{\alpha p} < +\infty$ and $\sum_{i=1}^{+\infty} b_i^{\alpha q} < +\infty$, then the following inequality holds

$$\sum_{i=1}^{+\infty} a_i^\alpha b_i^\alpha \leq \left(\sum_{i=1}^{+\infty} a_i^{\alpha p} \right)^{\frac{1}{p}} \left(\sum_{i=1}^{+\infty} b_i^{\alpha q} \right)^{\frac{1}{q}}.$$

In particular, a two-variable version of the fractal Hölder's inequality is given in the next lemma.

Lemma 2.2. *Let $1/p + 1/q = 1$, $p > 1$, and let $h, F, G \in C_\alpha(\mathbb{R}_+^2)$ be non-negative functions. If*

$$0 < \sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) F^p(m, n) < +\infty, \quad 0 < \sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) G^q(m, n) < +\infty,$$

then the following inequality holds

$$(2.1) \quad \sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) F(m, n) G(m, n) \leq \left(\sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) F^p(m, n) \right)^{\frac{1}{p}} \times \left(\sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) G^q(m, n) \right)^{\frac{1}{q}}.$$

Proof. The inequality (2.1) is trivially true in the case when h or F or G is identically zero. Suppose that

$$\left(\sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) F^p(m, n) \right) \left(\sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) G^q(m, n) \right) \neq 0.$$

Applying the following α -Young's inequality

$$x_i^{\frac{\alpha}{p}} y_i^{\frac{\alpha}{q}} \leq \frac{x_i^\alpha}{p^\alpha} + \frac{y_i^\alpha}{q^\alpha}, \quad x_i, y_i \geq 0, \quad \text{and} \quad \frac{1}{p} + \frac{1}{q} = 1, \quad p > 1,$$

to

$$x^\alpha := \frac{h(m, n) F^p(m, n)}{\sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) F^p(m, n)}$$

and

$$y^\alpha := \frac{h(m, n) G^q(m, n)}{\sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) G^q(m, n)},$$

we can obtain

$$\begin{aligned} & \frac{[h(m, n)]^{\frac{1}{p}} F(m, n) [h(m, n)]^{\frac{1}{q}} G(m, n)}{\left(\sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) F^p(m, n)\right)^{\frac{1}{p}} \left(\sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) G^q(m, n)\right)^{\frac{1}{q}}} \\ & \leq \frac{1}{p^\alpha} \cdot \frac{h(m, n) F^p(m, n)}{\sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) F^p(m, n)} + \frac{1}{q^\alpha} \cdot \frac{h(m, n) G^q(m, n)}{\sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) G^q(m, n)}. \end{aligned}$$

Summarizing both side of the obtained inequality, we have

$$\begin{aligned} & \frac{\sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) F(m, n) G(m, n)}{\left(\sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) F^p(m, n)\right)^{\frac{1}{p}} \left(\sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) G^q(m, n)\right)^{\frac{1}{q}}} \\ & \leq \frac{1}{p^\alpha} \cdot \frac{\sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) F^p(m, n)}{\sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) F^p(m, n)} + \frac{1}{q^\alpha} \cdot \frac{\sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) G^q(m, n)}{\sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} h(m, n) G^q(m, n)} \\ & = \frac{1}{p^\alpha} + \frac{1}{q^\alpha} = 1^\alpha. \end{aligned}$$

This directly gives the desired inequality (2.1). The proof is completed. \square

Besides, we introduce the following notation and definition (see [21]).

Definition 2.1. Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^\alpha$. If the following inequality

$$(2.2) \quad f(\lambda x_1 + (1 - \lambda)x_2) \leq \lambda^\alpha f(x_1) + (1 - \lambda)^\alpha f(x_2)$$

holds, for any $x_1, x_2 \in I$ and $\lambda \in [0, 1]$, then f is said to be a generalized convex function on I .

Mo et al. [21] proved the following generalized Hermite-Hadamard inequality for local fractional integral. Let $f \in I_x^{(\alpha)}[a, b]$ be a generalized convex function on $[a, b]$ with $a < b$. Then

$$(2.3) \quad f\left(\frac{a+b}{2}\right) \leq \frac{\Gamma(1+\alpha)}{(b-a)^\alpha} {}_a I_b^{(\alpha)} f \leq \frac{f(a) + f(b)}{2^\alpha}.$$

Applying above inequality we can prove next lemma.

Lemma 2.3. If $f \in I_x^{(\alpha)}(\mathbb{R}_+)$, $f^{(\alpha)}(t) < 0$, $f^{(2\alpha)}(t) > 0$, $t \in (1/2, +\infty)$, then we have

$$(2.4) \quad \frac{1}{\Gamma(1+\alpha)} \int_1^{+\infty} f(t)(dt)^\alpha \leq \frac{1}{\Gamma(1+\alpha)} \sum_{n=1}^{+\infty} f(n) \leq \frac{1}{\Gamma(1+\alpha)} \int_{\frac{1}{2}}^{+\infty} f(t)(dt)^\alpha.$$

Proof. Setting $a = n - \frac{1}{2}$, $b = n + \frac{1}{2}$, the generalized Hermite-Hadamard inequality (2.3) yields

$$(2.5) \quad \frac{f(n)}{\Gamma(1+\alpha)} \leq \frac{1}{\Gamma(1+\alpha)} \int_{n-\frac{1}{2}}^{n+\frac{1}{2}} f(t)(dt)^\alpha.$$

Similarly, for $a = n$, $b = n + 1$, from (2.3) we get

$$(2.6) \quad \frac{1}{\Gamma(1+\alpha)} \int_n^{n+1} f(t)(dt)^\alpha \leq \frac{f(n)}{\Gamma(1+\alpha)}.$$

Now, from (2.5) and (2.6) we obtain

$$\frac{1}{\Gamma(1+\alpha)} \int_n^{n+1} f(t)(dt)^\alpha \leq \frac{f(n)}{\Gamma(1+\alpha)} \leq \frac{1}{\Gamma(1+\alpha)} \int_{n-\frac{1}{2}}^{n+\frac{1}{2}} f(t)(dt)^\alpha.$$

Furthermore, we can obtain

$$\begin{aligned} \sum_{n=1}^{+\infty} \frac{1}{\Gamma(1+\alpha)} \int_n^{n+1} f(t)(dt)^\alpha &= \frac{1}{\Gamma(1+\alpha)} \int_1^{+\infty} f(t)(dt)^\alpha \\ &\leq \frac{1}{\Gamma(1+\alpha)} \sum_{n=1}^{+\infty} f(n) \leq \sum_{n=1}^{+\infty} \frac{1}{\Gamma(1+\alpha)} \int_{n-\frac{1}{2}}^{n+\frac{1}{2}} f(t)(dt)^\alpha = \frac{1}{\Gamma(1+\alpha)} \int_{\frac{1}{2}}^{+\infty} f(t)(dt)^\alpha, \end{aligned}$$

which implies (2.4) holds. This completes the proof. \square

Suppose that $r > 0$ and $K(x, y)$ is strictly decreasing and generalized convex function in both variables on \mathbb{R}_+ . Using chain rule for local fractional derivative (the formula (b3) from Introduction) yields

$$\frac{\partial^\alpha}{\partial x^\alpha} K(x, n) x^{-\alpha r} = \frac{1}{x^{\alpha r}} \cdot \frac{\partial^\alpha}{\partial x^\alpha} [K(x, n)] - \frac{\Gamma(1+r\alpha)}{\Gamma(1+(r-1)\alpha)} \cdot \frac{K(x, n)}{x^{\alpha(r+1)}} < 0$$

and

$$\begin{aligned} \frac{\partial^{2\alpha}}{\partial x^{2\alpha}} K(x, n) x^{-\alpha r} &= \frac{1}{x^{\alpha r}} \cdot \frac{\partial^{2\alpha}}{\partial x^{2\alpha}} [K(x, n)] - \frac{\Gamma(1+r\alpha)}{\Gamma(1+(r-1)\alpha)} \cdot \frac{K(x, n)}{x^{\alpha(r+1)}} \\ &\quad \times \frac{\partial^\alpha}{\partial x^\alpha} [K(x, n)] > 0, \end{aligned}$$

for $x > 0$ and $n \in \mathbb{N}$. In this way (see also [22], Corollary 1) we obtain the following result.

Lemma 2.4. *Let $r > 0$, $m, n \in \mathbb{N}$, and $K(x, y)$ be strictly decreasing and generalized convex function in both variables on \mathbb{R}_+ . Then*

$$K(m, y)y^{-\alpha r} \quad \text{and} \quad K(x, n)x^{-\alpha r}$$

are strictly decreasing and generalized convex function on \mathbb{R}_+ .

In what follows we suppose that $K \in C_\alpha(\mathbb{R}_+^2)$ is a non-negative homogeneous function of degree $-\alpha s$, $s > 0$. Further, we define

$$(2.7) \quad k(\beta) = \frac{1}{\Gamma(1+\alpha)} \int_1^{+\infty} K(1, t)t^{-\alpha\beta}(dt)^\alpha,$$

under assumption $k(\beta) < +\infty$.

To prove our main results we need some technical lemma.

Lemma 2.5. *Let $1/p + 1/q = 1$, $p > 1$, and let $K \in C_\alpha(\mathbb{R}_+^2)$ be a non-negative homogeneous function of degree $-\alpha s$, $s > 0$. If K is strictly decreasing and generalized*

convex function in both variables on \mathbb{R}_+ , then

$$(2.8) \quad \omega_m(pA_2) := \sum_{n=1}^{+\infty} K(m, n) \left(\frac{m}{n}\right)^{\alpha p A_2} \leq \Gamma(1 + \alpha) m^{\alpha(1-s)} k(pA_2)$$

and

$$(2.9) \quad \bar{\omega}_n(qA_1) := \sum_{m=1}^{+\infty} K(m, n) \left(\frac{n}{m}\right)^{\alpha q A_1} \leq \Gamma(1 + \alpha) n^{\alpha(1-s)} k(2 - s - qA_1),$$

where $A_1 \in (\max\{(1-s)/q, 0\}, 1/q)$ and $A_2 \in (\max\{(1-s)/p, 0\}, 1/p)$.

Proof. Applying Lemma 2.2 and Lemma 2.4 we get

$$\omega_m(pA_2) \leq \Gamma(1 + \alpha) \frac{1}{\Gamma(1 + \alpha)} \int_0^{+\infty} K(m, x) \left(\frac{x}{m}\right)^{-\alpha p A_2} (dx)^\alpha.$$

Further, using homogeneity of function K and substituting $u = x/m$, we have

$$\begin{aligned} \omega_m(pA_2) &\leq \Gamma(1 + \alpha) m^{\alpha(1-s)} \frac{1}{\Gamma(1 + \alpha)} \int_0^{+\infty} K(1, u) u^{-\alpha p A_2} (du)^\alpha \\ &= \Gamma(1 + \alpha) m^{\alpha(1-s)} k(pA_2), \end{aligned}$$

which implies (2.8), where we used the definition of $k(\beta)$ in equation (2.7). Similarly, we obtain (2.9). \square

The main results are stated below.

Theorem A. Let $1/p + 1/q = 1$, $p > 1$, and let $(a_m)_{m \in \mathbb{N}}$ and $(b_n)_{n \in \mathbb{N}}$ be non-negative real sequences. If $K(x, y)$, A_1 , A_2 are defined as in Lemma 2.5, then the following inequalities hold and are equivalent

$$(2.10) \quad I := \sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} K(m, n) a_m^\alpha b_n^\alpha \leq L \left(\sum_{m=1}^{+\infty} m^{\alpha(1-s) + \alpha p(A_1 - A_2)} a_m^\alpha \right)^{\frac{1}{p}} \times \left(\sum_{n=1}^{+\infty} n^{\alpha(1-s) + \alpha q(A_2 - A_1)} b_n^\alpha \right)^{\frac{1}{q}}$$

and

$$(2.11) \quad J := \left(\sum_{n=1}^{+\infty} n^{\alpha(s-1)(p-1) + \alpha p(A_1 - A_2)} \left(\sum_{m=1}^{+\infty} K(m, n) a_m^\alpha \right)^p \right)^{\frac{1}{p}} \leq L \left(\sum_{m=1}^{+\infty} m^{\alpha(1-s) + \alpha p(A_1 - A_2)} a_m^\alpha \right)^{\frac{1}{p}},$$

where $L = \Gamma(1 + \alpha) k(pA_2)^{1/p} k(2 - s - qA_1)^{1/q}$.

Proof. By using the local fractional Hölder's inequality (2.1), we have

$$\begin{aligned}
 I &= \sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} K(m, n) a_m^\alpha \frac{m^{\alpha A_1}}{n^{\alpha A_2}} b_n^\alpha \frac{n^{\alpha A_2}}{m^{\alpha A_1}} \\
 &\leq \left(\sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} K(m, n) \frac{m^{\alpha p A_1}}{n^{\alpha p A_2}} a_m^{\alpha p} \right)^{\frac{1}{p}} \left(\sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} K(m, n) \frac{n^{\alpha q A_2}}{m^{\alpha q A_1}} b_n^{\alpha q} \right)^{\frac{1}{q}} \\
 &= \left(\sum_{m=1}^{+\infty} \left(\sum_{n=1}^{+\infty} K(m, n) \left(\frac{m}{n} \right)^{\alpha p A_2} \right) m^{\alpha p(A_1 - A_2)} a_m^{\alpha p} \right)^{\frac{1}{p}} \\
 &= \left(\sum_{n=1}^{+\infty} \left(\sum_{m=1}^{+\infty} K(m, n) \left(\frac{n}{m} \right)^{\alpha q A_1} \right) n^{\alpha q(A_2 - A_1)} b_n^{\alpha q} \right)^{\frac{1}{q}}.
 \end{aligned}$$

Now, from Lemma 2.5, we get the inequality (2.10).

We suppose that the inequality (2.10) is valid. To obtain (2.11), we set

$$b_n^\alpha := n^{\alpha(s-1)(p-1) + \alpha p(A_1 - A_2)} \left(\sum_{m=1}^{+\infty} K(m, n) a_m^\alpha \right)^{p-1}.$$

It follows that

$$J^p = \sum_{n=1}^{+\infty} n^{\alpha(1-s) + \alpha q(A_2 - A_1)} b_n^{\alpha q}.$$

By using the inequality (2.10), we have

$$\begin{aligned}
 &\sum_{n=1}^{+\infty} n^{\alpha(s-1)(p-1) + \alpha p(A_1 - A_2)} \left(\sum_{m=1}^{+\infty} K(m, n) a_m^\alpha \right)^p = J^p = I \\
 &\leq L \left(\sum_{m=1}^{+\infty} m^{\alpha(1-s) + \alpha p(A_1 - A_2)} a_m^{\alpha p} \right)^{\frac{1}{p}} \left(\sum_{n=1}^{+\infty} n^{\alpha(1-s) + \alpha q(A_2 - A_1)} b_n^{\alpha q} \right)^{\frac{1}{q}},
 \end{aligned}$$

which implies the inequality (2.11) holds. By using the two dimensional Hölder's inequality in Lemma 2.1, we have

$$\begin{aligned}
 I &= \sum_{n=1}^{+\infty} \left(n^{\alpha(s-1)\frac{1}{q} + \alpha(A_1 - A_2)} \left(\sum_{m=1}^{+\infty} K(m, n) a_m^\alpha \right) \right) n^{\alpha(1-s)\frac{1}{q} + \alpha(A_2 - A_1)} b_n^\alpha \\
 &\leq J \left(\sum_{n=1}^{+\infty} n^{\alpha(1-s) + \alpha q(A_2 - A_1)} b_n^{\alpha q} \right)^{\frac{1}{q}}.
 \end{aligned}$$

From (2.11) and the above inequality, we have (2.10). Therefore, the inequalities (2.11) and (2.10) are equivalent. \square

Now, we consider some special choices of the parameters A_1 and A_2 . More precisely, let the parameters A_1 and A_2 satisfy condition

$$(2.12) \quad pA_2 + qA_1 = 2 - s.$$

Then, the constant L from Theorem A becomes

$$(2.13) \quad L^* = \Gamma(1 + \alpha)k(pA_2).$$

Further, the inequalities (2.10) and (2.11) take form

$$(2.14) \quad \sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} K(m, n) a_m^\alpha b_n^\alpha \leq L^* \left(\sum_{m=1}^{+\infty} m^{-\alpha + \alpha p q A_1} a_m^{\alpha p} \right)^{\frac{1}{p}} \left(\sum_{n=1}^{+\infty} n^{-\alpha + \alpha p q A_2} b_n^{\alpha q} \right)^{\frac{1}{q}}$$

and

$$(2.15) \quad \left(\sum_{n=1}^{+\infty} n^{\alpha(p-1)(1-pqA_2)} \left(\sum_{m=1}^{+\infty} K(m, n) a_m^\alpha \right)^p \right)^{\frac{1}{p}} \leq L^* \left(\sum_{m=1}^{+\infty} m^{-\alpha + \alpha p q A_1} a_m^{\alpha p} \right)^{\frac{1}{p}}.$$

In the following theorem we show that, if the parameters A_1 and A_2 satisfy condition (2.12), then one obtains the best possible constant.

Theorem B. *Let s , A_1 , A_2 and $K(x, y)$ be defined as in Theorem A. If the parameters A_1 and A_2 satisfy condition $pA_2 + qA_1 = 2 - s$, then the constant $L^* = \Gamma(1 + \alpha)k(pA_2)$ in inequalities (2.14) and (2.15) is the best possible.*

Proof. For this purpose, set $\tilde{a}_m^\alpha = m^{-\alpha q A_1 - \frac{\alpha \varepsilon}{p}}$ and $\tilde{b}_n^\alpha = n^{-\alpha p A_2 - \frac{\alpha \varepsilon}{q}}$ where $0 < \varepsilon < \frac{1-pA_2}{q}$. Now, let us suppose that the inequality (2.14) is valid. By using Lemma 2.2, we have

$$\begin{aligned} \frac{1}{\Gamma(1 + \alpha)\varepsilon^\alpha} &= \frac{1}{\Gamma(1 + \alpha)} \int_1^{+\infty} u^{-\alpha - \alpha \varepsilon} (du)^\alpha \leq \frac{1}{\Gamma(1 + \alpha)} \sum_{m=1}^{+\infty} m^{-\alpha - \alpha \varepsilon} \\ &= \frac{1}{\Gamma(1 + \alpha)} \sum_{m=1}^{+\infty} m^{-\alpha + \alpha p q A_1} \tilde{a}_m^{\alpha p} \\ &\leq \frac{1}{\Gamma(1 + \alpha)} \int_{\frac{1}{2}}^{+\infty} u^{-\alpha - \alpha \varepsilon} (du)^\alpha + \frac{1}{\Gamma(1 + \alpha)} \int_1^{+\infty} u^{-\alpha - \alpha \varepsilon} (du)^\alpha. \end{aligned}$$

Hence, we obtain

$$(2.16) \quad \frac{1}{\Gamma(1 + \alpha)} \sum_{m=1}^{+\infty} m^{-\alpha + \alpha p q A_1} \tilde{a}_m^{\alpha p} \leq \frac{1}{\varepsilon^\alpha \Gamma(1 + \alpha)} + O(1),$$

and similarly

$$(2.17) \quad \frac{1}{\Gamma(1 + \alpha)} \sum_{n=1}^{+\infty} n^{-\alpha + \alpha p q A_2} \tilde{b}_n^{\alpha q} \leq \frac{1}{\varepsilon^\alpha \Gamma(1 + \alpha)} + O(1).$$

Suppose that the constant L^* , defined by (2.13), is not the best possible in inequality (2.14). That implies that there exists constant M , smaller than L^* , such that the inequality (2.14) is still valid if we replace L^* with M . Hence, if we insert relations (2.16) and (2.17) in inequality (2.14), with the constant M instead of L^* , we have

$$(2.18) \quad \sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} K(m, n) \tilde{a}_m^\alpha \tilde{b}_n^\alpha \leq \frac{1}{\varepsilon^\alpha} (M + o(1)).$$

Further, we estimate the left-hand side of inequality (2.14). Namely, if we insert the above defined sequences $(\tilde{a}_m^\alpha)_{m \in \mathbb{N}}$ and $(\tilde{b}_n^\alpha)_{n \in \mathbb{N}}$ in the left-hand side of (2.14), we easily obtain the inequality

$$(2.19) \quad \begin{aligned} J_\varepsilon &:= \frac{1}{\Gamma^2(1+\alpha)} \sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} K(m, n) \tilde{a}_m^\alpha \tilde{b}_n^\alpha \\ &\geq \frac{1}{\Gamma(1+\alpha)} \int_1^{+\infty} x^{-\alpha q A_1 - \frac{\alpha \varepsilon}{p}} \left(\frac{1}{\Gamma(1+\alpha)} \int_1^{+\infty} K(x, y) y^{-\alpha p A_2 - \frac{\alpha \varepsilon}{q}} (dy)^\alpha \right) (dx)^\alpha, \end{aligned}$$

where we used Lemma 2.2. By using the substitution $u = y/x$, we obtain

$$(2.20) \quad J_\varepsilon \geq \frac{1}{\Gamma(1+\alpha)} \int_1^{+\infty} x^{-\alpha - \alpha \varepsilon} \left(\frac{1}{\Gamma(1+\alpha)} \int_{\frac{1}{x}}^{+\infty} K(1, u) u^{-\alpha p A_2 - \frac{\alpha \varepsilon}{q}} (du)^\alpha \right) (dx)^\alpha.$$

Further, since the kernel K is strictly decreasing in both variables, it follows that $K(1, 0) > K(1, t)$, for $t > 0$, so we have

$$\begin{aligned} &\frac{1}{\Gamma(1+\alpha)} \int_{\frac{1}{x}}^{+\infty} K(1, u) u^{-\alpha p A_2 - \frac{\alpha \varepsilon}{q}} (du)^\alpha \\ &> \frac{1}{\Gamma(1+\alpha)} \int_0^{+\infty} K(1, u) u^{-\alpha p A_2 - \frac{\alpha \varepsilon}{q}} (du)^\alpha - \frac{K(1, 0)}{\Gamma(1+\alpha)} \int_0^{\frac{1}{x}} K(1, u) u^{-\alpha p A_2 - \frac{\alpha \varepsilon}{q}} (du)^\alpha \\ &= k \left(p A_2 + \frac{\varepsilon}{q} \right) - \frac{K(1, 0)}{\Gamma(1+\alpha)(1 - p A_2 - \frac{\varepsilon}{q})^\alpha} x^{\alpha p A_2 + \frac{\alpha \varepsilon}{q} - \alpha} \end{aligned}$$

and, consequently,

$$(2.21) \quad J_\varepsilon \geq \frac{1}{\varepsilon^\alpha} \cdot \frac{k \left(p A_2 + \frac{\varepsilon}{q} \right)}{\Gamma(1+\alpha)} + \frac{K(1, 0)}{\Gamma^2(1+\alpha)} \cdot \frac{1}{(1 - p A_2 - \frac{\varepsilon}{q})^\alpha (p A_2 - \frac{\varepsilon}{p} - 1)^\alpha}.$$

Now, the relations (2.19), (2.20) and (2.21) yield the estimate for the left-hand side of inequality (2.14):

$$(2.22) \quad \sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} K(m, n) \tilde{a}_m^\alpha \tilde{b}_n^\alpha > \frac{1}{\varepsilon^\alpha} (L^* + o(1)).$$

Finally, by comparing (2.18) and (2.22), and by letting $\varepsilon \rightarrow 0^+$, we get that $L^* \leq M$, which contradicts with the assumption that the constant M is smaller than L^* .

The equivalence of inequalities (2.14) and (2.15) means that the constant L^* is the best possible in the inequality (2.15). The proof is now completed. \square

As corollaries of Theorem B we have the following results. We processed with the kernel $K_1(x, y) = (x + y)^{-\alpha s}$, $s > 0$. By using local fractional calculus, we have

$$\frac{\partial^\alpha}{\partial x^\alpha} \cdot \frac{1}{(m + x)^{\alpha s}} = - \frac{\Gamma(1 + s\alpha)}{\Gamma(1 + (s - 1)\alpha)} \cdot \frac{1}{(m + x)^{\alpha(s+1)}} < 0, \quad x > 0,$$

and similarly

$$\frac{\partial^{2\alpha}}{\partial x^{2\alpha}} \cdot \frac{1}{(m+x)^{\alpha s}} = \frac{\Gamma(1+(s+1)\alpha)}{\Gamma(1+(s-1)\alpha)} \cdot \frac{1}{(m+x)^{\alpha(s+2)}} > 0, \quad x > 0.$$

Now, by applying Lemma 2.4 we obtain

$$\frac{\partial^\alpha}{\partial x^\alpha} K_1(x, y) x^{-\alpha r} < 0 \quad \text{and} \quad \frac{\partial^{2\alpha}}{\partial x^{2\alpha}} K_1(x, y) x^{-\alpha r} > 0$$

and

$$\frac{\partial^\alpha}{\partial y^\alpha} K_1(x, y) y^{-\alpha r} < 0 \quad \text{and} \quad \frac{\partial^{2\alpha}}{\partial y^{2\alpha}} K_1(x, y) y^{-\alpha r} > 0,$$

for $r > 0$.

In what follows we suppose that

$$(2.23) \quad A_1 = \frac{2-s}{2q}, \quad A_2 = \frac{2-s}{2p}.$$

Then, based on equation (2.23), the constant L^* from Theorem B becomes

$$\begin{aligned} L^* &= \Gamma(1+\alpha)k(pA_2) = \Gamma(1+\alpha)k\left(1 - \frac{s}{2}\right) \\ &= \Gamma(1+\alpha) \frac{1}{\Gamma(1+\alpha)} \int_0^{+\infty} \frac{u^{-\alpha-\frac{\alpha s}{2}}}{(1+u)^{\alpha s}} (du)^\alpha = \Gamma(1+\alpha) B_\alpha\left(\frac{s}{2}, \frac{s}{2}\right). \end{aligned}$$

Now, from Theorem B, we get the following result.

Corollary 2.1. *Let $1/p + 1/q = 1$, $p > 1$, $0 < s < 2$, and $(a_m)_{m \in \mathbb{N}}$ and $(b_n)_{n \in \mathbb{N}}$ be non-negative real sequences. Then the following inequalities hold and are equivalent*

$$\begin{aligned} \sum_{m=1}^{+\infty} \sum_{n=1}^{+\infty} \frac{a_m^\alpha b_n^\alpha}{(m+n)^{\alpha s}} &\leq \Gamma(1+\alpha) B_\alpha\left(\frac{s}{2}, \frac{s}{2}\right) \\ &\times \left(\sum_{m=1}^{+\infty} m^{\alpha p(1-\frac{s}{2})-\alpha} a_m^{\alpha p} \right)^{\frac{1}{p}} \left(\sum_{n=1}^{+\infty} n^{\alpha q(1-\frac{s}{2})-\alpha} b_n^{\alpha q} \right)^{\frac{1}{q}} \end{aligned}$$

and

$$\left(\sum_{n=1}^{+\infty} n^{\frac{\alpha p s}{2}-\alpha} \left(\sum_{m=1}^{+\infty} \frac{a_m^\alpha}{(m+n)^{\alpha s}} \right)^p \right)^{\frac{1}{p}} \leq \Gamma(1+\alpha) B_\alpha\left(\frac{s}{2}, \frac{s}{2}\right) \left(\sum_{m=1}^{+\infty} m^{\alpha p(1-\frac{s}{2})-\alpha} a_m^{\alpha p} \right)^{\frac{1}{p}},$$

where the constant $\Gamma(1+\alpha) B_\alpha(s/2, s/2)$ is the best possible.

3. CONCLUSION

In this paper, we have firstly obtained a fractal Hölder's inequality and some related inequalities. According to the basic results, some new discrete local fractional Hilbert-type inequalities have been investigated. At the same time, some new fractional Hilbert-type inequalities with homogeneous kernels have been given.

4. ACKNOWLEDGMENTS

The authors would like to thank the editors and the anonymous referees for their constructive comments and suggestions which led to improvement of the original manuscript. This work was supported by the High-level Talent Fund Project of Sanmenxia Polytechnic under Grant No. SZYGCCRC-2021-009.

REFERENCES

- [1] G. H. Hardy, J. E. Littlewood and G. Pólya, *Inequalities*, Second Edition, Cambridge University Press, Cambridge, 1967.
- [2] F. Jarad, T. Abdeljawad and J. Alzabut, *Generalized fractional derivatives generated by a class of local proportional derivatives*, European Physical Journal Special Topics **226** (2017), 3457–3471. <https://doi.org/10.1140/epjst/e2018-00021-7>
- [3] J. V. C. Sousa, D. S. Oliveira and E. C. Oliveira, *Grüss-type inequalities by means of generalized fractional integrals*, Bull. Braz. Math. Soc. (N.S.) **50** (2019), 1029–1047. <https://doi.org/10.1007/s00574-019-00138-z>
- [4] W. Yang, *Certain new Chebyshev and Grüss-type inequalities for unified fractional integral operators via an extended generalized Mittag-Leffler function*, Fractal and Fractional **6**(4) (2022), Paper ID 182, 27 pages. <https://doi.org/10.3390/fractalfract6040182>
- [5] W. Yang, *Certain new weighted Young and Pólya-Szegő-type inequalities for unified fractional integral operators via an extended generalized Mittag-Leffler function with applications*, Fractals **30**(6) (2022), Paper ID 2250106, 37 pages. <https://doi.org/10.1142/S0218348X22501067>
- [6] X. J. Yang, *Local Fractional Functional Analysis and its Applications*, Asian Academic Publisher Limited, Hong Kong, 2011.
- [7] X. J. Yang, *Advanced Local Fractional Calculus and its Applications*, World Science Publisher, New York, 2012.
- [8] X. J. Yang, D. Baleanu and H. M. Srivastava, *Advanced analysis of local fractional calculus applied to the rice theory in fractal fracture mechanics*, in: J. Singh, H. Dutta, D. Kumar, D. Baleanu and J. Hristov (Eds.), *Methods of Mathematical Modelling and Computation for Complex Systems*, Studies in Systems, Decision and Control, Vol. 373. Springer, Cham, 2022.
- [9] M. Krnić and P. Vuković, *Multidimensional Hilbert-type inequalities obtained via local fractional calculus*, Acta Appl. Math. **169**(1) (2020), 667–680. <https://doi.org/10.1007/s10440-020-00317-x>
- [10] G. Jumarie, *Fractional Euler's integral of first and second kinds. Application to fractional Hermite's polynomials and to probability density of fractional order*, J. Appl. Math. Inform. **28** (2010), 257–273.
- [11] Q. Liu, *A Hilbert-type fractional integral inequality with the kernel of Mittag-Leffler function and its applications*, Math. Inequal. Appl. **21**(3) (2018), 729–737. <https://dx.doi.org/10.7153/mia-2018-21-52>
- [12] Q. Liu and D. Chen, *A Hilbert-type integral inequality on the fractal spaces*, Integral Transforms Spec. Funct. **28**(10) (2017), 772–780. <https://doi.org/10.1080/10652469.2017.1359588>
- [13] T. Batbold, M. Krnić and P. Vuković, *A unified approach to fractal Hilbert-type inequalities*, J. Inequal. Appl. **2019** (2019), Paper ID 117, 13 pages. <https://doi.org/10.1186/s13660-019-2076-9>
- [14] Y. Li, S. Rashid, Z. Hammouch, D. Baleanu and Y. Chu, *New Newton's type estimates pertaining to local fractional integral via generalized p -convexity with applications*, Fractals **29**(5) (2021), Paper ID 2140018, 20 pages. <https://doi.org/10.1142/S0218348X21400181>

- [15] M. Z. Sarikaya, T. Tunc and H. Budak, *On generalized some integral inequalities for local fractional integrals*, Appl. Math. Comput. **276** (2016), 316–323. <https://doi.org/10.1016/j.amc.2015.11.096>
- [16] S. Iftikhar, P. Kumam and S. Erden, *Newton's-type integral inequalities via local fractional integrals*, Fractals **28**(3) (2020), Paper ID 2050037, 13 pages. <https://doi.org/10.1142/S0218348X20500371>
- [17] A. Akkurt, M. Z. Sarikaya, H. Budak and H. Yildirim, *Generalized Ostrowski type integral inequalities involving generalized moments via local fractional integrals*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. **111**(3) (2017), 797–807. <https://doi.org/10.1007/s13398-016-0336-9>
- [18] M. Z. Sarikaya and H. Budak, *Generalized Ostrowski type inequalities for local fractional integrals*, Proc. Amer. Math. Soc. **145**(4) (2017), 1527–1538. <http://dx.doi.org/10.1090/proc/13488>
- [19] W. Sun, *Hermite-Hadamard type local fractional integral inequalities with Mittag-Leffler kernel for generalized preinvex functions*, Fractals **29**(8) (2021), Paper ID 2150253, 13 pages. <https://doi.org/10.1142/S0218348X21502534>
- [20] J. Choi, E. Set and M. Tomar, *Certain generalized Ostrowski type inequalities for local fractional integrals*, Commun. Korean Math. Soc. **32**(3) (2017), 601–617. <https://doi.org/10.4134/CKMS.c160145>
- [21] H. Mo, X. Sui and D. Yu, *Generalized convex functions on fractal sets and two related inequalities*, Abstr. Appl. Anal. **2014** (2014), Paper ID 636751, 7 pages. <https://doi.org/10.1155/2014/254737>
- [22] H. Budak, M. Z. Sarikaya and H. Yildirim, *New inequalities for local fractional integrals*, Iran. J. Sci. Technol. Trans. A Sci. **41**(4) (2017), 1039–1046. <https://doi.org/10.1007/s40995-017-0315-9>

¹UNIVERSITY OF ZAGREB,
 FACULTY OF TEACHER EDUCATION,
 SAVSKA CESTA 77, 10000 ZAGREB, CROATIA
Email address: predrag.vukovic@ufzg.hr

²NORMAL SCHOOL,
 SANMENXIA POLYTECHNIC,
 SANMENXIA 472000, CHINA
Email address: yangwg8088@163.com