

ON CENTRALLY-EXTENDED GENERALIZED JORDAN  
\*-DERIVATIONS IN RINGS

BHARAT BHUSHAN<sup>1</sup>, GURNINDER S. SANDHU<sup>2</sup>, SHAKIR ALI<sup>3</sup>,  
AND BRUNO LEONARDO MACEDO FERREIRA<sup>4,5,6</sup>

ABSTRACT. Let  $R$  be an associative ring with an involution  $'*$ '. In this article, we introduce the notions of *centrally-extended generalized Jordan  $*$ -derivation*, *centrally extended Jordan left  $*$ -centralizer* and characterize these mappings in involutive prime rings.

1. INTRODUCTION

Throughout this study,  $R$  is an associative ring with center  $Z(R)$ .  $R$  is called *prime*, if for any  $a, b \in R$ ,  $aRb = (0)$  implies either  $a = 0$  or  $b = 0$ ; and it is called *semiprime*, if  $aRa = (0)$ , implies  $a = 0$ . Clearly, every prime ring is semiprime ring but the converse need not be true, for instance  $\mathbb{Z} \times \mathbb{Z}$ . The symmetric ring of quotients of  $R$  is denoted by  $Q_s$  with center  $C$ , which is known as the extended centroid of  $R$ ; clearly  $R \subseteq Q_s$  and  $Z(R) \subseteq C$ . It is well-known that if  $R$  is prime then  $Q_s$  is prime and  $C$  is a field. The central closure of  $R$  is denoted by  $A(= RC + C)$ ; for more details of these objects, we refer the reader to [6]. For any  $x, y \in R$ , the commutator (resp. anti-commutator) of  $x, y$  is defined as  $[x, y] = xy - yx$  (resp.  $x \circ y = xy + yx$ ). It is established knowledge that  $R$  satisfies  $s_4$  (the standard identity in four noncommuting variables), if for all  $x_1, x_2, x_3, x_4 \in R$ , the equation

$$s_4(x_1, x_2, x_3, x_4) = \sum_{\sigma \in S_4} (-1)^\sigma x_{\sigma(1)} x_{\sigma(2)} x_{\sigma(3)} x_{\sigma(4)} = 0,$$

---

*Key words and phrases.* Prime ring, involution, centrally-extended Jordan  $*$ -derivation, centrally-extended generalized Jordan  $*$ -derivation, centrally-extended Jordan left  $*$ -centralizer.

2020 *Mathematics Subject Classification.* Primary: 16W10. Secondary: 16N60, 16W25.  
<https://doi.org/10.46793/KgJMat2608.1329B>

*Received:* September 17, 2024.

*Accepted:* December 16, 2024.

where  $S_4$  is the symmetric group of degree 4 and  $(-1)^\sigma$  is the sign of permutation  $\sigma \in S_4$ . For some interesting equivalent forms of  $s_4$ , one can refer [11, Lemma 1].

For any  $n \in \mathbb{Z}^+$ ,  $R$  is called an  $n$ -torsion free if for any  $x \in R$ ,  $nx = 0$  implies  $x = 0$ . An anti-automorphism  $'*$  of  $R$  is called *involution* if  $(x^*)^* = x$  for all  $x \in R$ . A ring equipped with an involution  $'*$  is called *\*-ring* or *ring with involution* or *involutive ring*. An element  $x$  in a \*-ring  $R$  is called *symmetric* if  $x^* = x$ , and it is called *skew-symmetric* if  $x^* = -x$ . The set of symmetric and skew symmetric elements of a ring  $R$  is denoted by  $H(R)$  and  $S(R)$ , respectively. Moreover, if  $R$  is a prime ring endowed with the involution  $'*$ , then  $'*$  can be uniquely extended to  $Q_s(R)$  (see [15, page 4]).

Let  $R$  be a \*-ring, an additive mapping  $d : R \rightarrow R$  is called *\*-derivation* if  $d(xy) = d(x)y^* + xd(y)$  for all  $x, y \in R$  and it is called *Jordan \*-derivation* if  $d(x^2) = d(x)x^* + xd(x)$  for all  $x \in R$ . These notions are first mentioned in [12]. Note that the mapping  $x \mapsto xa - ax^*$ , where  $a$  is a fixed element of  $R$ , is an example of Jordan \*-derivation, called *inner Jordan \*-derivation*. Moreover, if  $a \in Q_s$ , then such a map is called *X-inner Jordan \*-derivation*. The study of Jordan \*-derivations has been originated from the problem of representability of quadratic forms by bilinear forms (see [28, 30]). Thereafter, some significant studies have taken place on the structure of Jordan \*-derivations in rings (see [5, 14, 16, 21]).

In [2], Ali introduced the notion of *generalized \*-derivation*, which is a self-map  $F$  of  $R$  associated with a \*-derivation  $d$  satisfying  $F(xy) = F(x)y^* + xd(y)$  for all  $x, y \in R$ . In addition to this, a self-map  $F$  of  $R$  is called a *generalized Jordan \*-derivation* associated with a Jordan \*-derivation  $d$  if  $F(x^2) = F(x)x^* + xd(x)$  for all  $x \in R$  (see [16], [19]). For any fixed  $a, b \in R$ , a map  $x \mapsto ax^* + xb$  is a basic example of generalized Jordan \*-derivation, which is called *generalized inner Jordan \*-derivation*. Further, if  $a, b$  comes from  $Q_s$ , then this map is called *generalized X-inner Jordan \*-derivation*. There has been an ongoing interest in the investigation of such mappings, for more details we refer the reader to [1, 4, 5, 16, 19] and references therein.

A mapping  $f$  of  $R$  is called *centralizing* (resp. *commuting*) on a subset  $S$  of  $R$  if  $[f(x), x] \in Z(R)$  (resp.  $[f(x), x] = 0$ ) for all  $x \in S$ . To our best knowledge, Divinsky [17] initiated the study of commuting and centralizing mappings in rings by proving a classical result which states that a simple Artinian ring is commutative if it admits a commuting nontrivial automorphism. Since then, many significant results on commuting and centralizing mappings have been established by Posner [27], Mayne [25], Bell and Martindale [8], Brešar [10]. Moreover, let  $R$  be a \*-ring, a mapping  $f : R \rightarrow R$  is called *\*-centralizing* (*\*-commuting*) on a subset  $S$  of  $R$  if  $[f(x), x^*] \in Z(R)$  (resp.  $[f(x), x^*] = 0$ ) for all  $x \in S$  (see [3]).

Bell and Daif [7] introduced *centrally-extended derivations* and discussed their existence. Accordingly, a mapping  $d : R \rightarrow R$  is called *centrally-extended derivation* if  $d(x + y) - d(x) - d(y) \in Z(R)$  and  $d(xy) - d(x)y - xd(y) \in Z(R)$  for all  $x, y \in R$ . Motivated by this, El-Deken and Nabel [31] introduced that a mapping  $d$  is called *centrally-extended \*-derivation* if  $d(x + y) - d(x) - d(y) \in Z(R)$  and  $d(xy) -$

$d(x)y^* - xd(y) \in Z(R)$  for all  $x, y \in R$ . Furthermore, a mapping  $F$  is called *centrally-extended generalized \*-derivation* associated with a centrally-extended \*-derivation  $d$  if  $F(x+y) - F(x) - F(y) \in Z(R)$  and  $F(xy) - F(x)y^* - xd(y) \in Z(R)$  for all  $x, y \in R$ . In a recent paper [9], we introduced and studied the concept of *centrally-extended Jordan \*-derivation*, which is a mapping  $d : R \rightarrow R$  such that  $d(x+y) - d(x) - d(y) \in Z(R)$  and  $d(x \circ y) - d(x)y^* - d(y)x^* - xd(y) - yd(x) \in Z(R)$  for all  $x, y \in R$ . Nowadays, centrally-extended mappings are getting attention of many researchers, consequently there has been rising literature on these mappings in rings under different settings, for instance, see [7, 9, 18, 26, 31–34].

In this article, we shall introduce centrally-extended generalized Jordan \*-derivation in rings and discuss their existence in noncommutative prime ring under suitable torsion conditions. We also investigate some specific functional identities involving centrally-extended generalized Jordan \*-derivations. Precisely, in Section 4 of this article, we prove a structural result on centrally-extended generalized Jordan \*-derivations which plays a key role in Sections 5 and 6, where we study centralizing and hypercommuting conditions involving centrally-extended generalized Jordan \*-derivations, respectively.

## 2. PRELIMINARIES

The following lemmas constitute a set of results that will be instrumental in the development of the paper.

**Lemma 2.1** (Brauer's Trick). *Let  $G$  be a group and  $H_1, H_2$  be subgroups of  $G$ . If  $G = H_1 \cup H_2$ , then either  $G = H_1$  or  $G = H_2$ .*

**Lemma 2.2.** *If  $R$  is a prime ring, then  $Z(R)$  has no proper zero divisor.*

**Lemma 2.3.** *Let  $R$  is a prime for any  $a \in Z(R)$ . If there exists  $b \in R$  such that  $ab \in Z(R)$ , then either  $a = 0$  or  $b \in Z(R)$ .*

**Lemma 2.4.** ([1, Proposition 2.3]). *Let  $R$  be a 2-torsion free semiprime ring with involution  $'*$ . If  $f : R \rightarrow R$  is an additive map such that  $f(x^2) = f(x)x^*$  for all  $x \in R$ , then there exists  $q \in Q_r(R)$  such that  $f(x) = qx^*$  for all  $x \in R$ .*

**Lemma 2.5.** ([3, Lemma 2.2]). *Let  $R$  be a 2-torsion free semiprime ring with involution  $'*$ . If an additive mapping  $f$  of  $R$  into itself such that  $[f(x), x^*] \in Z(R)$  for all  $x \in R$ , then  $[f(x), x^*] = 0$  for all  $x \in R$ .*

**Lemma 2.6.** ([6, Theorem 6.4.6]). *Let  $R$  be a prime ring with extended centroid  $C$ , anti-automorphism  $g$  and maximal right ring of quotients  $Q_{mr}(R) = Q$ . If  $0 \neq \phi = \phi(x_1, \dots, x_n, g(x_1), \dots, g(x_n)) \in Q_C \langle X \cup g(X) \rangle$  is a  $g$ -identity on  $K$  ideal of  $R$ , then  $\phi$  is a  $g$ -identity on  $Q_s = Q_s(R)$ .*

**Lemma 2.7.** ([9, Theorem 4.6]). *Let  $R$  be a 2-torsion free noncommutative prime ring. If  $R$  admits a non-zero centrally-extended Jordan derivation  $\mathfrak{d} : R \rightarrow R$  such*

that  $[\mathfrak{d}(x), x] \in Z(R)$  for all  $x \in R$ , then either  $\mathfrak{d} = 0$  or  $R$  is an order in a central simple algebra of dimension at most 4 over its center.

**Lemma 2.8.** ([9, Theorem 4.7]). *Let  $R$  be a 2-torsion free noncommutative prime ring with involution  $*$ . If  $R$  admits a non-zero centrally-extended Jordan derivation  $\mathfrak{d} : R \rightarrow R$  such that  $[\mathfrak{d}(x), x^*] \in Z(R)$  for all  $x \in R$ , then either  $\mathfrak{d} = 0$  or  $R$  is an order in a central simple algebra of dimension at most 4 over its center.*

**Lemma 2.9.** ([10, Proposition 3.1]). *Let  $R$  be a 2-torsion free semiprime ring and  $U$  be a Jordan subring of  $R$ . If an additive mapping  $F$  of  $R$  into itself is centralizing on  $U$ , then  $F$  is commuting on  $U$ .*

**Lemma 2.10.** ([10, Theorem 3.2]). *Let  $R$  be a prime ring. If an additive mapping  $F : R \rightarrow R$  is commuting on  $R$ , then there exists  $\lambda \in C$  and an additive  $\xi : R \rightarrow C$ , such that  $F(x) = \lambda x + \xi(x)$  for all  $x \in R$ .*

**Lemma 2.11.** ([19, Theorem 2.2]). *Let  $R$  be a 2-torsion free prime ring with involution  $'*$ . Let  $F : R \rightarrow R$  be a generalized Jordan  $*$ -derivation associated with a Jordan  $*$ -derivation  $d$ . Then,  $F$  is of the form  $F(x) = qx^* + d(x)$  for all  $x \in R$  and some  $q \in Q_s(R)$ .*

**Lemma 2.12.** ([20, Theorem 1]). *Let  $R$  be a prime ring with involution  $'*$  and center  $Z(R)$ . If  $d$  is a non-zero derivation such that  $[d(h), h] \in Z(R)$  for all  $h \in H(R)$ , then  $R$  satisfies  $s_4$ .*

**Lemma 2.13.** ([20, Theorem 3]). *Let  $R$  be a prime ring with involution  $'*$  and center  $Z(R)$ . If  $n$  be a fixed natural number such that  $x^n \in Z(R)$  for all  $x \in H(R)$ , then  $R$  satisfies  $s_4$  identity.*

**Lemma 2.14.** ([20, Theorem 6]). *Let  $R$  be a prime ring with involution  $'*$  and center  $Z(R)$ . If  $d$  is a non-zero derivation on  $R$  such that  $d(x)x + xd(x) \in Z(R)$  for all  $x \in H(R)$ , then  $R$  satisfies  $s_4$  identity.*

**Lemma 2.15.** ([20, Theorem 7]). *Let  $R$  be a prime ring with involution  $'*$  and center  $Z(R)$ . If  $d$  is a non-zero derivation on  $R$  such that  $d(x)x + xd(x) \in Z(R)$  for all  $x \in S(R)$ , then  $R$  satisfies  $s_4$  identity.*

**Lemma 2.16.** [21, Theorem 1.3]. *Let  $R$  be a 2-torsion free noncommutative prime ring with involution  $'*$ , then any Jordan  $*$ -derivation on  $R$  is  $X$ -inner.*

**Lemma 2.17.** *Let  $R$  be a 2-torsion free prime ring. If  $q_1 \in Q_s(R)$  such that  $[q_1, h] \in C$  for all  $h \in H(R)$ , then  $R$  satisfy  $s_4$  identity or  $q_1 \in C$ .*

*Proof.* Let us consider

$$(2.1) \quad [q_1, h] \in C, \quad \text{for all } h \in H(R).$$

Replacing  $h$  by  $h^2$ , where  $h \in H(R)$ , we obtain  $[q_1, h]h + h[q_1, h] \in C$ , i.e.,  $d(h)h + hd(h) \in C$  for all  $h \in H(R)$ , where  $d(x) = [q_1, x]$ . If  $d \neq 0$ , we have the result by Lemma 2.14. If  $d = 0$ , then we conclude  $q_1 \in C$ , as desired.  $\square$

**Lemma 2.18.** *Let  $R$  be a 2-torsion free prime ring with involution. If  $[h, k] = 0$  for all  $h \in H(R)$ ,  $k \in S(R)$ , then  $R$  satisfy  $s_4$  identity.*

*Proof.* Suppose that  $R$  does not satisfy  $s_4$  identity. By hypothesis, we have  $[h, k] = 0 \in C$  for all  $h \in H(R)$  and  $k \in S(R)$ . In view of Lemma 2.17, it follows that either  $R$  satisfies  $s_4$  or  $k \in Z(R)$  for all  $k \in S(R)$ . Under the given hypothesis, we left with  $S(R) \subseteq Z(R)$ . Clearly  $h \circ k \in S(R)$  for all  $h \in H(R)$  and  $k \in S(R)$ , therefore we have  $[h \circ k, k] = 0$ , i.e.,  $[h, k]k + k[h, k] = 0$  for all  $h \in H(R)$  and  $k \in S(R)$ . For a fixed  $h \in H(R)$ , we have  $d(k)k + kd(k) = 0$  for all  $k \in S(R)$ , where  $d(x) = [x, h]$  for all  $x \in R$ . If  $d \neq 0$ , then we have the result by Lemma 2.15. In case  $d = 0$ , we conclude  $H(R) \subseteq Z(R)$  and hence by Lemma 2.13, we have a contradiction. Hence,  $R$  must satisfies  $s_4$  identity. □

### 3. DEFINITIONS AND EXAMPLES

We begin our discussions with the definition of centrally-extended generalized Jordan \*-derivations of rings with involution.

**Definition 3.1.** Let  $R$  be a ring with involution  $'*$ '. A mapping  $F : R \rightarrow R$  is called *centrally-extended generalized Jordan \*-derivation* associated with an centrally-extended Jordan \*-derivation  $d$ , if

- (A)  $F(x + y) - F(x) - F(y) \in Z(R)$ ,
- (B)  $F(x \circ y) - F(x)y^* - F(y)x^* - xd(y) - yd(x) \in Z(R)$ ,

for all  $x, y \in R$ .

*Remark 3.1.* If  $R$  is 2-torsion free noncommutative prime ring with involution  $'*$ ', then for an additive mapping  $F$  to be a centrally-extended generalized Jordan \*-derivation, it is sufficient to satisfy the condition  $F(x^2) - F(x)x^* - xd(x) \in Z(R)$  for all  $x \in R$ .

*Example 3.1.* We now show the existence of centrally-extended generalized Jordan \*-derivations in certain rings.

(I) Let  $\mathbb{Z}$  be the ring of integers and  $R = \left\{ \begin{pmatrix} x & y \\ z & t \end{pmatrix} : x, y, z, t \in \mathbb{Z} \right\}$ , a noncommutative prime ring. Then the mapping  $*$  :  $R \rightarrow R$  such that  $\begin{pmatrix} x & y \\ z & t \end{pmatrix}^* = \begin{pmatrix} t & -y \\ -z & x \end{pmatrix}$ ,  $F : R \rightarrow R$  such that  $F \begin{pmatrix} x & y \\ z & t \end{pmatrix} = \begin{pmatrix} x - t & y \\ z & 0 \end{pmatrix}$  with associated mapping  $d : R \rightarrow R$  defined as  $d \begin{pmatrix} x & y \\ z & t \end{pmatrix} = \begin{pmatrix} x & 0 \\ 0 & x \end{pmatrix}$ . One can easily verify that  $F$  is a centrally-extended generalized Jordan \*-derivation with associated centrally-extended Jordan \*-derivation  $d$ .

(II) Let  $R$  be a ring defined as  $R := \mathfrak{R} \times \mathbb{Z}$ , where  $\mathfrak{R} = \left\{ \begin{pmatrix} x & y \\ 0 & z \end{pmatrix} : x, y, z \in \mathbb{Z}_2 \right\}$ . For any  $X = \begin{pmatrix} x & y \\ 0 & z \end{pmatrix} \in \mathfrak{R}$ , let us define  $X^* = \begin{pmatrix} z & y \\ 0 & x \end{pmatrix}$  and hence  $(X, k)^* = (X^*, k)$ , which is an involution on  $R$ . Define  $F : R \rightarrow R$  such that  $F \left( \begin{pmatrix} x & y \\ 0 & z \end{pmatrix}, k \right) = \left( \begin{pmatrix} 0 & x \\ 0 & x \end{pmatrix}, 1 \right)$  with associated mapping  $d : R \rightarrow R$  defined as  $d \left( \begin{pmatrix} x & y \\ 0 & z \end{pmatrix}, k \right) = \left( \begin{pmatrix} z & y \\ 0 & x \end{pmatrix}, 1 \right)$ . We observe that  $F$  is a centrally-extended generalized Jordan  $*$ -derivation with an associated centrally-extended Jordan  $*$ -derivation.

*Remark 3.2.* It can be seen from the above example that every centrally-extended generalized Jordan  $*$ -derivation is a centrally-extended Jordan  $*$ -derivation, but the converse does not necessarily hold in general.

**Definition 3.2.** Let  $R$  be a ring with involution  $'*$ '. A mapping  $T : R \rightarrow R$  is called *centrally-extended Jordan  $*$ -left centralizer* (resp. *centrally-extended Jordan  $*$ -right centralizer*) if

- (A)  $T(x + y) - T(x) - T(y) \in Z(R)$ ,
- (B)  $T(x^2) - T(x)x^* \in Z(R)$  (resp.  $T(x^2) - x^*T(x) \in Z(R)$ ),

for all  $x, y \in R$ . Moreover,  $T$  is called *centrally-extended Jordan  $*$ -centralizer* if it is both a centrally-extended Jordan  $*$ -left centralizer and a centrally-extended Jordan  $*$ -right centralizer.

*Example 3.2.* Let  $R$  be a ring defined as  $R := M_2(\mathbb{R}) \times \mathbb{C}$ , where  $M_2(\mathbb{R})$  denotes the ring of  $2 \times 2$  matrices over real numbers. For any  $r = (x, z_1), s = (y, z_2) \in R$ , we define  $r^* = (x^*, \bar{z}_1)$ , where  $x^*$  is defined as in Example 3.1 (I) and  $\bar{z}_1$  is the complex conjugate of  $z_1$ . Define a mapping  $T : R \rightarrow R$  such that  $T(x, z) = (0, 1)$  for all  $(x, z) \in R$ . Then it can be easily verified that  $T$  is a centrally-extended Jordan  $*$ -left centralizer of  $R$ .

#### 4. AUXILIARY RESULTS

In this section, we shall mainly prove the following theorem which is crucial for the results proved in the subsequent sections.

**Theorem 4.1.** *Let  $R$  be a 2-torsion free noncommutative prime ring with involution  $'*$ ' and  $F$  be a centrally-extended generalized Jordan  $*$ -derivation of  $R$  associated with a centrally-extended Jordan  $*$ -derivation  $\mathfrak{d}$ . Then, there exists  $q \in Q_s(R)$  such that  $F(x) = qx^* + \mathfrak{d}(x)$  for all  $x \in R$ .*

After proving few more facts in this regard, we shall return to the proof of Theorem 4.1.

**Proposition 4.1.** *Let  $R$  be a 2-torsion free prime ring with involution  $'*'$  and  $d : R \rightarrow R$  a Jordan  $*$ -derivation. Then,  $d$  can be uniquely extended to  $A$ , unless  $R$  satisfies  $s_4$ .*

*Proof.* Suppose that  $R$  does not satisfy  $s_4$  identity. By Lemma 2.16, there exists  $a \in Q_s(R)$  such that  $d(x) = xa - ax^*$  for all  $x \in R$ . Let us define  $\bar{d}(q) = qa - aq^*$  for all  $q \in A$ . Clearly,  $\bar{d}$  is a well-defined map and it is an extension of  $d$ . Now, we claim that the extension of  $d$  is unique.

Let  $D$  be also an extension of  $d$ . Since  $A$  is a noncommutative 2-torsion free prime ring,  $D$  is  $X$ -inner on  $A$ , i.e., there exists  $b \in A$  such that  $D(u) = ub - bu^*$  for all  $u \in A$ . As  $D$  is an extension of  $d$ , so we have

$$(4.1) \quad xb - bx^* = xa - ax^*, \quad \text{for all } x \in R.$$

In particular by taking  $h$  for  $x$  in (4.1), where  $h \in H(R)$ , we obtain  $[h, b - a] = 0$  for all  $h \in H(R)$ . Using Lemma 2.17, we have  $b - a \in C$ . That means there exists  $c \in C$  such that  $b = a + c$ . From (4.1), we have  $xc - cx^* = 0$  for all  $x \in R$ . Replacing  $x$  by  $k$  in the last expression, where  $k \in S(R)$ , we get  $2ck = 0$  for all  $k \in S(R)$ . If  $c \neq 0$ , then by using Lemma 2.2, we have  $S(R) = (0)$ , thence  $H(R) = R$ ; which implies that for any  $x, y \in R$ , we have  $xy = (xy)^* = y^*x^* = yx$ , which is a contradiction. In case  $c = 0$ , we have  $a = b$ . It proves our claim. □

**Proposition 4.2.** *Let  $R$  be a 2-torsion free prime ring with involution  $'*'$  and  $F : R \rightarrow R$  a generalized Jordan  $*$ -derivation associated with  $d$  a Jordan  $*$ -derivation. Then,  $F$  can be uniquely extended to  $A$ , unless  $R$  satisfies  $s_4$ .*

*Proof.* By Lemma 2.11, there exists  $a \in Q_s(R)$  such that  $F(x) = ax^* + d(x)$  for all  $x \in R$ . We define  $\tilde{F}(u) = au^* + d(u)$  for all  $u \in A$ . By Proposition 4.1,  $\tilde{F}$  is a well defined map and also it is an extension of  $F$ . Now, we will show that extension of  $F$  is unique. Let  $G$  be also extension of  $F$ . Using Lemma 2.11, there exists  $b \in Q_s$  and  $g$  a Jordan  $*$ -derivation on  $A$  such that  $G(u) = bu^* + g(u)$  for all  $u \in A$ . Since,  $G$  is an extension of  $F$ . Therefore

$$(4.2) \quad ax^* + d(x) = bx^* + g(x), \quad \text{for all } x \in R.$$

By Lemma 2.16, there exists  $c, q \in Q_s$  such that  $d(x) = xc - cx^*$  and  $g(x) = xq - qx^*$  for all  $x \in R$ . From (4.2), we have  $(a - c)x^* + xc = (b - q)x^* + xq$  for all  $x \in R$ . As the preceding equation is a  $g$ -identity on  $R$ , application of Lemma 2.6 yields  $(a - c)x^* + xc = (b - q)x^* + xq$  for all  $x \in Q_s(R)$ . Replacing  $x$  by 1, we obtain  $a = b$ . From (4.2), we find  $d(x) = g(x)$  for all  $x \in R$ . In fact, Proposition 4.1 gives  $d(x) = g(x)$  for all  $x \in A$ . It completes the proof. □

For the sake of brevity, we omit the proof of the following result, as it follows proceeding along the same lines as the proof of Lemma 4.4 of [9], with insignificant variations.

**Proposition 4.3.** *Let  $R$  be a ring with involution  $*$  and with no non-zero central ideal. If  $F$  is a centrally-extended generalized Jordan  $*$ -derivation associated with centrally-extended Jordan  $*$ -derivation  $\mathfrak{d}$  of  $R$ , then  $F$  is additive.*

**Corollary 4.1.** *Let  $R$  be a noncommutative prime ring with involution  $'*$ '. If  $F$  is a centrally-extended generalized Jordan  $*$ -derivation of  $R$  associated with centrally-extended Jordan  $*$ -derivation  $\mathfrak{d}$ , then  $F$  is additive.*

**Proposition 4.4.** *Let  $R$  be a 2-torsion free noncommutative prime ring with involution  $'*$ ' and  $T : R \rightarrow R$  a centrally-extended Jordan  $*$ -left centralizer. Then, there exists  $q \in Q_s(R)$  such that  $T(x) = qx^*$  for all  $x \in R$ .*

*Proof.* Note that  $T$  is additive by Proposition 4.1. If  $Z(R) = (0)$ , then  $T(x^2) = T(x)x^*$  for all  $x \in R$ . Thus, we conclude the desired result by Lemma 2.4. In case  $Z(R) \neq (0)$ , we first claim that there exists  $0 \neq z \in Z(R)$  such that  $z^* = z$ .

Let us suppose that  $0 \neq z_c \in Z(R)$ . If  $z_c^* = z_c$ , then we are done. If  $z_c^* \neq z_c$ , then we take  $z_1 = z_c + z_c^*$ ; and observe that  $z_1 = z_1^*$ . Therefore, we can say that there exists  $0 \neq z \in Z(R)$  such that  $z^* = z$ .

By the assumption, we have

$$(4.3) \quad T(x^2) - T(x)x^* \in Z(R), \quad \text{for all } x \in R.$$

Polarizing (4.3), we get

$$(4.4) \quad T(xy + yx) - T(x)y^* - T(y)x^* \in Z(R), \quad \text{for all } x, y \in R.$$

Replacing  $y$  by  $z^2$ , where  $z \in H(R) \cap Z(R)$  in (4.4) to get

$$T(xz^2 + z^2x) - T(x)z^2 - T(z^2)x^* \in Z(R), \quad \text{for all } x \in R.$$

It implies

$$T(xz^2 + z^2x) - T(x)z^2 - T(z)zx^* - c_1x^* \in Z(R), \quad \text{for all } x \in R,$$

where  $c_1$  is the corresponding central element. It implies

$$(4.5) \quad T(4xz^2) - 2T(x)z^2 - 2T(z)zx^* - 2c_1x^* \in Z(R).$$

Also,  $4xz^2 = z(xz + zx) + (xz + zx)z$ . From (4.4), we have

$$T(z(xz + zx) + (xz + zx)z) - T(z)(xz + zx)^* - T(xz + zx)z \in Z(R).$$

Again using (4.4) in the last summand of the above relation, we get

$$(4.6) \quad T(4xz^2) - T(z)(x^*z + zx^*) - (T(x)z + T(z)x^* + c_2)z \in Z(R),$$

where  $c_2$  is the corresponding central element. Comparing (4.5) and (4.6) to obtain

$$T(x)z^2 - T(z)x^*z - c_1x^* \in Z(R).$$

It implies

$$([T(x)z, x^*] - [T(z)x^*, x^*])z = 0.$$

Application of Lemma 2.2 yields

$$(4.7) \quad [T(x)z - T(z)x^*, x^*] = 0.$$

Define  $\mathfrak{G}(x) = T(x)z - T(z)x^*$  for all  $x \in R$ . From (4.7), we find  $\mathfrak{G}$  is an additive \*-commuting map. Applying involution in (4.7) and using Lemma 2.10 in order to get  $\mathfrak{G}(x)^* = \lambda x + \sigma(x)$  for all  $x \in R$ , for some  $\lambda \in C$  and  $\sigma : R \rightarrow C$ . It implies  $\mathfrak{G}(x) = T(x)z - T(z)x^* = \lambda^*x^* + \sigma(x)^*$ . Therefore,  $T(x)z = (T(z) + \lambda^*)x^* + \sigma(x)^*$  for all  $x \in R$ . It can also be written as  $T(x) = z^{-1}(T(z) + \lambda^*)x^* + \sigma(x)^*z^{-1}$ . Thus,  $T(x) = qx^* + \sigma'(x)$  for all  $x \in R$  where  $q = z^{-1}(T(z) + \lambda^*)$  and  $\sigma'(x) = z^{-1}\sigma(x)^*$ . From (4.3), we have  $q(x^2)^* + \sigma'(x^2) - (qx^* + \sigma'(x))x^* \in Z(R)$  for all  $x \in R$ . It implies

$$(4.8) \quad x^*\sigma'(x) \in C, \quad \text{for all } x \in R.$$

For any fixed  $x \in R$ , using Lemma 2.3 in (4.8), we have either  $x \in Z(R)$  or  $\sigma'(x) = 0$ . As  $\sigma'$  is additive function, by application of Lemma 2.1, we find either  $x \in Z(R)$  for all  $x \in R$  or  $\sigma'(x) = 0$  for all  $x \in R$ . Since  $R$  is noncommutative, we have  $\sigma' = 0$ . Thus,  $T(x) = qx^*$  for all  $x \in R$ . □

*Proof of Theorem 4.1.* Let  $T(x) = (F - \mathfrak{d})(x)$  for all  $x \in R$ . Then for any  $x \in R$ ,  $T(x^2) = (F - \mathfrak{d})(x^2) = F(x)x^* + x\mathfrak{d}(x) + c_1 - \mathfrak{d}(x)x^* - x\mathfrak{d}(x) - c_2$  where  $c_1$  and  $c_2$  are corresponding central element. It turns out to be  $T(x^2) - T(x)x^* = c_1 + c_2 \in Z(R)$  for all  $x \in R$ . Therefore  $T$  is a centrally-extended Jordan \*-left centralizer. Invoking Proposition 4.4, we get  $T(x) = qx^*$  for all  $x \in R$ , where  $q \in Q_s(R)$ . Thus  $F(x) = qx^* + \mathfrak{d}(x)$  for all  $x \in R$ . □

### 5. CENTRALIZING CONDITIONS

An astonishing result of Posner [27] states that a prime ring  $R$  is commutative if it possesses a non-zero derivation  $d$  which is centralizing on  $R$  (i.e.  $[d(x), x] \in Z(R)$ ). Proceeding this investigation, Mayne [24, 25] studied automorphisms and derivations which are centralizing on appropriate subsets of a prime ring. Later on, Bell and Martindale [8] examined centralizing mappings of semiprime rings. Since then several authors have extended these results in different directions. In 2014, Ali and Dar [3] introduced the notion of \*-centralizing mappings in prime rings with involution. Motivated by these studies, in this section, we are intended to describe the structure of centralizing and \*-centralizing centrally-extended generalized Jordan \*-derivations of a prime ring with involution  $*$ .

**Theorem 5.1.** *Let  $R$  be a 2-torsion free prime ring with involution  $'*$  and  $F : R \rightarrow R$  a centrally-extended generalized Jordan \*-derivation associated with centrally-extended Jordan \*-derivation  $\mathfrak{d}$ . If  $[F(x), x] \in Z(R)$  for all  $x \in R$ , then there exists  $\lambda \in C$  such that  $F(x) = \lambda x$  for all  $x \in R$ , unless  $R$  satisfies  $s_4$ .*

*Proof.* Suppose that  $R$  does not satisfies  $s_4$ . By hypothesis, we have

$$[F(x), x] \in Z(R), \quad \text{for all } x \in R.$$

If  $Z(R) = (0)$ , then from Lemma 2.11, we find that there exists  $a \in Q_s(R)$  such that

$$(5.1) \quad F(x) = ax^* + \mathfrak{d}(x), \quad \text{for all } x \in R.$$

Since  $F$  is a additive and centralizing map, by using Lemma 2.9 and Lemma 2.10, there exists  $\lambda \in C$  and a map  $\sigma : R \rightarrow C$  such that

$$(5.2) \quad F(x) = \lambda x + \sigma(x), \quad \text{for all } x \in R.$$

In view of Proposition 4.2, it follows from equations (5.1) and (5.2) that

$$(5.3) \quad \lambda x + \sigma(x) = ax^* + \mathfrak{d}(x), \quad \text{for all } x \in A.$$

By Lemma 2.16 and Proposition 4.1, there exists  $b \in Q_s(R)$  such that  $\mathfrak{d}(x) = xb - xb^*$  for all  $x \in A$ . Therefore, from (5.3), we have

$$(5.4) \quad \lambda x + \sigma(x) = ax^* + xb - bx^*, \quad \text{for all } x \in A.$$

Taking 1 instead of  $x$  in (5.4), we find

$$(5.5) \quad a \in C.$$

Replacing  $x$  by  $h$  in (5.4), we obtain

$$(5.6) \quad \lambda h + \sigma(h) = ah + d(h), \quad \text{for all } h \in H(A),$$

where  $d(x) = [x, b]$ . Since  $a \in C$ , from (5.6), we see that  $[d(h), h] = 0$  for all  $h \in H(A)$ . If  $d \neq 0$ , then Lemma 2.12 leads us to a contradiction.

On the other hand, let  $d = 0$ , i.e.,  $b \in C$ . Moreover, from (5.6), we get  $(\lambda - a)h \in C$ . Since  $\lambda - a \in C$ , in view of Lemma 2.3, we have either  $\lambda = a$  or  $H(R) \subseteq Z(R)$ . In the latter case, we have a contradiction by Lemma 2.13, so we left with  $\lambda = a$ . With this, from (5.4) we obtain

$$(5.7) \quad 2\lambda k - 2bk \in C, \quad \text{for all } k \in S(R).$$

As  $\lambda, b \in C$ , it implies either  $S(R) \subseteq Z(R)$  or  $\lambda = b$ . In the former case, we have the desired result by Lemma 2.18 and in the latter case, (5.1) yields  $F(x) = \lambda x^* + x\lambda - \lambda x^*$  for all  $x \in R$ , i.e.,  $F(x) = \lambda x$  for all  $x \in R$ , as desired.

In case  $Z(R) \neq \{0\}$ , we have  $0 \neq h_c \in Z(R)$  such that  $h_c^* = h_c$ . By the assumption  $[F(x), x] \in Z(R)$  for all  $x \in R$ . With the aid of Lemma 2.9 and Proposition 4.3, we have

$$(5.8) \quad [F(x), x] = 0, \quad \text{for all } x \in R.$$

From Lemma 2.10 and (5.8), we have

$$(5.9) \quad F(x) = \lambda x + \sigma(x), \quad \text{for all } x \in R,$$

for some  $\lambda \in C$  and a map  $\sigma : R \rightarrow C$ . Application of Proposition 4.1 in (5.9) yields

$$(5.10) \quad qx^* + \mathfrak{d}(x) = \lambda x + \sigma(x), \quad \text{for all } x \in R,$$

for some  $q \in Q_s(R)$ . From (B), we find

$$(5.11) \quad F(x \circ h_c) - F(x)h_c - F(h_c)x^* - x\mathfrak{d}(h_c) - h_c\mathfrak{d}(x) \in Z(R), \quad \text{for all } x \in R.$$

Using (5.9) in (5.11), we find

$$\lambda(x \circ h_c) - \lambda x h_c - \lambda h_c x^* - \sigma(h_c)x^* - x\mathfrak{d}(h_c) - h_c\mathfrak{d}(x) \in C, \quad \text{for all } x \in R.$$

It implies

$$(5.12) \quad \lambda(x - x^*)h_c - \sigma(h_c)x^* - x\mathfrak{d}(h_c) - h_c\mathfrak{d}(x) \in C, \quad \text{for all } x \in R.$$

Replacing  $x$  by  $h_c$  in (5.12), we conclude

$$(5.13) \quad \mathfrak{d}(h_c) \in Z(R).$$

Replacing  $x$  by  $h$  in (5.12), where  $h \in H(R)$ , we obtain

$$(5.14) \quad -\sigma(h_c)h - h\mathfrak{d}(h_c) - h_c\mathfrak{d}(h) \in C.$$

It implies

$$(5.15) \quad \mathfrak{d}(h_c)[h, x] + \sigma(h_c)[h, x] + h_c[\mathfrak{d}(h), x] = 0, \quad \text{for all } h \in H(R), x \in R.$$

Replacing  $h$  by  $h^2$  in (5.15), we find

$$(5.16) \quad (\mathfrak{d}(h_c) + \sigma(h_c))([h, x]h + h[h, x]) + h_c[\mathfrak{d}(h)h + h\mathfrak{d}(h), x] = 0, \quad \text{for all } h \in H(R), x \in R.$$

Using (5.13) and (5.15) in (5.16), we obtain

$$h_c(\mathfrak{d}(h)[h, x] + [h, x]\mathfrak{d}(h)) = 0, \quad \text{for all } h \in H(R), x \in R.$$

It implies that

$$(5.17) \quad \mathfrak{d}(h)[h, x] + [h, x]\mathfrak{d}(h) = 0, \quad \text{for all } h \in H(R), x \in R.$$

Polarizing  $h$  in (5.17), we find

$$(5.18) \quad \mathfrak{d}(h_1)[h, x] + \mathfrak{d}(h)[h_1, x] + [h, x]\mathfrak{d}(h_1) + [h_1, x]\mathfrak{d}(h) = 0, \quad \text{for all } h, h_1 \in H(R), x \in R.$$

In particular, replacing  $h_1$  by  $h_c$  in (5.18) and thereby using (5.13), we conclude

$$(5.19) \quad 2\mathfrak{d}(h_c)[h, x] = 0, \quad \text{for all } h \in H(R), x \in R.$$

If  $\mathfrak{d}(h_c) \neq 0$ , then Lemma 2.2 in (5.19) implies  $H(R) \subseteq Z(R)$ . With the aid of Lemma 2.13, we arrive at a contradiction.

In case  $\mathfrak{d}(h_c) = 0$ , we obtain from (5.12) that

$$2\lambda kh_c + \sigma(h_c)k - h_c\mathfrak{d}(k) \in C, \quad \text{for all } k \in S(R).$$

It implies  $h_c[\mathfrak{d}(k), k] = 0$ , for all  $k \in S(R)$ . From the fact  $h_c \neq 0$  and Lemma 2.2, it follows that

$$(5.20) \quad [\mathfrak{d}(k), k] = 0, \quad \text{for all } k \in S(R).$$

By using  $\mathfrak{d}(h_c) = 0$  and replacing  $h$  by  $k^2$  in (5.14), where  $k \in S(R)$ , we have

$$-\sigma(h_c)k^2 - h_c\mathfrak{d}(k^2) \in C.$$

It implies

$$-\sigma(h_c)k^2 + h_c[\mathfrak{d}(k), k] \in C, \quad \text{for all } k \in S(R).$$

From (5.20), we find

$$(5.21) \quad \sigma(h_c)k^2 \in C, \quad \text{for all } k \in S(R).$$

If  $\sigma(h_c) \neq 0$ , then Lemma 2.3 in (5.21) implies  $k^2 \in Z(R)$  for all  $k \in S(R)$ . For any fixed  $x \in R$ , we have  $[k, x]k + k[k, x] = d(k)k + kd(k) = 0$  for all  $k \in S(R)$ , where  $d(y) = [y, x]$  for all  $y \in R$ . If  $d \neq 0$ , then Lemma 2.15 yields a contradiction. Now, if  $d = 0$ , then we have  $[x, y] = 0$  for all  $x, y \in R$ , hence  $R$  is commutative, which is a contradiction.

In case  $\sigma(h_c) = 0$ , using fact  $\mathfrak{d}(h_c) = 0$ , from (5.14), we have  $h_c\mathfrak{d}(h) \in Z(R)$  for all  $h \in H(R)$ . Since  $h_c \neq 0$ , we have  $\mathfrak{d}(h) \in Z(R)$  for all  $h \in H(R)$  by Lemma 2.2. For any fixed  $h \in H(R)$ , using Lemma 2.2 in (5.17) we find that for each  $h \in H(R)$ , either  $\mathfrak{d}(h) = 0$  or  $h \in Z(R)$ . Using Lemma 2.1, we have either  $\mathfrak{d}(h) = 0$  for all  $h \in H(R)$  or  $H(R) \subseteq Z(R)$ . In latter case, we have a contradiction by Lemma 2.13. In case  $\mathfrak{d}(h) = 0$  for all  $h \in H(R)$ , we have

$$(5.22) \quad F(h^2) - F(h)h \in C, \quad \text{for all } h \in H(R).$$

Using (5.9) in (5.22), we conclude

$$(5.23) \quad \sigma(h)h \in C, \quad \text{for all } h \in H(R).$$

For any fixed  $h \in H(R)$ , using Lemma 2.3 in (5.23), we have either  $\sigma(h) = 0$  or  $h \in Z(R)$ . Aid of Lemma 2.1, we have either  $\sigma(h) = 0$  for all  $h \in H(R)$  or  $H(R) \subseteq Z(R)$ . In the latter case, we have the desired result by using Lemma 2.13. In case  $\sigma(h) = 0$ , replacing  $x$  by  $h$ , where  $h \in H(R)$  in (5.10), we get

$$(5.24) \quad qh = \lambda h \text{ for all } h \in H(R).$$

Replacing  $h$  by  $h_c$  in (5.24) and thereby using Lemma 2.2, we conclude  $q = \lambda$ . Replacing  $x$  by  $k$  in (5.10), where  $k \in S(R)$ , we find

$$(5.25) \quad \mathfrak{d}(k) = 2\lambda k + \sigma(k), \quad \text{for all } k \in S(R).$$

Using (B), we have

$$(5.26) \quad F(h \circ k) + F(h)k - F(k)h - h\mathfrak{d}(k) \in Z(R), \quad \text{for all } k \in S(R), h \in H(R).$$

Using (5.9) and (5.25) in (5.26), we conclude

$$\lambda(h \circ k) + \lambda hk - \lambda kh - \sigma(k)h - h(2\lambda k + \sigma(k)) \in C, \quad \text{for all } k \in S(R), h \in H(R).$$

It implies

$$(5.27) \quad \sigma(k)h \in C, \quad \text{for all } k \in S(R), h \in H(R).$$

If there exists  $k \in S(R)$  such that  $\sigma(k) \neq 0$ , then using Lemma 2.3 in (5.27), we find  $H(R) \subseteq Z(R)$ . With the aid of Lemma 2.13, we get a contradiction. In case  $\sigma(k) = 0$  for all  $k \in S(R)$ . From (5.9) and fact  $\sigma(k) = \sigma(h) = 0$ , we find  $F(x) = F(h + k) = F(h) + F(k) = \lambda h + \lambda k = \lambda(h + k) = \lambda x$  for all  $x \in R$ .  $\square$

**Theorem 5.2.** *Let  $R$  be a 2-torsion free prime ring with involution  $'*$ ' and  $F : R \rightarrow R$  a centrally-extended generalized Jordan  $*$ -derivation with an associated centrally-extended Jordan  $*$ -derivation  $\mathfrak{d}$ . If  $[F(x), x^*] \in Z(R)$  for all  $x \in R$ , then there exists  $\lambda \in C$  such that  $F(x) = \lambda x^*$  for all  $x \in R$ , unless  $R$  satisfies  $s_4$ .*

*Proof.* Suppose that  $R$  does not satisfies  $s_4$ . If  $Z(R) = (0)$ , then by Lemma 2.11, we find

$$(5.28) \quad F(x) = ax^* + \mathfrak{d}(x), \quad \text{for all } x \in R,$$

for some  $a \in Q_s(R)$ . From Lemma 2.5 and assumption, we find  $[F(x), x^*] = 0$  for all  $x \in R$ . Applying involution, we obtain  $[F(x)^*, x] = 0$  for all  $x \in R$ . By Lemma 2.10, we have

$$F(x)^* = \lambda x + \sigma(x), \quad \text{for all } x \in R.$$

It implies

$$(5.29) \quad F(x) = \lambda^* x^* + \sigma(x)^*, \quad \text{for all } x \in R.$$

By Proposition 4.2, Eq. (5.28) and (5.29), we conclude

$$(5.30) \quad \lambda^* x^* + \sigma(x)^* = ax^* + \mathfrak{d}(x), \quad \text{for all } x \in A.$$

Replacing  $x$  by 1 in (5.30), we obtain

$$(5.31) \quad a \in C.$$

Using (5.31) in (5.30), we obtain

$$[\mathfrak{d}(x), x^*] = 0, \quad \text{for all } x \in R.$$

Application of Lemma 2.8 implies  $\mathfrak{d} = 0$  or  $R$  satisfy  $s_4$  identity. For non trivial solution, we have  $\mathfrak{d} = 0$ . Thus using it in (5.28), we obtain  $F(x) = ax^*$  for all  $x \in R$ , where  $a \in C$  as desired.

Let  $Z(R) \neq \{0\}$ . Then there exists  $0 \neq h_c \in Z(R)$  such that  $h_c^* = h_c$ . By the assumption, we have  $[F(x), x^*] \in Z(R)$  for all  $x \in R$ . With the aid of Lemma 2.5 and Lemma 4.3, we have

$$(5.32) \quad [F(x), x^*] = 0, \quad \text{for all } x \in R.$$

Applying involution both sides in (5.32), we find

$$(5.33) \quad [F(x)^*, x] = 0, \quad \text{for all } x \in R.$$

Application of Lemma 2.10 in (5.33) implies

$$(5.34) \quad F(x)^* = \lambda x + \sigma(x), \quad \text{for all } x \in R,$$

for some  $\lambda \in C$  and a mapping  $\sigma : R \rightarrow C$ . Applying involution both sides in (5.34), we get

$$(5.35) \quad F(x) = \lambda^* x^* + \sigma(x)^*, \quad \text{for all } x \in R.$$

From (B), we have

$$(5.36) \quad F(x \circ y) - F(x)y^* - F(y)x^* - x\mathfrak{d}(y) - y\mathfrak{d}(x) \in Z(R), \quad \text{for all } x, y \in R.$$

Using (5.35) in (5.36), we find

$$\lambda^*(x \circ y)^* - \lambda^*(x)^*y^* - \sigma(x)^*y^* - \lambda^*y^*x^* - \sigma(y)^*x^* - x\mathfrak{d}(y) - y\mathfrak{d}(x) \in C, .$$

for all  $x, y \in R$ . It implies

$$(5.37) \quad -\sigma(x)^*y^* - \sigma(y)^*x^* - x\mathfrak{d}(y) - y\mathfrak{d}(x) \in C, \quad \text{for all } x, y \in R.$$

Replacing  $x$  and  $y$  by  $h_c$  in (5.37), we find  $h_c\mathfrak{d}(h_c) \in Z(R)$ . It forces

$$(5.38) \quad \mathfrak{d}(h_c) \in Z(R).$$

Replacing  $y$  by  $h_c$  in (5.37) and using (5.38), we find

$$(5.39) \quad -\sigma(h_c)^*x^* - x\mathfrak{d}(h_c) - h_c\mathfrak{d}(x) \in Z(R), \quad \text{for all } x \in R.$$

Replacing  $x$  by  $h$  in (5.39), where  $h \in H(R)$ , we have

$$-\sigma(h_c)^*h - h\mathfrak{d}(h_c) - h_c\mathfrak{d}(h) \in C.$$

Commuting with  $x$  and using (5.38), we obtain

$$(5.40) \quad \mathfrak{d}(h_c)[h, x] + \sigma(h_c)^*[h, x] + h_c[\mathfrak{d}(h), x] = 0, \quad \text{for all } x \in R, h \in H(R).$$

Replacing  $h$  by  $h^2$  in (5.40) and using it, we conclude

$$(5.41) \quad \mathfrak{d}(h)[h, x] + [h, x]\mathfrak{d}(h) = 0.$$

Polarizing (5.41), we find

$$(5.42) \quad \mathfrak{d}(h_1)[h, x] + \mathfrak{d}(h)[h_1, x] + [h, x]\mathfrak{d}(h_1) + [h_1, x]\mathfrak{d}(h) = 0, \quad \text{for all } h, h_1 \in H(R).$$

In particular, replacing  $h_1$  by  $h_c$  in (5.42) to obtain

$$(5.43) \quad 2\mathfrak{d}(h_c)[h, x] = 0, \quad \text{for all } h \in H(R).$$

If  $\mathfrak{d}(h_c) \neq 0$ , then using Lemma 2.2 and (5.38) in (5.43), we find  $H(R) \subseteq Z(R)$ . With the aid of Lemma 2.13, we have contradiction.

In case  $\mathfrak{d}(h_c) = 0$ , from (5.39), we find  $[\mathfrak{d}(x), x^*] = 0$  for all  $x \in R$ . For non trivial solution, Lemma 2.8 yields  $\mathfrak{d} = 0$ . Using it in (5.37), we get

$$\sigma(y)^*[x^*, y^*] = 0, \quad \text{for all } x, y \in R.$$

For any fixed  $y \in R$ , we have either  $\sigma(y) = 0$  or  $[y^*, R] = 0$ . Application of Lemma 2.1 implies that either  $\sigma = 0$  or  $[y, R] = (0)$  for all  $y \in R$ . But  $R$  is noncommutative, so we left with  $\sigma = 0$ . Thus, from (5.35) we have  $F(x) = ax^*$ , where  $a = \lambda^* \in C$ . It completes the proof.  $\square$

## 6. HYPERCOMMUTING CONDITIONS

A pair of mappings  $f$  and  $g$  satisfying the condition  $f(x)x - xg(x) = 0$  (resp.  $f(x)x^* - x^*g(x) = 0$ ) on an appropriate subset  $K$  of a ring (resp. ring with involution)  $R$  is called hypercommuting (resp.  $*$ -hypercommuting). Obviously, it is a more general concept than that of commuting ( $*$ -commuting) mappings. In this section, we study a pair  $(F, G)$  of centrally-extended generalized Jordan  $*$ -derivations which is hypercommuting or  $*$ -hypercommuting.

**Theorem 6.1.** *Let  $R$  be a 2-torsion free prime ring with involution  $'*$ ' and  $F, G : R \rightarrow R$  are centrally-extended generalized Jordan  $*$ -derivations with an associated centrally-extended Jordan  $*$ -derivations  $\mathfrak{d}, \mathfrak{g}$ , respectively. If  $F(x)x - xG(x) = 0$  for all  $x \in R$ , then there exists  $\lambda \in C$  such that  $F(x) = G(x) = \lambda x$  for all  $x \in R$ , unless  $R$  satisfies  $s_4$ .*

*Proof.* Assume that  $R$  does not satisfy  $s_4$ . By the hypothesis, we have

$$F(x)x - xG(x) = 0, \quad \text{for all } x \in R.$$

Suppose that  $Z(R) = (0)$ . Clearly in this case  $F$  and  $G$  becomes generalized Jordan  $*$ -derivations. Thus with the aid of Lemma 2.11, we have  $F(x) = ax^* + \mathfrak{d}(x)$ ,  $G(x) = bx^* + \mathfrak{g}(x)$  for all  $x \in R$ , where  $a, b \in Q_s(R)$ . Using it in our hypothesis, we find

$$(ax^* + \mathfrak{d}(x))x - x(bx^* + \mathfrak{g}(x)) = 0, \quad \text{for all } x \in R.$$

The fact of Lemma 2.16 yields  $\mathfrak{d}(x) = xc - cx^*$ ,  $\mathfrak{g}(x) = xd - dx^*$  for all  $x \in R$  for some  $c, d \in Q_s(R)$ . In this view it follows that  $R$  satisfies the functional identity

$$(6.1) \quad (ax^* + xc - cx^*)x = x(bx^* + xd - dx^*), \quad \text{for all } x \in R.$$

Application of Lemma 2.6 in (6.1) yields

$$(6.2) \quad (ax^* + xc - cx^*)x = x(bx^* + xd - dx^*), \quad \text{for all } x \in A.$$

Replacing  $x$  by 1 in (6.2), we find  $a = b$ . Polarizing (6.2), we obtain  $(ax^* + xc - cx^*)y + (ay^* + yc - cy^*)x = y(bx^* + xd - dx^*) + x(by^* + yd - dy^*)$  for all  $x, y \in A$ . Replacing  $y$  by 1, we get  $ax^* + xc - cx^* + ax = bx^* + xd - dx^* + xb$  for all  $x, y \in A$ . By using the fact  $a = b$  in preceding equation, we find

$$(6.3) \quad xc - cx^* - xd + dx^* + [a, x] = 0, \quad \text{for all } x \in A.$$

Replacing  $x$  by  $h$  in (6.3), where  $h \in H(R)$ , to obtain

$$(6.4) \quad -[c, h] + [d, h] + [a, h] = 0.$$

It implies  $[d - c + a, h] = 0$ . With the aid of Lemma 2.17, we have  $d - c + a \in C$ . Replacing  $x$  by  $k$  in (6.3), where  $k \in S(R)$  and thereby using the fact  $d - c + a \in C$ , we find  $k \circ (c - d) + [c - d, k] = 0$  for all  $k \in S(R)$ . It implies

$$(6.5) \quad (c - d)k = 0, \quad \text{for all } k \in S(R).$$

Replacing  $k$  by  $h \circ k$  in (6.5), where  $h \in H(R)$ ,  $k \in S(R)$  and using (6.5), we find

$$(6.6) \quad (c - d)hk = 0, \quad \text{for all } h \in H(R), k \in S(R).$$

From (6.5) and (6.6), we get

$$(6.7) \quad (c - d)[k, h] = 0, \quad \text{for all } k \in S(R), h \in H(R).$$

Replacing  $k$  by  $k \circ h_1$  in (6.7), we obtain

$$(6.8) \quad (c - d)h_1[k, h] + (c - d)[k, h]h_1 + (c - d)k[h, h_1] + (c - d)[h, h_1]k = 0.$$

Using (6.5) and (6.7) in (6.8), we find

$$(6.9) \quad (c-d)h_1[k, h] = 0, \quad \text{for all } k \in S(R), h, h_1 \in H(R).$$

Equation (6.5) also implies

$$(6.10) \quad (c-d)k_1[k, h] = 0, \quad \text{for all } k, k_1 \in S(R), h \in H(R).$$

From (6.9) and (6.10), we have

$$\begin{aligned} 2(c-d)x[k, h] &= (c-d)(2x)[k, h] \\ &= (c-d)(h_1 + k_1)[k, h] \\ &= (c-d)h_1[k, h] + (c-d)k_1[k, h] \\ &= 0, \quad \text{for all } x \in R, h \in H(R), k \in S(R). \end{aligned}$$

Thus, we have  $(c-d)R[k, h] = (0)$  for all  $k \in S(R)$  and  $h \in H(R)$ . In case  $c \neq d$ , primeness of  $R$  implies  $[H(R), S(R)] = (0)$ . Lemma 2.18 leads us to the contradiction. In case  $c = d$ , we find  $\mathfrak{d} = \mathfrak{g}$ . Using the fact  $a = b$ , we conclude  $F(x) = G(x)$  for all  $x \in R$ . Thus, we have the desired result by Theorem 5.1.

Let  $Z(R) \neq (0)$ . Then there exists  $0 \neq h_c \in Z(R)$  such that  $h_c^* = h_c$ . By the assumption, we have

$$(6.11) \quad F(x)x - xG(x) = 0, \quad \text{for all } x \in R.$$

Replacing  $x$  by  $h_c$  in (6.11), we find

$$(6.12) \quad F(h_c) - G(h_c) = 0.$$

Proposition 4.1 yields

$$(6.13) \quad F(x) = q_1x^* + \mathfrak{d}(x), \quad \text{for all } x \in R,$$

$$(6.14) \quad G(x) = q_2x^* + \mathfrak{g}(x), \quad \text{for all } x \in R,$$

for some  $q_1, q_2 \in Q_s(R)$ . Using (6.13) and (6.14) in (6.12), we obtain

$$(6.15) \quad (q_1 - q_2)(h_c) + (\mathfrak{d} - \mathfrak{g})(h_c) = 0.$$

Polarizing (6.11) to obtain

$$(6.16) \quad F(x)y - yG(x) + F(y)x - xG(y) = 0, \quad \text{for all } x, y \in R.$$

Replacing  $y$  by  $h_c$  in (6.16) to obtain

$$(6.17) \quad (F(x) - G(x))h_c + F(h_c)x - xG(h_c) = 0, \quad \text{for all } x \in R.$$

Using (6.12) in (6.17), we have

$$(6.18) \quad (F(x) - G(x))h_c + [F(h_c), x] = 0, \quad \text{for all } x \in R.$$

Replacing  $h_c$  by  $h_c^2$  in (6.18) and using Lemma 2.2, we conclude

$$(6.19) \quad [\mathfrak{d}(h_c), x] = 0, \quad \text{for all } x \in R.$$

It implies  $\mathfrak{d}(h_c) \in Z(R)$ . In the same way, we compute  $\mathfrak{g}(h_c) \in Z(R)$ . Using (6.13) and (6.19) in (6.18), we obtain

$$(F(x) - G(x) + [q_1, x])h_c = 0, \quad \text{for all } x \in R.$$

It implies

$$(6.20) \quad F(x) - G(x) + [q_1, x] = 0, \quad \text{for all } x \in R.$$

Since  $F$  and  $G$  are centrally-extended generalized Jordan \*-derivation, replacing  $x$  by  $h \circ h_c$  in (6.20) to obtain

$$(6.21) \quad \begin{aligned} &(F(h) - G(h))h_c + (F(h_c) - G(h_c))h + h_c(\mathfrak{d}(h) - \mathfrak{g}(h)) \\ &+ h(\mathfrak{d}(h_c) - \mathfrak{g}(h_c)) + 2[q, h]h_c \in Z(R), \quad \text{for all } h \in H(R). \end{aligned}$$

Using (6.12) and (6.20) in (6.21), we see that

$$(6.22) \quad (\mathfrak{d}(h) - \mathfrak{g}(h))h_c + [q, h]h_c + h(\mathfrak{d}(h_c) - \mathfrak{g}(h_c)) \in Z(R), \quad \text{for all } h \in H(R).$$

Replacing  $h$  by  $h^2$  in (6.22), we obtain

$$(6.23) \quad \begin{aligned} &(\mathfrak{d}(h) - \mathfrak{g}(h))hh_c + h(\mathfrak{d}(h) - \mathfrak{g}(h))h_c + [q_1, h]hh_c + h[q_1, h]h_c \\ &+ h^2(\mathfrak{d}(h_c) - \mathfrak{g}(h_c)) \in Z(R), \quad \text{for all } h \in H(R). \end{aligned}$$

From (6.22) and (6.23), we conclude

$$(6.24) \quad h^2(\mathfrak{d}(h_c) - \mathfrak{g}(h_c)) \in Z(R).$$

If  $\mathfrak{d}(h_c) - \mathfrak{g}(h_c) \neq 0$ , then using Lemma 2.3 in (6.24), we have  $h^2 \in Z(R)$  for all  $h \in H(R)$ . Thus, we have the result by Lemma 2.13. If  $\mathfrak{d}(h_c) - \mathfrak{g}(h_c) = 0$ , then from (6.15), we obtain

$$(6.25) \quad (q_1 - q_2)h_c = 0.$$

Using Lemma 2.2 in (6.25), we have  $q_1 = q_2$ . With the aid of (6.13), (6.14) and the fact  $q_1 = q_2$  in (6.20), we find

$$(6.26) \quad \mathfrak{d}(x) - \mathfrak{g}(x) + [q_1, x] = 0, \quad \text{for all } x \in R.$$

Replacing  $x$  by  $h \circ k$  where  $h \in H(R)$ ,  $k \in S(R)$  in (6.26), we obtain

$$(6.27) \quad \begin{aligned} &(\mathfrak{d}(h) - \mathfrak{g}(h))(-k) + k(\mathfrak{d}(h) - \mathfrak{g}(h)) + (\mathfrak{d}(k) - \mathfrak{g}(k))h + h(\mathfrak{d}(k) - \mathfrak{g}(k)) \\ &+ [q_1, h]k + k[q_1, h] + [q_1, k]h + h[q_1, k] \in Z(R). \end{aligned}$$

Using (6.26) in (6.27) to conclude

$$(6.28) \quad 2[q_1, h]k \in Z(R), \quad \text{for all } h \in H(R), k \in S(R).$$

We now split the proof into the following two parts.

**Case 1.** If mapping induced on centroid is non identity map, then there exists  $0 \neq z \in C$  such that  $z^* \neq z$ . Replacing  $h$  by  $x + x^*$  and  $k$  by  $y - y^*$  in (6.28), where  $x, y \in R$  in order to obtain

$$(6.29) \quad 2[q_1, x + x^*](y - y^*) \in Z(R), \quad \text{for all } x, y \in R.$$

With the aid of Lemma 2.6 in (6.29), we have

$$(6.30) \quad [q_1, x + x^*](y - y^*) \in Z(R), \quad \text{for all } x, y \in Q_s(R).$$

Replace  $y$  by  $z$  in (6.30), we have

$$(6.31) \quad [q_1, x + x^*](z - z^*) \in C, \quad \text{for all } x \in Q_s(R).$$

Using Lemma 2.3 in (6.31), we have

$$(6.32) \quad [q_1, x + x^*] \in Z(R), \quad \text{for all } x \in Q_s(R).$$

Replacing  $x$  by  $h$ , where  $h \in H(R)$  in (6.32), we obtain  $[q_1, h] \in Z(R)$ . Using Lemma 2.17, we conclude  $q_1 \in Z(R)$ . From (6.20), we have  $F(x) = G(x)$  for all  $x \in R$ . Hence, by Theorem 5.1, we get the desired result.

**Case 2.** If mapping induced on centroid is an identity map, then  $c^* = c$  for all  $c \in C$ . From (6.28), we have

$$([q_1, h]k)^* = [q_1, h]k, \quad \text{for all } h \in H(R), k \in S(R).$$

It implies

$$(6.33) \quad [q_1, h]k - k[q_1^*, h] = 0, \quad \text{for all } h \in H(R), k \in S(R).$$

Replacing  $k$  by  $k \circ h_1$  in (6.33), where  $k \in S(R)$ ,  $h_1 \in H(R)$ , we obtain

$$(6.34) \quad [q_1, h]h_1k + [q_1, h]kh_1 - kh_1[q_1^*, h] - h_1k[q_1^*, h] = 0.$$

Using (6.28) and (6.33) in (6.34) to conclude

$$(6.35) \quad [q_1, h]h_1k - kh_1[q_1^*, h] = 0, \quad \text{for all } h, h_1 \in H(R), k \in S(R).$$

Replacing  $h_1$  by  $k_1^2$  in (6.35), where  $k_1 \in S(R)$ , we find

$$([q_1, h]k_1)k_1k - kk_1(k_1[q_1^*, h]) = 0.$$

It implies

$$(6.36) \quad ([q_1, h]k_1)k_1k - kk_1([q_1, h]k_1)^* = 0, \quad \text{for all } h \in H(R), k, k_1 \in S(R).$$

Using (6.28) and (6.33) in (6.36), we obtain

$$(6.37) \quad [q_1, h]k_1[k_1, k] = 0.$$

For fixed  $k_1 \in S(R)$ , from the fact  $[q_1, h]k_1 \in C$ , using Lemma 2.2 in (6.37), we have either  $[q_1, h]k_1 = 0$  for all  $h \in H(R)$  or  $[k_1, k] = 0$  for all  $k \in S(R)$ . Invoking Lemma 2.1 yields that either  $[q_1, h]k_1 = 0$  for all  $h \in H(R)$ ,  $k_1 \in S(R)$  or  $[k_1, k] = 0$  for all  $k, k_1 \in S(R)$ . In the latter case, replace  $k_1$  by  $h \circ k$  to obtain  $[h, k]k + k[h, k] = 0$  for all  $h \in H(R)$ ,  $k \in S(R)$ . For any fixed  $h \in H(R)$ , we obtain  $d(k)k + kd(k) = 0$ , where  $d(x) = [h, x]$  for all  $k \in S(R)$ . With the aid of Lemma 2.15, we find either  $R$  satisfies  $s_4$  identity or  $d = 0$ . In case  $d = 0$ , we have  $H(R) \subseteq Z(R)$ . Application of Lemma 2.13 gives a contradiction. If  $[q_1, h]k_1 = 0$ , then using the similar arguments as in (6.5), we find  $[q_1, H(R)] = (0)$ . With the aid of Lemma 2.17, we have  $q_1 \in C$ . From (6.20), we have  $F(x) = G(x)$  for all  $x \in R$ . Thus, we get the desired conclusion from Theorem 5.1.  $\square$

**Theorem 6.2.** *Let  $R$  be a 2-torsion free prime ring with involution  $'*$  and  $F, G : R \rightarrow R$  are centrally-extended generalized Jordan  $*$ -derivations associated with centrally-extended Jordan  $*$ -derivations  $\mathfrak{d}, \mathfrak{g}$ , respectively. If  $F(x)x^* - x^*G(x) = 0$  for all  $x \in R$ , then there exists  $\lambda \in C$  such that  $F(x) = G(x) = \lambda x^*$ , unless  $R$  satisfies  $s_4$ .*

*Proof.* Suppose that  $R$  does not satisfies  $s_4$ . By the hypothesis, we have  $F(x)x^* - x^*G(x) = 0$  for all  $x \in R$ .

If  $Z(R) = (0)$ , then application of Lemma 2.11 yields  $F(x) = ax^* + \mathfrak{d}(x)$ ,  $G(x) = bx^* + \mathfrak{g}(x)$  for all  $x \in R$ , where  $a, b \in Q_s(R)$ . Using it in our hypothesis, we find

$$(6.38) \quad (ax^* + \mathfrak{d}(x))x^* - x^*(bx^* + \mathfrak{g}(x)) = 0, \quad \text{for all } x \in R.$$

With the aid of Lemma 2.16, we conclude  $\mathfrak{d}(x) = xc - cx^*$ ,  $\mathfrak{g}(x) = xd - dx^*$  for all  $x \in R$  for some  $c, d \in U(R)$ . Using it in (6.38), we find

$$(6.39) \quad (ax^* + xc - cx^*)x^* = x^*(bx^* + xd - dx^*), \quad \text{for all } x \in R.$$

Using Lemma 2.6 in (6.39), we conclude

$$(6.40) \quad (ax^* + xc - cx^*)x^* = x^*(bx^* + xd - dx^*), \quad \text{for all } x \in A.$$

Replacing  $x$  by 1 in (6.40), we get  $a = b$ . Polarizing (6.40), we obtain

$$(6.41) \quad \begin{aligned} & (ax^* + xc - cx^*)y^* + (ay^* + yc - cy^*)x^* \\ &= y^*(bx^* + xd - dx^*) + x^*(by^* + yd - dy^*), \quad \text{for all } x, y \in R. \end{aligned}$$

Replacing  $y$  by 1 in (6.41), we find

$$(6.42) \quad ax^* + xc - cx^* + ax^* = bx^* + xd - dx^* + x^*b, \quad \text{for all } x, y \in R.$$

Using the fact  $a = b$  in (6.42), we conclude

$$(6.43) \quad xc - cx^* - xd + dx^* + [a, x^*] = 0, \quad \text{for all } x \in R.$$

Replacing  $x$  by  $h$  in (6.43), where  $h \in H(R)$ , we obtain

$$(6.44) \quad -[c, h] + [d, h] + [a, h] = 0.$$

Since (6.44) is the same as (6.4), using similar arguments, we can reach our conclusion.

Let  $Z(R) \neq (0)$ . Then there exists  $0 \neq h_c \in Z(R)$  such that  $h_c^* = h_c$ . By the assumption, we have

$$(6.45) \quad F(x)x^* - x^*G(x) = 0, \quad \text{for all } x \in R.$$

Replacing  $x$  by  $h_c$  in (6.45), we obtain

$$(6.46) \quad F(h_c) - G(h_c) = 0.$$

Invoking Proposition 4.1, we have

$$(6.47) \quad F(x) = q_1x^* + \mathfrak{d}(x), \quad \text{for all } x \in R,$$

$$(6.48) \quad G(x) = q_2x^* + \mathfrak{g}(x), \quad \text{for all } x \in R,$$

for some  $q_1, q_2 \in Q_s(R)$ . Using (6.47) and (6.48) in (6.46), we obtain

$$(6.49) \quad (q_1 - q_2)(h_c) + (\mathfrak{d} - \mathfrak{g})(h_c) = 0.$$

Polarizing (6.45) to get

$$(6.50) \quad F(x)y^* - y^*G(x) + F(y)x^* - x^*G(y) = 0, \quad \text{for all } x, y \in R.$$

Replacing  $y$  by  $h_c$  in (6.50), we see that

$$(6.51) \quad (F(x) - G(x))h_c + F(h_c)x^* - x^*G(h_c) = 0, \quad \text{for all } x \in R.$$

Using (6.46) in (6.51), we have

$$(6.52) \quad (F(x) - G(x))h_c + [F(h_c), x^*] = 0, \quad \text{for all } x \in R.$$

Replacing  $h_c$  by  $h_c^2$  in (6.52), we obtain

$$(6.53) \quad [\mathfrak{d}(h_c), x^*] = 0, \quad \text{for all } x \in R.$$

It implies  $\mathfrak{d}(h_c) \in Z(R)$ . In the same way, one can easily observe that  $\mathfrak{g}(h_c) \in Z(R)$ . With the aid of (6.47), (6.53) in (6.52), we have

$$(6.54) \quad (F(x) - G(x) + [q_1, x^*])h_c = 0, \quad \text{for all } x \in R.$$

It yields

$$(6.55) \quad F(x) - G(x) + [q_1, x^*] = 0, \quad \text{for all } x \in R.$$

Replacing  $x$  by  $h \circ h_c$  in (6.55) where  $h \in H(R)$ , we obtain

$$(6.56) \quad (F(h) - G(h))h_c + (F(h_c) - G(h_c))h + h_c(\mathfrak{d}(h) - \mathfrak{g}(h)) \\ + h(\mathfrak{d}(h_c) - \mathfrak{g}(h_c)) + 2[q, h]h_c \in Z(R), \quad \text{for all } h \in H(R).$$

Using (6.46) and (6.55) in (6.56), we conclude

$$(6.57) \quad (\mathfrak{d}(h) - \mathfrak{g}(h))h_c + [q, h]h_c + h(\mathfrak{d}(h_c) - \mathfrak{g}(h_c)) \in Z(R), \quad \text{for all } h \in H(R).$$

Replacing  $h$  by  $h^2$  in (6.57), we obtain

$$(6.58) \quad (\mathfrak{d}(h) - \mathfrak{g}(h))hh_c + h(\mathfrak{d}(h) - \mathfrak{g}(h))h_c + [q_1, h]hh_c + h[q_1, h]h_c \\ + h^2(\mathfrak{d}(h_c) - \mathfrak{g}(h_c)) \in Z(R), \quad \text{for all } h \in H(R).$$

From (6.57) and (6.58), we conclude

$$(6.59) \quad h^2(\mathfrak{d}(h_c) - \mathfrak{g}(h_c)) \in Z(R).$$

If  $\mathfrak{d}(h_c) - \mathfrak{g}(h_c) \neq 0$ , then using Lemma 2.3 in (6.59), we have  $h^2 \in Z(R)$  for all  $h \in H(R)$ . Thus, a contradiction follows from Lemma 2.13. In case  $\mathfrak{d}(h_c) - \mathfrak{g}(h_c) = 0$ , from (6.49), we obtain

$$(q_1 - q_2)h_c = 0,$$

and it implies  $q_1 = q_2$ . Now from (6.47), (6.48) and (6.55) we find

$$(6.60) \quad \mathfrak{d}(x) - \mathfrak{g}(x) + [q_1, x^*] = 0, \quad \text{for all } x \in R.$$

Replacing  $x$  by  $h \circ k$  in (6.60), where  $h \in H(R)$ ,  $k \in S(R)$ , we find

$$(6.61) \quad (\mathfrak{d}(h) - \mathfrak{g}(h))(-k) + k(\mathfrak{d}(h) - \mathfrak{g}(h)) + (\mathfrak{d}(k) - \mathfrak{g}(k))h + h(\mathfrak{d}(k) \\ - \mathfrak{g}(k)) - [q_1, h]k - k[q_1, h] - [q_1, k]h - h[q_1, k] \in Z(R).$$

Using (6.60) in (6.61), we conclude

$$(6.62) \quad -2k[q_1, h] \in Z(R), \quad \text{for all } h \in H(R), k \in S(R).$$

As (6.62) is same as (6.28), similar arguments are taking us to the desired conclusion.  $\square$

**Acknowledgements.** The first author would like to thank the University Grants Commission, New Delhi, for the financial support (Grant No. 402996). The third named author is supported by SERB-DST MATRICES Project (Grant No. MTR/2019/000603), India.

#### REFERENCES

- [1] A. Alahmadi, H. Alhazmi, S. Ali, N. A. Dar and A. N. Khan, *Additive maps on prime and semiprime rings with involution*, Hacettepe J. Math. Stat. **49**(3) (2020), 1126–1133. <https://doi.org/10.15672/hujms.661178>
- [2] S. Ali, *On generalized \*-derivation in \*-rings*, Palestine J. Math. **1** (2012), 32–37.
- [3] S. Ali and N. A. Dar, *On \*-centralizing mappings in rings with involution*, Georgian Math. J. **21**(1) (2014), 25–28. <https://doi.org/10.1515/gmj-2014-0006>
- [4] S. Ali, N. A. Dar and J. Vukman, *Jordan left \*-centralizers of prime and semiprime rings with involution*, Beitr Algebra Geom. **54**(2) (2013), 609–624. <https://doi.org/10.1007/S13366-012-0117-3>
- [5] S. Ali, A. Fošner, M. Fošner and M. S. Khan, *On generalized Jordan triple  $(\alpha, \beta)^*$ -derivations and related maps*, Mediterr. J. Math. **10** (2013), 1657–1668. <https://doi.org/10.1007/s00009-013-0277-x>
- [6] K. I. Beidar, W. S. Martindale III and A. V. Mikhalev, *Rings with Generalized Identities*, Pure Appl. Math. **196**, Marcel Dekker Inc., New York, 1996.
- [7] H. E. Bell and M. N. Daif, *On centrally-extended maps on rings*, Beitr Algebra Geom. **8** (2016), 129–136. <https://doi.org/10.1007/s13366-015-0244-8>
- [8] H. E. Bell and W. S. Martindale III, *Centralizing mappings of semiprime rings*, Canad. Math. Bull. **30** (1987), 92–101. <https://doi.org/10.4153/CMB-1987-014-x>
- [9] B. Bhushan, G. Sandhu, S. Ali and D. Kumar, *On centrally extended Jordan derivations and related maps in rings*, Hacettepe J. Math. Stat. **52**(1) (2023), 23–35. <https://doi.org/10.15672/hujms.1008922>
- [10] M. Brešar, *Centralizing mappings and derivations in prime rings*, J. Algebra **156**(2) (1993), 385–394. <https://doi.org/10.1006/jabr.1993.1080>
- [11] M. Brešar, *Commuting traces of biadditive mappings, commutativity-preserving mappings and Lie mappings*, Tran. Amer. Math. Soc. **335**(2) (1993), 525–546. <https://doi.org/10.1090/S0002-9947-1993-1069746-X>
- [12] M. Brešar and J. Vukman, *On some additive mappings in rings with involution*, Aeq. Math. **38** (1989), 178–185. <https://doi.org/10.1007/BF01840003>
- [13] M. Brešar, *On the distance of the composition of two derivations to the generalized derivations*, Glasgow Math. J. **33** (1991), 89–93. <https://doi.org/10.1017/S0017089500008077>
- [14] M. Brešar and B. Zalar, *On the structure of Jordan \*-derivations*, Colloq. Math. **63** (1992), 163–171. <https://doi.org/10.4064/cm-63-2-163-171>
- [15] C. L. Chuang, *\*-Differential identities of prime rings with involution*, Trans. Amer. Math. Soc. **316**(1) (1989), 251–279. <https://doi.org/10.1090/s0002-9947-1989-0937242-2>
- [16] N. A. Dar and S. Ali, *On the structure of generalized Jordan \*-derivations of prime rings*, Commun. Algebra **49**(4) (2021), 1422–1430. <https://doi.org/10.1080/00927872.2020.1837148>

- [17] N. Divinsky, *On commuting automorphisms of rings*, Trans. Royal Soc. Can. Sec. III **3**(49) (1955), 19–22. <https://doi.org/>
- [18] O. H. Ezzat, *Functional equations related to higher derivations in semiprime rings*, Open Math. **19** (2021), 1359–1365. <https://doi.org/10.1515/math-2021-0123>
- [19] A. N. Khan, N. A. Dar and A. Abbasi, *A note on generalized Jordan  $*$ -derivations on prime  $*$ -rings*, Bull. Iran. Math. Soc. **47**(2) (2021), 1–12. <https://doi.org/10.1007/s41980-020-00390-w>
- [20] T. K. Lee and P. H. Lee, *Derivations centralizing symmetric or skew elements*, Bull. Inst. Math. Acad. Sin. **14**(3) (1986), 249–256.
- [21] T. K. Lee and Y. Zhou, *Jordan  $*$ -derivations of prime rings*, J. Algebra Appl. **13**(4) (2014), Article ID 1350126. <https://doi.org/10.1142/S0219498813501260>
- [22] J.-H. Lin, *Weak Jordan derivations of prime rings*, Linear Multi. Algebra **69**(8) (2021), 1422–1445. <https://doi.org/10.1080/03081087.2019.1630061>
- [23] W. S. Martindale III, *Prime rings with involution and generalized polynomial identities*, J. Algebra **22** (1972), 502–516. [https://doi.org/10.1016/0021-8693\(72\)90164-0](https://doi.org/10.1016/0021-8693(72)90164-0)
- [24] J. H. Mayne, *Centralizing automorphisms of prime rings*, Canad. Math. Bull. **19**(1) (1976), 113–115. <https://doi.org/10.4153/CMB-1976-017-1>
- [25] J. H. Mayne, *Centralizing mappings of prime rings*, Canad. Math. Bull. **27**(1) (1984), 122–126. <https://doi.org/10.4153/CMB-1984-018-2>
- [26] N. Muthana and Z. Alkhmisi, *On centrally-extended multiplicative (generalized)- $(\alpha, \beta)$ -derivations in semiprime rings*, Hacettape J. Math. Stat. **49**(2) (2020), 578–585. <https://doi.org/10.15672/hujms.568378>
- [27] E. C. Posner, *Derivations in prime rings*, Proc. Amer. Math. Soc. **8** (1957), 1093–1100. <https://doi.org/10.2307/2032686>
- [28] P. Šemrl, *On Jordan  $*$ -derivations and an application*, Colloq. Math. **59** (1990), 241–251.
- [29] M. A. Siddeeqe, N. Khan and A. A. Abdullah, *Weak Jordan  $*$ -derivations of prime rings*, J. Algebra Appl. **22**(5) (2023), Article ID 230105. <https://doi.org/10.1142/S0219498823501050>
- [30] P. Šemrl, *Quadratic functionals and Jordan  $*$ -derivations*, Stud. Math. **97** (1991), 157–165. <https://doi.org/10.4064/SM-97-3-157-165>
- [31] S. F. El-Deken and H. Nabel, *Centrally-extended generalized  $*$ -derivations on rings with involution*, Beitr Algebra Geom. **60** (2019), 217–224. <https://doi.org/10.1007/S13366-018-0415-5>
- [32] S. F. El-Deken and M. M. El-Soufi, *On centrally-extended reverse and generalized reverse derivations*, Indian J. Pure Appl. Math. **51**(3) (2020), 1165–1180. <https://doi.org/10.1007/s13226-020-0456-y>
- [33] M. S. T. El-Sayiad, N. M. Muthana and Z. S. Alkhamisi, *On rings with some kind of centrally-extended maps*, Beitr Algebra Geom. **57**(3) (2016), 579–588. <https://doi.org/10.1007/s13366-015-0274-2>
- [34] M. S. T. El-Sayiad, A. Ageeb and A. M. Khalid, *What is the action of a multiplicative centrally-extended derivation on a ring?*, Georgian Math. J. **29**(4) (2022), 607–613. <https://doi.org/10.1515/gmj-2022-2164>

<sup>1</sup>DEPARTMENT OF MATHEMATICS,  
AKAL UNIVERSITY,  
TALWANDI SABO-151302, INDIA.  
*Email address:* [bharat\\_rs18@pbi.ac.in](mailto:bharat_rs18@pbi.ac.in)  
ORCID id: <https://orcid.org/0000-0001-6127-0731>

<sup>2</sup>DEPARTMENT OF MATHEMATICS,  
PATEL MEMORIAL NATIONAL COLLEGE,  
RAJPURA-140401, INDIA  
*Email address:* [gurninder\\_rs@pbi.ac.in](mailto:gurninder_rs@pbi.ac.in)  
ORCID id: <https://orcid.org/0000-0001-8618-6325>

<sup>3</sup>DEPARTMENT OF MATHEMATICS,  
ALIGARH MUSLIM UNIVERSITY,  
ALIGARH-202002, INDIA  
*Email address:* [asmith@example.com](mailto:asmith@example.com)  
ORCID id: <https://orcid.org/0000-0001-5162-7522>

<sup>4</sup>FEDERAL UNIVERSITY OF TECHNOLOGY-PARANÁ-BRAZIL,  
<sup>5</sup>FEDERAL UNIVERSITY OF ABC,  
<sup>6</sup>UNIVERSITY OF SÃO PAULO, BRAZIL  
*Email address:* [brunoferreira@utfpr.edu.br](mailto:brunoferreira@utfpr.edu.br)  
ORCID id: <https://orcid.org/0000-0003-1621-8197>